

Evaluation of best performing microbial culture in relation to growth, yield and quality of Sponge Gourd (*Luffa cylindrica* L.)

Abstract

The present investigation the effect of different microbial culture on growth, yield and quality of Sponge Gourd (*Luffa cylindrica* L.) was undertaken at Vegetable Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (UP) during 2022. The experiment was laid out in a Randomized Block Design with 10 treatments viz., T₀ (Control), T₁ (PSB - Photosynthetic Bacteria *Rhodospirillum rubrum*) - 20 ml/sq.m), T₂ (PSB - Photosynthetic Bacteria *Rhodospirillum rubrum*) - 30 ml/sq.m), T₃ (PSB - Photosynthetic Bacteria *Rhodospirillum rubrum*) - 40 ml/sq.m), T₄ (Vermiwash-20 ml/sq.m), T₅ (Vermiwash-300 ml/sq.m), T₆ (Vermiwash-40 ml/sq.m), T₇ (Azotobacter-8ml/sq.m), T₈ (Azotobacter - 10 ml/sq.m) and T₉ (Azotobacter- 15ml/sq.m) with three replications. Results revealed that treatment T₉ (Azotobacter- 15ml/sq.m) recorded significantly highest vine length, highest number of primary branches per plant is treatment T₄ (Vermiwash-20 ml/sq.m) and number of node per plant is treatment T₄ (Vermiwash-20 ml/sq.m). Earliness parameter such as days of first flowering is minimum days is T₉ (Azotobacter- 15ml/sq.m) and node number at first male flower appears was recorded in treatment T₉ (Azotobacter- 15ml/sq.m). Node number at first female flower appears was recorded in treatment T₉ (Azotobacter- 15ml/sq.m). Days to first fruit setting after anthesis recorded in treatment Azotobacter-10ml/sq.m (T₈). Yield parameter like number of fruit per plant were recorded under T₆ (Vermiwash-40 ml/sq. m). Average weight for 10 fruits (g) was recorded highest in treatment T₉ (Azotobacter-15 ml/sq.m). Fruit length was recorded highest in treatment (Vermiwash – 40 ml/sq.m T₆). Fruit girth was recorded highest in treatment T₉ (Azotobacter-15 ml/sq.m). Fruit yield per plot (kg) was recorded highest in treatment T₆ (Vermiwash – 40 ml/sq.m). and fruit yield /ha (q) was highest in was found in Vermiwash – 40 ml/sq.m (T₆)

Keywords; Photosynthetic Bacteria, Vermiwash, Azotobacter, Sponge Gourd, Monoecious and cucurbits.

Introduction

In India, number of major and minor cucurbits is cultivated, which share about 5.6% of the total vegetable production (Rai *et al.*, 2008). Through, authentic statistical records of area, production and productivity of Sponge gourd are not

available, but in the year 2012 India ranked the second place in the production of Sponge gourd (FAO, 2010). The Sponge Gourd (*Luffa cylindrica* Linn; 2n = 26) is one of the important tropical and subtropical cucurbitaceous crop grown extensively throughout India. It has a smooth surface and is one of the popular vegetable. It is a good source of vitamin A and C and has laxative properties. Fully ripen Sponge Gourds have high amount of fibers which are used as a cleansing agents (bathing and utensils) and making shoe soles, tablemats. The sponge of the mature fruit helps the skin in increasing the blood circulation and as a relief for rheumatic and arthritis sufferers. Also the fruits are used to cure jaundice, diabetes, to purify blood and to cure skin diseases.

Luffa is a fast-growing, long-season, warm-climate vine that can climb to a height of about 5m. It tolerates a wide range of climatic and soil conditions, although excessive rainfall during flowering and fruiting period can cause damage and reduce yield (Bal *et al.*, 2004). It is a warm-season plant that prefers average monthly temperatures in the range of 18 to 24°C, with daytime highs of 30 to 35°C. The plants are sensitive to frost. Sponge gourd prefers pH of around 6 to 6.8 and requires high level of K and P. It also grows well in green house and will grow on many soil types but well drained sandy loams are preferred (Okusanya, 1978).

Hence, it is to explore the potential of sponge gourd to develop as natural fiber and contribute to the industry as well as to the export market. Sponge gourd is monoecious, flowers open in the early morning and remain open for a day. The flowers are big and bright yellow, highly attractive and are a rich source of pollen for the foraging insects, chiefly bees (Chaudhari, 2000).

To study the effect of azotobactor ,PSB and vermiwash on the growth, yield and quality of sponge gourd

Materials and Methods

The experiment was conducted in the Vegetable Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (UP) during 2022-

23. The experiment was conducted on Sponge Gourd (*Luffa cylindrica* L.) with ten treatment including control and three replication in Randomized block design. The total number of plant included in the experiment were 120 and were spaced at 1.5m x 3m.

Statistical analysis was done by using method of analysis of variance (ANOVA) for randomized block design (RBD) by Fischer and Yates (1963). Whenever 'F' test was found significant for comparing the means of two treatments, critical difference (C. D. at 5%) was worked.

Result and Discussion

Growth parameters:-

The number of primary branches per plant ranged from 8.16 to 20.91. The maximum number of primary branches per plant was recorded under treatment T₄ - Vermiwash-20ml/sq.m (20.91) followed by T₃ - PSB (Photosynthetic Bacteria *Rhodospirillum rubrum*) - 40 ml/sq.m (16.83) respectively. The treatment T₁ and T₅, were statistically at par, but they differed significantly as compared to treatment T₀ (control) which recorded minimum number of primary branches per plant (8.16). Similar results were found by **Vimala and Natarajan (2000)** biofertilizer viz., Rhizobium @ 6 kg /ha and phosphobacteria 35ml/ sq. meter showed the highest values for primary branches, of per plant.

The No. of node per plant Vermiwash - 20 ml/sq.m (T₄) recorded the maximum number of node per plant (46.00) and was on par with those receiving PSB-40 ml/sq.m and PSB(Photosynthetic Bacteria *Rhodospirillum rubrum*)-20ml/sq.m (T₃ and T₁) was significantly superior to the rest of the treatments. The minimum number of node per plant (39.08) were recorded control (T₀) was significantly inferior to the remaining treatments. Similar results were found by **Das et al., (2015)** reported that the application of Azotobacter 5 kg/ha + PSB 5 kg/ha and recorded maximum No. of node per plant (41.00) in gourd.

The vine length ranged from 490.00 cm to 619.75 cm. The maximum vine length was recorded under treatment T₉- Azotobacter-15 ml/ sq.m (619.75 cm) followed by T₅ - Vermiwash-30 ml/ sq.m (600.00 cm), T₆- Vermiwash - 40 ml/sq.m (590.25 cm), respectively. The treatment T₅ and T₆, were statistically at

par, but they differed significantly as compared to treatment T₀ (control) which recorded minimum vine length (490.00) cm. Similar results were found by **Abdel-sattar and Mohamed(2017)**,

Table 1. Evaluation of best performing microbial culture in relation to growth of Sponge Gourd .

Treatments	No. of primary branches/plant	No. of node per plant	Vine length (cm)at harvest
Control(T ₀)	8.16	39.08	490.00
PSB- 20ml/sq.m (T ₁)	15.66	45.25	550.75
PSB- 30ml/sq.m (T ₂)	18	44.75	580.25
PSB- 40ml/sq.m (T ₃)	16.83	45.58	530.75
Vermiwash- 20 ml/sq.m (T ₄)	20.91	46.00	570.25
Vermiwash-30 ml/sq.m (T ₅)	15.41	43.83	600.00
Vermiwash-40 ml/sq.m (T ₆)	14.66	42.58	590.25
Azotobacter-8ml/sq.m (T ₇)	11.91	43.41	588.00
Azotobacter-10 ml/sq.m (T ₈)	12.66	44.41	590.00
Azotobacter-15 ml/sq.m (T ₉)	14.75	42.00	619.75
F-test	S	S	S
SE(d)	2.46	1.41	2.30
C.V.	20.25	3.95	0.49
CD at 5% level	5.21	2.98	4.88

Flowering parameter:-

Days to first flowering was maximum in control (T₀) followed by the treatment (Photosynthetic Bacteria *Rhodosudomonaspalustris*)- 20ml/ sq.m (T₁) which recorded 35.16 days and 33.50 days respectively. Azotobacter-15ml/sq.m (T₉) recorded the minimum days to appearance of first flowering 27.25 days

Formatted: Font: Italic

followed by Azotobacter-10ml/sq.m (T₈) 28.25 days. Similar findings were reported by **Bonnie et al., (2000)** in Azotobacter treated tomato plants with minimum day to flowering and **Karuppiah et al., (2005)** in ridge gourd.

The Node number at which first male flower appears Azotobacter-15ml/sq.m (T₉) recorded the minimum number of node number at which first male flower appears (4.25) and was on par with those receiving Vermiwash-40 ml/sq.m and Azotobacter- 8ml/sq.m (T₆= 4.5 and T₇ = 4.91) was significantly superior to the rest of the treatments. The maximum number of node number at which first male flower appears (10.83) were recorded control (T₀) was significantly inferior to the remaining treatments. Similar result found that the **Thrive et al., (2015)** biofertilizers (Azotobacter + Azospirillum + PSB) minimum node number in bitter gourd. The Node number at which first female flower appears Azotobacter-15ml/sq.m (T₉) recorded the minimum number of node number at which first female flower appears (5.00) and was on par with those receiving Vermiwash-40 ml/sq.m, Azotobacter- 8ml/sq.m and Azotobacter- 10ml/sq.m (T₆= 5.41, T₇ = 5.66 and T₈= 5.66) was significantly superior to the rest of the treatments. The maximum number of node number at which first female flower appears (14.25) were recorded control (T₀) was significantly inferior to the remaining treatments. Similar result **Patelet et al., (2018)** recorded that an application of vermicompost and Azotobacter, PSB each 5 kg ha⁻¹ to the bottle gourd crop found to minimum No. of node at first female flower.

The day to first fruit setting Azotobacter-10ml/sq.m (T₈) recorded the minimum number of day to first fruit setting (2.91) and was on par with those receiving PSB (Photosynthetic Bacteria *Rhodosudomonaspalustris*) – 20ml/sq.m (T₁=3.00), PSB (Photosynthetic Bacteria *Rhodosudomonaspalustris*) – 30ml/sq.m (T₂=3.00), PSB (Photosynthetic Bacteria *Rhodosudomonaspalustris*) – 40ml/sq.m (T₃=3.00), Vermiwash-30 ml/sq.m (T₄=3.00), Vermiwash-40 ml/sq.m (T₆=3.00), Azotobacter- 15ml/sq.m (T₉=3.00) was significantly superior to the rest of the treatments. The maximum number of day to first fruit setting (4.00) were recorded control (T₀) was significantly inferior to the remaining treatments. Similar result found that the **Sreenivas et al., (2000)** recorded

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

significantly days to first fruit setting with the application of Vermicompost in ridge gourd.

Table 2. Evaluation of best performing microbial culture in relation to flowering ofSponge Gourd.

Treatments	Days of first flowering	Node number at first male flower appears	Node number at first female flower appears	Days to first fruit setting after anthesis
Control(T ₀)	35.16	10.83	14.25	4.00
PSB- 20ml/sq.m (T ₁)	33.50	8.08	12	3.00
PSB- 30ml/sq.m (T ₂)	32.75	5.58	11.66	3.00
PSB- 40ml/sq.m (T ₃)	30.87	7.08	13.16	3.00
Vermiwash-20 ml/sq.m (T ₄)	31.00	5.75	7.66	3.08
Vermiwash-30 ml/sq.m(T ₅)	31.25	5.00	6.66	3.00
Vermiwash-40 ml/sq.m (T ₆)	29.62	4.50	5.41	3.00
Azotobacter-8ml/sq.m (T ₇)	28.50	4.91	5.66	3.16
Azotobacter-10 ml/sq.m (T ₈)	28.25	5.00	5.66	2.91
Azotobacter-15 ml/sq.m (T ₉)	27.25	4.25	5	3.00
F-test	S	S	S	S
SE(d)	0.69	1.26	1.01	0.22
C.V.	2.77	25.41	14.24	8.93
CD at 5% level	1.47	2.68	2.14	0.48

Yield parameter:-

The maximum number of fruit per plant (45.48) during were recorded under T₆ (Vermiwash-40 ml/sq. m) which was found statistically at par with its concentration *i.e.* Vermiwash-30ml/sq. m (T₅) and PSB (Photosynthetic Bacteria

Rhodosudomonaspalustris) -40ml/sq. m (T₃). Similar result found that the **Nayak et al., (2016)** reported that the application of biofertilizers (Azotobacter + Azospirillum + PSB), lime and organic manure (Vermicompost) in maximum number of fruit per plant (45.48). The maximum average weight for 10 fruits (1653.53 g) during were recorded under T₉ (Azotobacter-15 ml/sq.m) which was found statistically at par with its concentration *i.e.* Vermiwash-40ml/sq.m (T₆) and PSB (Photosynthetic Bacteria *Rhodosudomonaspalustris*-40 ml/sq.m (T₃). Similar result found that the **Lucas-Garcia et al., (2004)** reported that gourd plants supplied with Azotobacter recorded increased of fruits weight 1600.33 g.

Formatted: Font: Italic

Formatted: Font: Italic

Table 3. Evaluation of best performing microbial culture in relation to yield of Sponge Gourd .

Treatments	Number of fruits per plant	Av. Weight for 10 fruit (g)	Fruit length (cm)
Control(T ₀)	33.75	1626.66	20
PSB- 20ml/sq.m (T ₁)	41.25	1590.00	22
PSB- 30ml/sq.m (T ₂)	42.50	1546.66	23
PSB- 40ml/sq.m (T ₃)	43.66	1506.66	24
Vermiwash-20 ml/sq.m (T ₄)	42.66	1640.00	25
Vermiwash-30 ml/sq.m (T ₅)	44.33	1643.33	26
Vermiwash-40 ml/sq.m (T ₆)	45.48	1546.66	27.33
Azotobacter-8ml/sq.m (T ₇)	37.25	1513.33	25.00
Azotobacter-10 ml/sq.m (T ₈)	39.50	1563.33	26.00
Azotobacter-15 ml/sq.m(T ₉)	41.33	1653.33	26.66
F-test	S	S	S
SE(d)	0.84	22.35	1.93
C.V.	2.50	1.73	9.67
CD at 5% level	1.78	47.33	4.09

The maximum fruit length (27.33 cm) was found in Vermiwash – 40 ml/sq.m (T₆) followed by Azotobacter- 15ml/sq.m 26.66 cm (T₉) and Vermiwash– 30ml/sq.m 26.00 cm (T₅), Azotobacter- 15ml/sq.m 26.00 cm the minimum fruit length (20 cm) was observed in control plants (T₀). **Reddy et al., (2012)** reported similar **Nayak et**

al., (2016) reported that the application combination with biofertilizers (Azotobacter + Azospirillum + PSB), lime and organic manure (Vermicompost) in sponge gourd to increase the length length of fruit (22.50 cm).

Table 4. Evaluation of best performing microbial culture in relation to yield ofSponge Gourd.

Treatments	Fruit girth (mm)	Av. Fruit Yield per plot (kg)	Fruit yield/ha (q)
Control(T ₀)	23.66	19.33	124.07
PSB- 20ml/sq.m (T ₁)	25.33	19.69	145.75
PSB- 30ml/sq.m (T ₂)	27.00	21.86	145.75
PSB- 40ml/sq.m (T ₃)	30.00	25.18	148.24
Vermiwash-20 ml/sq.m (T ₄)	26.00	21.95	146.19
Vermiwash-30 ml/sq.m (T ₅)	27.00	24.37	155.47
Vermiwash-40 ml/sq.m (T ₆)	30.00	25.80	164.50
Azotobacter-8ml/sq.m (T ₇)	25.00	20.82	127.00
Azotobacter-10 ml/sq.m (T ₈)	30.00	21.33	137.28
Azotobacter-15 ml/sq.m(T ₉)	32.00	21.35	149.07
F-test	S	S	S
SE(d)	2.48	0.95	5.07
C.V.	11.02	5.25	5.05
CD at 5% level	3.05	2.01	10.74

The maximum fruit girth (32.00 mm) during were recorded under T₉ (Azotobacter-15ml/sq.m) which was found statistically at par with its concentration *i.e.* PSB (Photosynthetic Bacteria *Rhodospirillum rubrum*-40ml/sq.m (T₃), Vermiwash-40 ml/sq.m(T₆) and Azotobacter -10ml/sq.m (T₈). Nayak *et al.*, (2016) reported that the application combination with biofertilizers (Azotobacter + Azospirillum +

Formatted: Font: Italic

PSB), lime and organic manure (Vermicompost) in sponge gourd to increase the length length of fruit (31.50 mm).Maximum average fruit yield per plot (25.80 kg) was found in Vermiwash - 40ml/sq.m (T₆) followed by PSB (Photosynthetic Bacteria *Rhodosudomonaspalustris*- 40 ml/sq.m 25.18 kg (T₃) andVermiwash – 30 ml/sq.m 10.77 kg (T₅) the minimum average fruit yield per plot (19.33 kg) was observed in control plants (T₀).**Reddy et al., (2012)** reported similar results in muskmelon, **Yogesh et al., (2009)** in cucumber.Maximum fruit yield/ ha (q)(164.50) was found in Vermiwash – 40 ml/sq.m (T₆) followed by Vermiwash – 30 ml/sq.m 155.47 q (T₅) andAzotobacter – 15ml/sq.m 149.07 q (T₉) the minimum fruit yield/ ha (124.07 q) was observed in control plants (T₀). Similar result found that the **Yobo et al., (2004)**, who recorded significantly higher fruit yield in gourd.

Formatted: Font: Italic

CONCLUSION:-

From The results of the present investigation concluded that the different microbial cultureof Sponge Gourd(*Luffa cylindrica* L.)could be sources without affecting growth performance and yield.Results revealed that plant growth characteristicstreatment T₉ (Azotobacter-15 ml/sq.m) (619.75) recorded significantly highest vine length .

Yield parameter like Fruit yield/ha(q.) was recorded highest in treatment T₆ (Vermiwash-40ml/plant)(164.50q/ha).

quality parameter like total soluble solid (tss)(obrix) was recorded highest in treatment T₃(photosynthetic bacteria-40ml/plant) (21.50).

References:-

- Abdelhafez Ahmed, A.M. and AbdelmonsiefRihan, A. (2006).** Effect of VAM inoculation on growth, yield and nutrient content of cantaloupe and cucumber under different water regimes. *Research Journal of Agriculture and Biological Science*, 2(6): 503-508.
- Bal, K. J., Hari, B. K. C., Radha, K. T., Madhusudan, G., Bhuwon, R. S. and Madhusudan, P. U. (2004).** Descriptors for Sponge Gourd (*Luffa cylindrica* L.) Roem. NARC, LIBIRD and IPGRI.
- Chaudhari, H. K. (2000).** Elementary principles of plant breeding, Oxford and IBH Publishing, New Delhi, India
- Das, R., Mandal A.R., Priya A., Das S.P. and Kabira, J. (2015).** Evaluation of integrated nutrient management on performance of bottle gourd *Lagenariasiceraria* (Molina) Standl.). *Journal of Applied and Natural Science*, 7(1):18-25.
- Fischer, R. R. and Yates, F.(1963).** Stastical tables for biological, agricultural and medical research. Sixth edition, Oliver and Boyd, Tweeddale Court, Edinberg, pp. 747-777.
- Karuthamani, M., Natarajan, S. and Thamburaj, S. (2005).** Effect of inorganic and bio-fertilizer on growth, flowering and yield of pumpkin (*Cucurbita moschata*). *South Indian Horticulture*. 43(516):134-136.
- Lucas-Garcia, N. and Suma, R. (2004).** Effect of genotype of Sponge Gourd (*Luffa cylindrical*) on growth and yield. *International Journal of Chemical Studies* 6(5): 3062-3065.
- Nayak, D. A., Pradhan M., Mohanty S., Paridaand A. K. and Mahapatra P. (2016).** Effect of integrated nutrient management on productivity and profitability of pointed gourd (*Trichosanthesdioica* Roxb.). *Journal of Crop & Weed*, 12 (1):25-31.

- Okusanya, O. T., (1978).** The effects of light and temperature on the germination and growth of *Luffa aegyptica (L. cylindrica)*. *Pysiol. Plant*, 44(4):404-408.
- Patel, B.J., Wagh, A.P., Umbarkar, P.S. and Bondre, S.V. (2018).** Integrated nutrient management studies in bottle gourd. *Journal of Pharmacognosy and Phytochemistry*, 7(5): 1383-1385.
- Rai, N. and Bansiwat, K. (2008).** Vermiwash : an excellent source of nutrition for plant growth, *Electronic Journal of Environment Sciences*, 1:19-21.
- Reddy, P.K. and Rao, P.V. (2012).** Growth and yield of bitter gourd (*Momordicacharantia L.*) as influenced by vermicompost and nitrogen management practices. *Journal of Research, ANGRAU*, 32 (3): 15-20.
- Sreenivas, C.H., Muralidhar, S. and Singarao, M. (2000).** Yield and quality of ridge gourd fruits as influenced by different levels of inorganic fertilizer and vermicompost. *Annals of Agricultural Research*, 21(2): 262-266.
- Thriveni, V., Mishra H.N., Pattanayak S.K., Sahoo G.S. and Thomson T. (2015).** Effect of inorganic, organic fertilizers and bio-fertilizers on growth, flowering, yield and quality attributes of bitter gourd (*Momordicacharantia L.*). *International Journal of Farm Sciences*, 5(1): 24-29.
- Vimala, B. and Natarajan, S. (2000).** Effect of nitrogen, phosphorus and biofertilizers on pod characters, yield and quality in pea (*Pisum sativum L. spp.* Hortense). *South Indian Horticulture* 48(1-6): 60-63.
- Yobo, A. and Tabar, S.M. (2004).** Genotype performance Sponge Gourd (*Luffa cylindrical*) fruit development can improve fruit quality. *Horticultural Environmental Biotechnology*, 50(3): 1-6.