

# Study Of Potential Utility Of Predictive Laboratory Parameters For Early Triage Of SARI Patients

## ABSTRACT

**INTRODUCTION:** The novel corona virus (COVID-19), which began in Wuhan, China, in December 2019, has been declared pandemic by the World Health Organization (WHO) on 11th March 2021. Till now, even when nations are in race for developing vaccine against COVID-19, diagnosing the cases and isolating them still remain the only way to contain the disease spread. Hence, effective triage of the suspected COVID 19 patients, becomes an effective tool in preventing disease spread, especially in developing nations. Simple hematological parameters like leukocyte counts, platelet counts, hemoglobin and parameters derived from complete blood counts namely NLR, derived NLR, PLR and liver function tests can be studied and analysed to differentiate COVID 19 patients from other non-COVID 19 Severe Acute Respiratory Illness (SARI) patients. These are the investigations at first point of contact with the patient and are relatively cost effective, keeping in mind the population of developing nations.

**PATIENTS & METHODS:** Retrospective case control study conducted over a period of one month including 286 consecutive Severe Acute Respiratory Illness(SARI) patients admitted in a tertiary care hospital in Delhi. The patients were divided into COVID-19 positive and COVID-19 negative groups based on their results of reverse transcriptase-polymerase chain reaction (RT-PCR) for COVID-19. The demographic

data, and routine hematological and biochemical parameters including derived parameters like Neutrophil lymphocyte ratio (NLR), derived NLR ratio, Platelet lymphocyte ratio (PLR) and Prognostic nutrition index (PNI) were analyzed and compared between the two groups using SPSS version 23.0. A p value of <0.05 was considered significant.

RESULTS: The COVID-19 positive patients had a significantly higher lymphocyte percentage (p value=0.017) and Serum Glutamate Pyruvate Transaminase(SGPT)( p value=0.033), and a significantly lower Absolute Neutrophil Count (ANC)( p value=0.023), NLR(p value=0.033) and platelet count(p valu=0.001). No significant difference in other haematological and biochemical parameters was found.

CONCLUSION: NLR, platelet count, lymphocyte percentage and SGPT are the simple laboratory biomarkers of inflammation that can be utilized for screening and triage in COVID 19 disease with variable sensitivity and specificity.

KEYWORDS: Age, COVID-19, lymphocyte percentage, Neutrophil to Lymphocyte ratio(NLR), platelet count, routine, Serum Glutamate Pyruvate Transaminase(SGPT), triage

#### INTRODUCTION:

The novel corona virus (COVID-19), which began in Wuhan, China, in December 2019, was declared pandemic by the World Health Organization (WHO) on 11th March 2021.<sup>1</sup> Globally it has resulted in 764,474,387 confirmed cases including

6,915,286 deaths<sup>2</sup> and has threatened global public health. India itself has reported 44,905,827 confirmed cases with 531,369 deaths. Also as on 10 April 2023, a total of 2,206,624,273 vaccine doses have been administered.<sup>2</sup> In spite of rapid development of knowledge, precautionary measures, and clinical trials, regulatory bodies and government administrations are facing a great challenge globally in various aspects to contain the spread of the disease. Development of vaccine and its worldwide administration still remains a global challenge. Although the containment measures carried out, have atleast for the moment decreased the risk of contagion significantly, but a truthful diagnostic testing still remains the most important step. Hence, even at present, a key to slow the disease spread is rapid diagnosis and isolation of cases and this relies on the effective triage protocols for identifying suspected COVID-19 patients and their isolation or diagnosis for a timely management and optimal utilization of the limited human and technical resources for testing and management of COVID-19 disease.<sup>3</sup> Currently, the gold standard of diagnosis is the RT-PCR test, which detects the viral RNA and provide results within 6 hours. However, false negative results due to low viral load and a relative shortage of detection kits are the major limitations due to which many patients are left undetected and undiagnosed<sup>4</sup>.

Immune dysregulation and damage to lymphocytes have been observed in patients of Coronavirus disease 2019(COVID 19). Increased serum levels of proinflammatory cytokines and chemokines (TNF, IL-1,IL-6,IL-8) suggest a possible role of hyperinflammatory state in the pathogenesis of COVID -19<sup>5</sup>. Neutrophil/lymphocyte ratio (NLR) and Platelet/Lymphocyte ratio (PLR) are well established inflammatory indices<sup>6</sup>. NLR and PLR have been used to evaluate systemic inflammation in

neoplastic and cardiovascular diseases<sup>7,8</sup>. Recent published studies done have shown the prognostic utility of NLR and PLR in differentiating severe and non-severe COVID 19 patients. However there is very limited evidence on use of these simple parameters in differentiating COVID 19 patients from other Severe Acute Respiratory Illness (SARI) patients<sup>9</sup>.

The aim of this article is to study the hematological and biochemical laboratory parameters (Neutrophil counts, lymphocyte count, platelet counts, NLR, PLR, SGOT, SGPT) in COVID-19 positive patients and compare them with COVID-19 negative severe acute respiratory illness (SARI) patients to assess their potential utility as soft pointers for COVID-19 infection in resource constraint settings.

#### MATERIALS & METHODS:

**Study Design:** It is a retrospective case control study conducted over a period of one month (September 5, 2020 to October 4, 2020) including 286 consecutive SARI patients admitted in ABVIMS & Dr Ram Manohar Lohia hospital, a designated nodal centre for management of COVID-19 patients in New Delhi.

The study included all adult patients who were diagnosed as SARI at our fever clinics and at SARI casualty as per WHO case definition<sup>5</sup> and screened for COVID-19. The patients were divided into cases and control groups based on their results of real time reverse transcriptase-polymerase chain reaction (RT-PCR) for COVID-19, i.e. COVID-19 positive (cases) and COVID-19 negative (control group). The information recorded included the demographic data, hematological and biochemical

parameters of inflammation including Complete Blood Counts (CBC), Renal function tests, Liver function tests, Total protein, and Serum albumin. Neutrophil lymphocyte ratio (NLR), derived NLR ratio, Platelet lymphocyte ratio (PLR) and Prognostic nutrition index (PNI), were calculated as per the case definitions described. All the data was compared between the COVID positive and COVID-19 negative groups. Also the accuracy of significant laboratory parameters in predicting cases with positive RT-PCR was evaluated using area under the ROC curve (AUC). The optimal cut off values, sensitivity and specificity of the statistically significant parameters were also calculated after the analysis of ROC curve.

Definitions: NLR is the ratio of absolute neutrophil count to absolute lymphocyte count. dNLR is the ratio of neutrophil count to WBC count minus neutrophil count ( $ANC/TLC-ANC$ ). PLR is the ratio of platelet count to absolute lymphocyte count. PNI (Prognostic Nutrition Index) is a common inflammatory biomarker calculated as  $(10 \times \text{serum albumin g/dL} + 0.005 \times \text{total lymphocyte count})$ .

Statistical analysis: The data was entered in MS EXCEL spreadsheet and analysis was done using Statistical Package for Social Sciences (SPSS) version 23.0. Categorical variables were presented in number and percentage (%) and continuous variables were presented as mean  $\pm$ SD and median. Normality of data was tested by Kolmogorov-Smirnov test. If the normality was rejected then non parametric test was used. Quantitative variables were compared using Mann-Whitney Test (as the data sets were not normally distributed) between the two groups. Qualitative variables were compared using Chi-Square test/Fisher Exact test. Receiver operating

characteristic curve was used to find out cut off point of parameters for predicting positive findings. A p value of <0.05 was considered statistically significant.

## RESULTS:

Demographic Characteristics: Two hundred eighty six SARI cases were included in the study. The result of RT-PCR for COVID-19 was positive in 127 (44.4%) cases and negative in 159 (55.5%). The cases ranged from 15 to 85 years in age with a mean age of  $50.76 \pm 15.77$  years for positive group and  $46.47 \pm 18.92$  years for negative group. There was a significant age difference between the two with COVID-19 positivity being more frequent in patients >50 years ( $p=0.045$ ). (Figure 1) Groups of cases with positive and negative RT-PCR showed no significant gender preference ( $p=0.13$ ). (Figure 2)

## Blood Routine Parameters:

Table 1 compares the laboratory parameters of COVID-19 positive patients with the negative group. The COVID-19 positive patients had a significantly higher lymphocyte percentage (19.7% vs 16.7%,  $p$  value<0.05) and SGPT (74.21 vs 42.21 U/L,  $p$  value<0.05), and a significantly lower Absolute Neutrophil

Count(ANC)[7255.4 vs 8894.1/ $\mu$ l], a significantly reduced NLR( 5.74 vs 6.98, p value<0.05) and platelet count (1.8 vs 2.16 lacs/ $\mu$ l]. Though not statistically significant, but we also found that the mean total leukocyte count was lower in COVID positive patients as compared to the control group (9417 vs 11108/ $\mu$ l] but not reaching the cutoffs for leucopenia. Also the PLR was lower in COVID positive patients (155.8 vs 183.3) but with a p value of 0.091, hence not statistically significant.

No significant difference in other hematological and biochemical parameters was found between the patient and the control group.

**Receiver operating curves (ROC)** were drawn for the laboratory parameters that showed a significant p value. Taking the null hypothesis into account as true area = 0.5, the parameters that showed value in predicting cases with positive RT-PCR for COVID-19 were age (AUC=0.544), lymphocyte count (AUC=0.588), NLR (AUC=0.579), platelet count (AUC=0.598) and SGPT (AUC=0.602). (Figure3, table 2). The ANC could not be used as a potential diagnostic marker for subsequent analysis because its AUC was less than 0.50.

Appropriate cut off values for the significant parameters (Age, lymphocyte count, NLR, platelet count & SGPT) were statistically derived using ROC curves. (Table 3).

The cut off values were 15.5% for lymphocyte percentage (sensitivity=70.2% and specificity=58%), 1.74 lacs/ $\mu$ l for platelet count (sensitivity=64.2% and specificity=55.9%), 29.5U/L for SGPT (sensitivity=70.2% and specificity=55.9%),

and 3.8 for NLR (sensitivity=62.9% and specificity=50.4%). Lymphocyte percentage and SGPT had the highest sensitivity while SGPT had the highest specificity in diagnosing COVID 19 positive patients.

## DISCUSSION:

Considering the rapid spread and the global health impact of COVID 19, it is required to continue research on new triage and diagnostic modalities to improve case identification. Molecular identification of SARS-CoV-2 in low-resource settings remains centralized and time-consuming, resulting in a delayed diagnosis of COVID-19. The dysregulation of immune response and excessive inflammation are the key elements of pathogenesis of COVID-19<sup>5</sup>. The blood biomarkers of systemic inflammation reflect the immune status and hence are potential indicators for prognosis and diagnosis of COVID-19 patients. This study aimed to identify frequently performed laboratory parameters that can assist in COVID-19 case identification. The study revealed a significantly raised lymphocyte percentage and SGPT levels, and a significantly lower ANC, NLR and platelet count in COVID 19 positive patients. All the other hematological and biochemical parameters did not show any role in predicting COVID-19 positive cases.

Various studies have found similar results in severe COVID-19 patients. A recent study on eighty lab confirmed COVID 19 patients admitted in ICU of a tertiary care hospital in India, found a positive correlation between NLR and severe critically ill ICU patients (p value= 0.012)<sup>10</sup>. A similar study focusing on the identification of risk

factors associated with COVID 19 severity, found an elevated NLR (cut off value=6.8) in 51.9% of study subjects. They also found a positive correlation of raised NLR with LDH ( pvalue=0.016) and CRP(p value=0.026)<sup>11</sup>. But, there are studies which have found results similar to our results. They found a significantly decreased NLR in COVID positive vs negative patients. They observed an AUC of 0.739, cut off =1.8, sensitivity of 0.59 and specificity of 0.76, and proposed the use of NLR as an early diagnostic marker for COVID 19.<sup>12</sup> Also studies have found a decreased mean ANC 3.89(2.61-6.33) and mean NLR of 3.21(1.94-7.11) in all the COVID positive patients vs a mean ANC and NLR of 7.41 and 12.41 respectively in severe COVID-19 patients (non- survivors).<sup>13</sup> Another study found a normal or mildly reduced NLR and ANC <1.2 and <4000/ $\mu$ l respectively in asymptomatic to mild cases of Covid-19. Severe cases were coupled with leucocytosis and increased NLR >3.13 or higher than >5.0.<sup>14</sup> Our study included a heterogeneous population of patients ranging from mild to severe category of COVID-19 which may explain our results.

A meta-analysis by including three studies in patients with severe or fatal COVID 19 found an increase in leucocyte and neutrophil count and a decrease in lymphocyte and platelet count<sup>15</sup>. However relative to non-COVID-19 infections, significantly lower absolute leucocyte or neutrophil and platelet counts during early stages of the disease have been demonstrated in a meta analysis of seven studies although the mean leucocyte count of the studies included did not exceed the lower limit for categorization as leucopenia or neutropenia. They found that with COVID-19 disease progression, both leukocyte and neutrophil counts were significantly higher in the

severe groups<sup>16</sup>. We too have found a lower mean leucocyte count in the COVID-19 cohort and a significantly lower ANC and NLR. Study by Huang et al revealed that asymptomatic COVID-19 patients with similar viral load as symptomatic patients, had a higher lymphocyte percentage indicative of a better immune response against the virus. This can also possibly explain a higher lymphocyte percentage during early course of the disease.<sup>17</sup>

However, many studies found neutrophilia, that is raised ANC and NLR in COVID-19 patients. It was initially thought to be due to secondary bacterial infections but recent studies have implicated excessive neutrophilia and neutrophil extracellular traps (NET) as the cause of hyper inflammation and lung injury.<sup>18</sup> More so the exacerbated NET formation is thought to drive a cascade of inflammatory reactions that favors micro thrombosis, thus suggesting raised ANC and NLR to be independent prognostic features as well as predictors of severe COVID-19 infection.<sup>19</sup> Yang et al. and Qu et al. in their respective studies had elaborated upon the predictive role of raised NLR as circulating biomarker that can represent viral hyper inflammation and immune status in severe COVID-19 and described it as an independent biomarker for poor prognosis.<sup>18,20</sup> . This study though caters only Chinese population with severe COVID - 19 disease, hence may not be representative of a heterogeneous population with mild, moderate and severe patients.

Thrombocytopenia has been the most common hematological change seen in patients with COVID-19 in various studies including ours. Xu P et al. proposed direct bone marrow suppression along with immune mediated destruction of platelets and platelet

aggregation in lungs to be the three mechanisms responsible for thrombocytopenia in COVID-19.<sup>21</sup> The immune dysfunction and consequent cytokine storm associated with COVID-19 infection leads to lung injury and destruction of hematopoietic destruction and bone marrow growth inhibition. The damaged lung tissues and pulmonary endothelial cells may activate platelets in the lungs, resulting in platelet aggregation and formation of micro thrombi. The COVID-19 infection may also increase the levels of auto antibodies and immune complex formation which gets deposited on the surface of platelets, and subsequent recognition and destruction by the reticuloendothelial system.

It can be noted that the ACE-2 protein has an abundant expression in intestinal epithelial cells, renal tubular epithelial cells, alveolar epithelial cells, heart, artery smooth muscle cells and gastrointestinal cells. Therefore apart from lungs, COVID-19 is also speculated to infect liver, gastrointestinal system and kidney which may lead to abnormal liver functions, diarrhea and acute kidney injury.<sup>22</sup> A meta-analysis of 521 studies found that patients with COVID-19 had higher than expected prevalence of liver injury, and the extent of the injury was associated with the severity of the disease. They found a substantial elevation in enzymes (>3 times) in almost 16% of the COVID-19 positive patients.<sup>23</sup> We found a significant elevation in SGPT levels of COVID-19 patients however SGOT, ALP and albumin did not show any significant difference. On the contrary, Chen et al., found that albumin had decreased, but SGPT and SGOT showed no significant changes.<sup>24</sup> Furthermore another meta-analysis of 128 studies by Kumar et al. found hypoalbuminemia [61.27% (48.24–72.87)] and elevations of SGPT [23.28% (19.92–27.01)], and SGOT

[23.41% (18.84–28.70)] to be the most frequent abnormalities.<sup>25</sup> The liver enzyme derangement can be due to direct viral injury, multiple antibiotics and antiviral drugs intake prior to presentation or due to the COVID-19 infection induced cytokine storm.

We found no significant differences or derangements in the renal function tests i.e. blood urea, serum creatinine and uric acid in COVID-19 positive and negative SARI patients. Sharma et al found similar findings of essentially normal renal profile in COVID-19 patients.<sup>26</sup> Nevertheless a close monitoring of the renal function in COVID-19 patients is required.

We made an attempt to study the impact of overall nutritional status as a marker of susceptibility to COVID-19 by comparing PNI between COVID-19 positive and negative SARI patients. No significant difference was found between the PNI of the two groups. However study by Wang et al found a significant difference in PNI between COVID-19 survivors and non survivors, and suggested it as a predictive biomarker for severity and progression in COVID-19 patients.<sup>13</sup> However no study done so far has utilized its utility as diagnostic marker for the purpose of triage of COVID-19 patients. Therefore we suggest that the nutritional status has no bearing on the susceptibility to COVID-19 infection, however it may have an impact on the prognosis of the disease.

The core strength of the present study is the inclusion of a large, contemporary, COVID-19 negative control population, which allowed us to critically reappraise the haematological features of patients with COVID-19.

There were some limitations in our study. First, it was a retrospective study and the sample size was small. Secondly, the lack of clinical data is one of the deficiencies of this study, so we can't rule out the influence of stress response and treatment on blood routine parameters. Further research is needed to overcome these limitations. Nevertheless, our results provided important illumination on this topic.

CONCLUSION: NLR, platelet count, lymphocyte percentage and SGPT are the simple, quick, inexpensive and reproducible laboratory biomarkers of inflammation that can be utilized for screening and triage in COVID-19 disease with variable sensitivity and specificity. Yet further studies with larger sample size are required to better validate the results.

#### Ethical Approval

The study protocol was approved by the Ethics Committee of ABVIMS & Dr Ram Manohar Lohia hospital. (ethical code: 403(52/2020)IEC/ABVIMS/RMLH136)

#### REFERENCES:

1. WHO Director-General's opening remarks at the media briefing on COVID-19 - 25 May 2020. <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---25-may-2020>.

2. WHO Coronavirus Disease (COVID-19) Dashboard [Internet]. Covid19.who.int. 2023 [last updated April 26, 2023]. Available from: <https://covid19.who.int/>
3. Li Q, Ding X, Xia G, et al. Eosinopenia and elevated C-reactive protein facilitate triage of COVID-19 patients in fever clinic: a retrospective case-control study [published online ahead of print, 2020 May 3]. *E Clinical Medicine*. 2020;100375.
4. Ou M, Zhu J, Ji P, Li H, Zhong Z, Li B, Pang J, Zhang J, Zheng X. Risk factors of severe cases with COVID-19: a meta-analysis. *Epidemiol Infect*. 2020 Aug 12;148:e175.
5. Qin C, Zhou L, Hu Z, et al. Dysregulation of Immune Response in Patients with Coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis*. 2020;71(15):762-768.
6. Damar Çakırca T, Torun A, Çakırca G, Portakal RD. Role of NLR, PLR, ELR and CLR in differentiating COVID-19 patients with and without pneumonia. *Int J Clin Pract*. 2021 Nov;75(11):e14781.
7. Diem S, Schmid S, Krapf M, Flatz L, Born D, Jochum W, Templeton AJ, Früh M. Neutrophil-to-Lymphocyte ratio (NLR) and Platelet-to-Lymphocyte ratio (PLR) as prognostic markers in patients with non-small cell lung cancer (NSCLC) treated with nivolumab. *Lung Cancer*. 2017 Sep;111:176-181.
8. Kurtul A, Ornek E. Platelet to Lymphocyte Ratio in Cardiovascular Diseases: A Systematic Review. *Angiology*. 2019 Oct;70(9):802-818.

9. Sarkar S, Khanna P, Singh AK. The Impact of Neutrophil-Lymphocyte Count Ratio in COVID-19: A Systematic Review and Meta-Analysis. *J Intensive Care Med.* 2022 Jul;37(7):857-869.
10. Neetu Goyal, I. M. Evaluation and comparison of utility of neutrophil-to-lymphocyte ratio (NLR) as a diagnostic marker in covid-19 infection. *European Journal of Molecular & Clinical Medicine*, 2022; 9(6): 2127-2134.
11. W Jelassi, N Abid, M Loukil, N Gader, M Ben Ali, I Chaabane, K Bouzaidi, H Ghrairi. Neutrophil-to-lymphocyte : a useful biomarker for predicting severity in coronavirus disease-2019 (COVID19). *European Respiratory Journal* Sep 2022, 60 (suppl 66)
12. Usul E, Şan İ, Bekgöz B, Şahin A. Role of hematological parameters in COVID-19 patients in the emergency room. *Biomark Med.* 2020;14(13):1207-1215.
13. Wang, R, He, M, Yin, W, et al. The Prognostic Nutritional Index is associated with mortality of COVID-19 patients in Wuhan, China. *J Clin Lab Anal.* 2020; 34:e23566.
14. Zahorec R, Hulin I, Zahorec P. Rationale Use of Neutrophil-to-lymphocyte ratio for early diagnosis and stratification of COVID-19. *Bratisl Lek Listy.* 2020;121(7):466-470.
15. Henry B, de Oliveira M, Benoit S, Plebani M, Lippi G. Hematologic, biochemical and immune biomarker abnormalities associated with severe illness and mortality in coronavirus disease 2019 (COVID-19): a meta-

- analysis. *Clinical Chemistry and Laboratory Medicine (CCLM)*. 2020;58(7):1021-1028.
16. Soraya GV, Ulhaq ZS. Crucial laboratory parameters in COVID-19 diagnosis and prognosis: An updated meta-analysis. *Med Clin (Barc)*. 2020.
  17. Huan Han, Zaichao Xu, Xiaoming Cheng, Youquan Zhong, Li Yuan, Fubing Wang, Yan Li, Fang Liu, Yingan Jiang, Chengliang Zhu, Yuchen Xia. Descriptive, Retrospective Study of the Clinical Characteristics of Asymptomatic COVID-19 Patients. *mSphere*, 2020; 5 (5)
  18. Yang A, Liu J, Tao W, Li H. The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients. *International Immunopharmacology*. 2020;84:106504.
  19. Wang J, Li Q, Yin Y, Zhang Y, Cao Y, Lin X et al. Excessive Neutrophils and Neutrophil Extracellular Traps in COVID-19. *SSRN Electronic Journal*. 2020. <https://doi.org/10.3389/fimmu.2020.02063>
  20. Qu R, Ling Y, Zhang Y, Wei L, Chen X, Li X et al. Platelet to lymphocyte ratio is associated with prognosis in patients with coronavirus disease 19. *Journal of Medical Virology*. 2020;92(9):1533-1541.
  21. Xu P, Zhou Q, Xu J. Mechanism of thrombocytopenia in COVID-19 patients. *Annals of Hematology*. 2020;99(6):1205-1208. doi: 10.1007/s00277-020-04019-0.
  22. Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of

different human organs vulnerable to 2019-nCoV infection. *Front med.* 2020;1-8. doi: 10.1007/s11684-020-0754-0

23. Ahmed J, Rizwan T, Malik F, Akhter R, Malik M, Ahmad J et al. COVID-19 and Liver Injury: A Systematic Review and Meta-Analysis. *Cureus.* 2020.
24. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *The Lancet.* 2020;395(10223):507-13. doi.org/10.1016/S0140-6736(20)30211-7
25. Kumar-M P, Mishra S, Jha D, Shukla J, Choudhury A, Mahindra R et al. Coronavirus disease (COVID-19) and the liver: a comprehensive systematic review and meta-analysis. *Herpetology International.* 2020.
26. Sharma P, Aggarwal V, Sapariya N, Baweja S, Nagori R, Chhabra M. Renal profile of COVID-19 infected patients admitted in a tertiary care hospital in Western Rajasthan. *International Surgery Journal.* 2020;7(7):2332.

**Table 1 : Comparison Of The Laboratory Parameters Between The Cases With Positive And Negative RT-PCR For COVID-19 Infection**

Parameter (Units; mean±SD)	COVID-19 RT-PCR		p value
	Negative (N=159)	Positive (N=127)	
Hemoglobin(gm/dL)	11.32±2.10	11.30±2.53	0.935

<b>Total leucocyte count(/<math>\mu</math>l)</b>	11108.00 $\pm$ 6618.24	9417.30 $\pm$ 8357.54	0.057
<b>Neutrophil count(%)</b>	77.48 $\pm$ 11.12	75.14 $\pm$ 11.79	0.087
<b>Lymphocyte count(%)</b>	16.74 $\pm$ 9.82	19.73 $\pm$ 11.25	<b>0.017</b>
<b>Absolute neutrophil count (ANC)(/<math>\mu</math>l)</b>	8894.10 $\pm$ 5824.97	7255.40 $\pm$ 6306.43	<b>0.023</b>
<b>Absolute lymphocyte count (ALC)(/<math>\mu</math>l)</b>	1616.70 $\pm$ 1078.90	1672.00 $\pm$ 1961.10	0.763
<b>Neutrophil lymphocyte ratio(NLR)</b>	6.98 $\pm$ 5.24	5.74 $\pm$ 6.98	<b>0.033</b>
<b>Derived NLR (dNLR)</b>	4.70 $\pm$ 3.12	4.20 $\pm$ 3.05	0.17
<b>Platelet count(/<math>\mu</math>l)</b>	216060 $\pm$ 92773.17	180980 $\pm$ 80426.78	<b>0.001</b>
<b>Platelet lymphocyte ratio (PLR)</b>	183.30 $\pm$ 131.69	155.88 $\pm$ 140.67	0.091
<b>Urea(mg/dL)</b>	59.57 $\pm$ 74.26	52.45 $\pm$ 53.03	0.364
<b>Serum creatinine(mg/dL)</b>	1.75 $\pm$ 2.61	1.71 $\pm$ 2.59	0.884
<b>Uric acid(mg/dL)</b>	7.02 $\pm$ 5.61	6.64 $\pm$ 3.38	0.497
<b>Total bilirubin(mg/dL)</b>	1.23 $\pm$ 3.60	0.74 $\pm$ 0.63	0.127
<b>Direct bilirubin(mg/dL)</b>	0.42 $\pm$ 0.87	0.36 $\pm$ 0.73	0.513

<b>Indirect bilirubin(mg/dL)</b>	0.81±3.15	0.45±0.41	0.197
<b>SGOT*(U/L)</b>	54.52±51.99	62.64±72.86	0.276
<b>SGPT**(U/L)</b>	42.21±45.58	74.21±179.45	<b>0.033</b>
<b>ALP***(U/L)</b>	123.61±84.8	126.33±127.24	0.83
<b>Total protein(gm/dL)</b>	11.58±56.75	6.59±0.78	0.335
<b>Serum albumin(gm/dL)</b>	3.19±1.1	3.21±0.61	0.903
<b>PNI****</b>	37.79±14.51	39.40±13.99	0.344

**Notes:**

Abbreviations: \*SGOT,Serum glutamic oxaloacetic transaminase \*\*SGPT,Serum glutamic pyruvic transaminase \*\*\*ALP,Alkaline phosphatase \*\*\*\*PNI,Prognostic nutrition index

**Table 2 : Area Under The Curve Of Significant Parameters**

Area Under the curve					
Test Result Variable(s)	Area under the curve(AUC)	Standard error	Asymptotic Sig <sup>b</sup>	Asymptotic 95% Confidence interval	
				Lower Bound	Upper Bound
Age	0.544	0.035		0.46	0.612

LYMPHOCYTE COUNT (%)	0.588	0.035	70.2%	0.52	0.656
NLR	0.579	0.034	62.9%	0.513	0.646
PLATELET COUNT (/µl)	0.598	0.034	64.2%	0.532	0.664
SGPT (U/L)	0.602	0.034	70.2%	0.534	0.669
<p>The test variables had at least one tie between the positive actual state group and the negative state group. Statistics may be biased.</p> <p>a. Under the non parametric assumption</p> <p>b. Null hypothesis : true area=0.5</p>					


**Table 3: Diagnostic Performance Of Laboratory Parameters In Differentiating COVID-19 Positive Cases From Controls**

Test Result Variable(s)	Positive if Greater Than or Equal To	Sensitivity	Specificity
LYMPHOCYTE COUNT (%)	12.5-15.5	70.2%	58%
NLR	3.895-4.325	62.9%	50.4%

PLATELET COUNT (/μl)	174000-188000	64.2%	55.9%
SGPT (U/L)	24.5-29.5	70.2%	59.4%


**Figure 1: Age Distribution of SARI Patients.**

Age Distribution




UNDEK

**Figure 2: Gender Distribution Of SARI Patients.**



**Figure 3: Area Under The Receiver Operating Characteristic Curve Of Different Laboratory Parameters In Predicting Cases With Positive RT-PCR For COVID-19.**

### ROC Curve



UNDER PEER