

# **Effect of Bio-fertilizer and Gibberellic Acid on growth and yield of baby corn (*Zea mays* L.)**

## ABSTRACT

The field experiment was conducted during *Rabi* season 2021-22 at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P). The soil of the experiment plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.2), low in organic carbon (0.48%), available N (171.48 kg/ha), available P (13.6 kg/ha) and available K (215.4 kg/ha). The treatments consisted of three levels of Gibberellic acid 50, 100 and 150 ppm of foliar application and (20 g and 10g per kg of water) and two levels of Bio fertilizer {Azospirillum seed inoculation (20 g/1kg of seed) and Azotobacter seed inoculation (20 g/1kg of seed)} respectively. The experiment was laid out in randomized block design with nine treatments and were replicated thrice. Results defined that maximum plant height (185.29 cm), dry weight (90.42 g/plant), Number of cobs per plant (2.64), cob length (19.76 cm), cob weight with husk (48.54 g/cob) and cob weight without husk (12.51 g/cob) Cob yield with husk (11.04 t/ ha), Cob yield without husk (4.15 t/ha) were recorded maximum with application of Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm treatment combination, respectively.

**Keywords:** *Gibberellic Acid, Biofertilizer, yield, Economics, Baby corn.*

## INTRODUCTION

Maize (*Zea mays* L.) is the third most important cereal crop in India after wheat and rice. It is grown all over the world under a widerange of climate. Currently it is cultivated in an area of 9.2 m ha with a production of 27.8 m t and productivity of 2965 kg/ha in India (IIMR 2022). Maize (*Zea mays* L.) is the third most important cereal crop in the agricultural economy after wheat and rice, in the world as well as in India. It is popularly known as “miracle crop” and “Queen of Cereals”. Maize is recognized as the “golden food” because of its higher potentiality of grain yield and higher nutritional value. Baby corn is an extremely easy crop to produce, and for a corn growing nation, it is surprising that baby corn is an imported crop. The reason behind locally non-production of baby corn is manual labour required for detasseling, harvesting and processing, which economically prevents large-scale production. However, locally produced fresh baby corn can have an advantage over imported baby corn.

Bio-fertilizers play an important role in the increasing availability of nitrogen and phosphorus. Among several bio agent *Azospirillum* is known to fix atmospheric nitrogen and increased about 10-15 % grain yield in maize (**Patil *et al.*, 2001**). *Azotobacter* was the first and is the most common bio- fertilizer for some plants such as maize, wheat, sorghum and rice which produces some plant growth promoting metabolites, enzymes and hormones (auxin, cytokinin and gibberellin) in addition to fixing air nitrogen.

The application of plant growth regulators is one of the most important factors in improving the growth, yield and flower quality. Plant growth regulators are found as two types; bioinhibitors such as methyljasmonate and (ABA) and promoters like gibberellins, auxins and cytokinins. Gibberellins (GA3) are group of plant hormones produced by the leaves of modern plant and developing the peaks in the roots and stem (**Takahashi, Phinney *et al.* 2012**). Contain gibberellic acid, which incited elongation of plant cells and configure the fruit without seed, which overcomes the genetic dwarf leg, and increases the production of side branches, especially floral which increases the number of flowers and fruits of production ris , and can be sprayed gibberellic acid equipped on specific types of plants to increase their growth

## MATERIALS AND METHODS

This experiment was carried out investigation entitled, Effect of Bio-fertilizer and Gibberellic Acid on growth and yield of baby corn (*Zea mays* L.) was laid out during rabi season of 2022 at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.). The crop research farm is situated at 25.570 N latitude, 87.190 E longitude and at an altitude of 98 m above mean sea level. The experiment was laid out in randomized block design comprised of levels of Biofertilizers and Gibberellic acid with nine treatments and each were replicated thrice viz. Treatment 1 (Azotobacter 20g/kg + Gibberellic acid 50 ppm), Treatment 2 (Azotobacter 20g/kg + Gibberellic acid 100 ppm), Treatment 3 (Azotobacter 20g/kg + Gibberellic acid 150 ppm), Treatment 4 (Azospirillum 20g/kg + Gibberellic acid 50 ppm), Treatment 5 (Azospirillum 20g/kg + Gibberellic acid 100 ppm), Treatment 6 (Azospirillum 20g/kg + Gibberellic acid 150 ppm), Treatment 7 (Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 50 ppm), Treatment 8 (Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 100 ppm), Treatment 9 (Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm).

## RESULTS AND DISCUSSION

### Growth attributes

Effect of Bio-fertilizer and Gibberellic Acid on growth and yield of baby corn (*Zea mays* L.) presented in below Table 1.

**Plant height-** Plant height increased significantly due to the application of biofertilizers and gibberellic acid. Application of Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm gives highest plant height (185.29cm). **Meena *et al.*, (2011)** revealed that bio-fertilizer (Azotobacter) recorded higher plants height

**Plant dry weight-** The maximum dry weight weight (90.42) was recorded with application of Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm. Phyto hormones which stimulate the formation of lateral roots and absorbent root hairs, which eventually helped in uptake of higher nutrients and minerals by plants and leads to increase in higher biomass accumulation and higher plant dry weight. The results were found to be in resonance with **Zhou *et al* (2014)**.

**Yield attributes and Yield-** Maximum Number of cobs per plant (2.64), cob length (19.76 cm), cob weight with husk (48.54 g/cob) and cob weight without husk (12.51 g/cob) Cob yield with husk (11.04 t/ ha), Cob yield without husk (4.15 t/ha) were recorded maximum with application of Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm treatment combination, respectively. **Ram and Mir (2006)** indicated that the application of Azotobacter significantly increased grains per spike, test weight, grain yield and straw yield over the control. Found that the application of Azotobacter significantly increased net returns and benefit cost ratio in wheat crop over the control.

**Economics-** Maximum gross return (Rs. 108,252/ha), net return (Rs. 70,787 /ha) and B: C ratio (1.89) was recorded in treatment T9 Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm. (Table 3).

## **CONCLUSION**

Application of Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm is a fitting practice for augmenting higher babycorn yields for farmer. The maximum Net returns and B:C ratio were recorded with the application of Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm.

## REFERENCES

- Meena, K. N., Kumar, A., Rana, D. S. and Meena, M. C. 2011. Productivity and nutrient uptake of maize (*Zea mays* L.) - wheat (*Triticum aestivum*) cropping system under different bio-sources and nitrogen levels. *Indian Journal of Agronomy*, **56**(3): 182-188.
- Nagavani, M., Tariq, A., Anjum, S.A., Randhawa, M. A., Ullah, E., Naeem, M., Qamar, R., Ashraf, U. and 2014, influence of Micro Nutrient and biofertilizer on Growth and Yield Behaviour of Maize (*Zea mays* L.) Hbrids. *American J. pl. Sci.*, **5**: 2646-2654.
- Patil, R. K., Goyal, S. N., Vora, M. and Vaishnav, P. R. 2001. Response of kharif maize to inoculation with azotobacter and azospirillum at varying levels of nitrogen. *GAU Res. J.*; **27**(1-2):13-17.
- Ram, T. and Mir, M. S. 2006. Effect of integrated nutrient management on yield and yield-attributing character of wheat (*Triticum aestivum*). *Indian Journal of Agronomy*, **51**(3): 189-192.
- Verma, S. 2017. Bio-efficacy of organic formulations on crop production-A review. *Int. J. Curr. Microbiol. App. Sci* **6**(5): 648-665.

**Table 1. Effect of bio-fertilizers and Gibberellic Acid on plant growth attributes of baby corn.**

<b>S. No.</b>	<b>Treatments</b>	<b>Plant height (Cm)</b>	<b>Plant Dry weight (g)</b>
1.	Azotobacter 20g/kg + Gibberellic acid 50 ppm	156.49	79.49
2.	Azotobacter 20g/kg + Gibberellic acid 100 ppm	164.28	81.42
3.	Azotobacter 20g/kg + Gibberellic acid 150 ppm	175.12	86.79
4.	Azospirillum 20g/kg + Gibberellic acid 50 ppm	152.25	72.76
5.	Azospirillum 20g/kg + Gibberellic acid 100 ppm	160.24	80.59
6.	Azospirillum 20g/kg + Gibberellic acid 150 ppm	168.28	82.28
7.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 50 ppm	173.65	83.56
8.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 100 ppm	183.33	88.26
9.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm	185.29	90.42
	F-test	S	S
	SEm(±)	0.68	0.77
	CD (p=0.05)	2.05	2.31

**Table 2. Effect of Gibberellic Acid and bio-fertilizers on yield attributes and yield of baby corn**

<b>S. No.</b>	<b>Treatments</b>	<b>No. of cobs plant</b>	<b>Length of the cob (cm)</b>	<b>Cob weight with husk(g)</b>	<b>Cob weight without husk(g)</b>	<b>Cob yield with husk (t/ha)</b>	<b>Cob yield without husk (t/ha)</b>
1.	Azotobacter 20g/kg + Gibberellic acid 50 ppm	1.95	15.98	42.42	10.31	7.50	3.64
2.	Azotobacter 20g/kg + Gibberellic acid 100 ppm	2.25	16.64	45.24	11.26	8.59	3.56
3.	Azotobacter 20g/kg + Gibberellic acid 150 ppm	2.60	18.43	45.61	12.34	10.10	3.83
4.	Azospirillum 20g/kg + Gibberellic acid 50 ppm	1.81	14.61	38.81	9.86	7.18	3.54
5.	Azospirillum 20g/kg + Gibberellic acid 100 ppm	2.30	15.84	40.83	10.99	7.97	3.62
6.	Azospirillum 20g/kg + Gibberellic acid 150 ppm	2.48	16.88	44.03	11.51	8.73	3.70
7.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 50 ppm	2.35	17.45	44.15	11.75	9.44	3.83
8.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 100 ppm	2.43	18.65	46.69	12.27	10.56	4.03
9.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm	2.64	19.76	48.54	12.51	11.04	4.15
	<b>F-test</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
	SEm(±)	0.20	0.48	0.67	0.11	0.17	0.11
	CD (p=0.05)	0.07	1.47	2.01	0.32	0.51	0.34

**Table 3. Economics of Baby Corn as influenced by bio fertilizers and Gibberellic Acid.**

<b>S. No.</b>	<b>Treatments</b>	<b>Total cost of cultivation (INR/ha)</b>	<b>Gross return (INR/ha)</b>	<b>Net return (INR/ha)</b>	<b>B:C ratio</b>
1.	Azotobacter 20g/kg + Gibberellic acid 50 ppm	37,165.00	86,363.00	49,198.00	1.32
2.	Azotobacter 20g/kg + Gibberellic acid 100 ppm	37,265.00	95,279.00	58,014.00	1.56
3.	Azotobacter 20g/kg + Gibberellic acid 150 ppm	37,315.00	1,02,508.00	65,193.00	1.75
4.	Azospirillum 20g/kg + Gibberellic acid 50 ppm	37,215.00	84,603.00	47,388.00	1.27
5.	Azospirillum 20g/kg + Gibberellic acid 100 ppm	37,315.00	90,053.00	52,738.00	1.41
6.	Azospirillum 20g/kg + Gibberellic acid 150 ppm	37,365.00	98,379.00	61,014.00	1.63
7.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 50 ppm	37,315.00	1,00,025.00	62,710.00	1.68
8.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 100 ppm	37,415.00	1,05,173.00	67,758.00	1.81
9.	Azotobacter 10g/kg + Azospirillum 10g/kg + Gibberellic acid 150 ppm	37,465.00	1,08,252	70,787.00	1.89