

Original Research Article

Influence of Varieties and Foliar spray of Zinc on Growth , Yield and Economics of Rice (*Oryza sativa* L.)

ABSTRACT

A field experiment was conducted during *Kharif* 2022 at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj (U.P) to determine the “Influence of Varieties and Foliar spray of Zinc on Growth , Yield and Economics of Rice (*Oryza sativa* L.)”. The results showed that treatment 9 [BPT5204+ Zinc(2.5%)] recorded significantly higher plant height (113.77 cm), , maximum number of tillers/hill (12.80), higher plant dry weight(g) (54.13),maximum number of effective tillers/hill (12.50) ,maximum number of panicles/plant (10.20),maximum number of grains/panicle (104.70), higher test weight (18.70 g), higher grain yield (5.81 t/ha), Maximum straw yield (5.92 t/ha) and higher harvest index (50.19 %) compare to other treatment. The maximum gross returns (1,62,773.00 INR/ha), maximum net return (1,10,546.50 INR/ha) and highest benefit cost ratio (2.12) was recorded in treatment 9 [BPT5204+ Zinc(2.5%)] as compared to other treatments.

Keywords: *Rice, Varieties, Zinc, Growth, Yield and Economics.*

Introduction

Rice (*Oryza sativa* L.) is the most consumed cereal grain in the world, constituting the dietary staple food of more than half of the world population and referred to as “Global grain”. The theme on rice as “Rice is life” was used for international year of rice- 2004, denoting its overwhelming importance as an item of food and commerce by the United Nations. Globally, 1.4 billion ha of cultivable land is available. In rice growing countries, India has largest area under rice in the world (**Kiran et al.2019**).It is tropical plant ,it flourishes comfortably in hot and humid climate. It is fundamentally a kharif crop in India. It demands temperature of around 25 degree Celsius and above and rainfall of more than 100cm. it is also grown through irrigation in those areas that receives comparatively less rainfall. In India, states located in the Eastern and Southern parts cultivate the majority of rice. West Bengal is the leading rice producer in India which is followed by Uttar Pradesh, Telangana, Andhra, Punjab, Orissa, Bihar, Chhattisgarh, Tamil Nadu, Assam and Haryana (**Singh et al.2020**) . The starch content of the rice plant is generally low during the early

growth stages and increases toward flowering. Generally, before flowering, the starch accumulates in the leaf sheath and culm, but after flowering, it accumulates in the panicle. Much of the starch accumulated in the leaf sheath and culm before flowering is translocated into the grains during ripening. Carbohydrates produced by photosynthetic tissues is either transported to other organs as soluble sugars, or accumulated in leaves as soluble sugars and starch during the different growth stages. Soluble carbohydrates and starch, which accumulates under normal conditions before the stress commonly, constitute the main resources for plants to supply energy during stress condition **(Debbarma and Abraham ,2015)**

Rice ranks first in area and second in terms of production after china and is grown in all most tropical and sub humid regions of different countries .Globally, rice covers an area about 878 Million hectares with production of 898 Million tonnes and the productivity of 3348 kg/ha (**USDA ,2022**). In India , rice grown cover an area about 43.78 Million hectares with production of 118.43 Million tonnes and productivity of 2708 kg/ha under 2019-2020 , During 2019-2020 total area coverage under rice in Uttar pradesh 5.74 lakh hectare with the production of 13.11 lakh tonnes and productivity of 2704kg/ha (**GOI ,2020**). According to government third advance estimates rice production in 2020-2021 is 121.46 Million tonnes (**GOI ,2021**).

zinc is the most commonly deficient micronutrient in agricultural soils. Analysis of over 256,000 soil samples from all over India showed that about 50% of the soils were deficient in zinc and it causes leaf bronzing and poor tillering at the early growth stages, leading to delayed maturity and significant yield loss , its main cause of deficiency of plant available Zn in soil is the precipitation or adsorption of Zn with various soil components, depending on the pH and redox potential (**Zinzala and Narwade ,2019**). Zinc is one of the most essential micronutrients required for the growth and development of plant and human beings. One-third of the human population, including children and women suffer from Zn deficiency related health problems such as growth retardation, lack of appetite, lack of immune function, hair loss, diarrhoea, vision, and skin lesions, weight loss, delayed healing of wounds, and mental lethargy , The lack of micronutrients has become the major nutritional problem affecting more than two billion people in both developed and developing countries of Asia, Africa and Latin America. Micronutrient deficiencies or "hidden hunger" affects approximately 38% of pregnant women and 43% of preschool children worldwide and the most widespread among developing countries (**Jatav and Singh ,2018**).

RNR 15048 (Telangana Sona) It is widely recognized for its super fine quality (short slender) of good cooking quality with grain yield of 6.5-7.0t/ha and high milling recovery (68-70%). *Kharif* Duration:110-120 days ,*Rabi* :125-130 days. It is said to be resistant to blast and tolerant to BPH

nearly saving Rs. 2000 to 3000 per acre expenditure on plant protection.

KNM 118 (Kunaram sannalu) The rice culture, KNM 118 is a good alternate variety to mega rice variety, MTU 1010 as it exhibits high yield potential (7-8 t/ha) with good test weight, and is less prone to grain shattering and lodging at the time of harvest when compared to MTU 1010. Plant height:100-106 cm. Duration: *Kharif* :120-125 days , *Rabi* :125-130 days. It is suitable for *kharif*, late *kharif* and *rabi* seasons. This culture has tolerance to leaf blast and neck blast.

BPT 5204 (Samba Masuri) The grain of samba masuri is of medium slender type and is semi dwarf variety suitable for rainfed shallow low lands of our country. Crop duration is 135-140 days and is suitable for *kharif* and *Rabi* seasons. The yield potential of BPT 5204 is 5.5 -6.0 t/ha. This variety is resistant to neck blast and tolerant to gall midge and is non-lodging and non-shattering type.

Zinc is an essential micro nutrient for crop plants. Zn is essential for several biochemical processes in rice plant, such as cytochrome and nucleotide synthesis, auxin metabolism, chlorophyll production, enzyme activation and membrane integrity. Zn- enrichment leads to more root surface area and the ability to change chemistry and biology of rhizosphere by releasing phytosiderophores from roots which ultimately increases Zn uptake by plants (Suvarna *et al* .2015). zinc is major component and activator of several enzymes involved in metabolic activities , its deficiency continues to be one of the key factors in determining rice production in several parts of country (Muthukumararaja and Sriramachandrasekharan 2012) . Foliar application of zinc usually applied under emergencies to save the crops to symptoms of Zn appear ,single foliar application may not be adequate to severe deficiency symptoms , foliar application of 0.5% aqueous solution of $ZnSO_4$ twice at 20 and 30 days after transplanting (Verma *et al*.2019) . Keeping in view the above fact, the experiment was conducted to find out Influence of Varieties and Foliar spray of Zinc on Growth , Yield and Economics of Rice “(*Oryza sativa* L.)”

MATERIALS AND METHODS:

The experiment was conducted during *Kharif* season of 2022 at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj (U.P). The soil of the field constituting a part of central gangetic alluvium is neutral and deep. The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH 7.8), low level of organic carbon (0.62%), available N (225 Kg/ha), P (38.2kg/ha) and K (240.7 kg/ha). The treatment consists of 3 different Zinc *viz.* (0.5%), (1.5%) and (2.5%) with combination of 3 Varieties *viz.* BPT5204, RNR15048, and KNM118. The experiment was laid out in RBD with 9 treatments each replicated thrice. The treatment combinations are Treatment 1[RNR 15048 + zinc (0.5%)], Treatment 2 [RNR 15048 + zinc(1.5%)], Treatment 3[RNR 15048 + zinc (2.5%)], Treatment 4[KNM 118 + zinc (0.5%)], Treatment 5 [KNM 118 + zinc (1.5%)], Treatment 6 [KNM 118 + zinc (2.5%)], Treatment 7[BPT 5204 + zinc (0.5%)], Treatment 8 [BPT 5204 + zinc (1.5%)],

T [BPT 5204 + zinc (2.5%)]. Data recorded on different aspects of crop , viz growth , yield attributes and yield were subjected to statistical analysis by analysis of variance method (**Gomez and Gomez , 1976**)

RESULT AND DISSCUSSION

Growth parameters

Plant height (cm):

The data revealed that significantly highest plant height (113.77 cm) was recorded in the treatment 9[BPT5204 + (Zinc 2.5%)] (Table 1) . Significant and higher plant height was recorded with the variety (BPT5204) might be due to variation in growth characters of varieties to genetic makeup, its adaptation to the environment. Similar results were reported by **Kiran et al. (2019)**. Further, in higher plant height with application of zinc (2.5%) may due to adequate supply of zinc which contributed to accelerate enzymatic activity and auxin metabolism in plants. Similar results Were reported by **Khan et al .(2007)**.

Number of tillers/hill:

The results revealed that significant and Maximum number of tillers/hill (12.80) was recorded in the treatment 9 [BPT5204 + Zinc(2.5%)]. However, treatment 4 [KNM118 + (Zinc 0.5 %)] was found to be statistically at par with treatment 9 [BPT5204 + (Zinc 2.5%)](Table 1). The significant and Maximum number of tillers / hill was observed with the variety (BPT5204) might be due to genetic character, The difference in growth attributes indicate that all rice varieties respond differently to the applied nutrient such as nutrient uptake and nutrient use efficiency. similar results were reported by **Pratap et al. (2020)**.Further, increased in tiller/hill with application of zinc (2.5%) may due to its application attributed to its role in various enzymatic activity and auxin metabolism which control growth of plant. Similar results were reported by **Singh et al. (2020)**.

Plant dry weight (g):

Data found that significant and higher plant dry weight (54.13 g) was recorded in the treatment 5 [KNM 118 +(Zinc 1.5%)]. However, treatment 8 [BPT 5204+(Zinc 1.5%)] was found to be statistically at par with treatment 5 [KNM 118 +(Zinc 1.5%)](Table 1). The Significant and higher plant dry weight was observed with the variety KNM118 might be due to their genetic potential, capacity towards uptake of nutrients by varieties, competition

of variety for space, light, nutrients and differential plant height. Similar results was also reported by **Patil et al. (2022)**. Further increased in plant dry weight with the application of zinc (2.5%) may due to important role of zinc for the activation of various types of enzymes, such as those required for the CO_2 assimilation pathway and chlorophyll biosynthesis. Similar results were reported by **Suvarna et al. (2015)**.

Crop Growth Rate ($\text{g/m}^2/\text{day}$):

The data recorded that significant and higher crop growth rate ($7.95 \text{ g/m}^2/\text{day}$) was recorded in treatment 7 [BPT5204 +(Zinc 0.5%)]. However, treatment 4 [KNM 118 +(Zinc 0.5%)] was found to be statistically at par with treatment 9 [BPT5204 +(Zinc 2.5%)] (Table1). The significant and higher crop growth rate was recorded with application of zinc (2.5%) might be due to its function of combined effects of genetic make up of plant, soil nutrient status, seedling vigor and environmental conditions under which it is grown, resulted increased in crop growth rate of the plant. Similar results were reported by **Mustafa et al (2011)**.

Yield Attributes & Yield

Number of effective tillers/hill :

The data showed that Significant and maximum number of effective tillers/hill (12.50) was recorded in the treatment 9 [BPT5204 + zinc (2.5%)] However, treatment 4 [KNM118 + (zinc 0.5%)] was found to be statistically at par with treatment 9 [BPT5204 + (zinc 2.5%)] (Table 2). The significant and Maximum number of effective tillers/hill was recorded with the variety (BPT5204) might be due to varietal character and adequate availability of photosynthates under transplanted system of cultivation, good maintenance of source sink relationship due to longer reproductive phase. Similar results were also reported by **Maniraj et al .(2022)**. Further, increased in effective tillers/hill with application of zinc (2.5%) may due to better utilization of applied zinc owing to developed root system. With the sufficient nutrition of zinc growth promoting substance like auxin in plant is produced and that may increase the overall growth of the plant. Similar results were reported by **Sardar et al.(2020)**.

Number of panicles/plant:

The data recorded that significant and Maximum number of panicles/plant (10.20)

was recorded in the treatment 9 [BPT5204 + zinc (2.5%)] However , treatment 8 [BPT5204 + (zinc 1.5%)] was found to be statistically at par with treatment 9 [BPT5204 +(zinc 2.5)] (Table 2).The significant and maximum number of panicles/plant was recorded with the variety (BPT5204) might be due to increased accumulation of photosynthetic from the source to the sink and the differences in growth characters due to genotypes may be attributed to their inherent characteristics, resulted increased in panicles/ plant. Similar results were reported by **Dangi et al. (2017)**. Further increased in panicles/plant with application of zinc (2.5%) may due to Participation of Zn in biosynthesis of indole acetic acid (IAA) and its role in initiation of primordial reproductive parts and partitioning of photosynthates towards them are responsible for increased in panicles/plant. Similar results were reported by **Ram et al .(2013)**.

Number of grains/panicle:

The data showed that significantly highest number of grains /panicle (104.70) was recorded in the treatment 9 [BPT5204 + zinc (2.5%)] (Table2). The significant and Maximum number of grains/panicle was recorded with the variety (BPT5204) might be due to Efficient utilization of growth resources, less intra species competition coupled with higher availability of nutrients among the widely spaced crop plants may be ascribed the reason for superiority in increasing of grains/panicle. Similar results were reported by **Rahem and Marzoka , (2020)**. Further increased in number of grains/panicle with application of zinc (2.5%) may due to it is synthesizer of protein and carbohydrate resulted into bolder seeds and increased more number of grains/panicle. Similar results were reported by **Singh et al.(2018)**.

Test weight (g):

Significant and higher test weight (18.70 g) was recorded in the treatment 9 [BPT5204 + zinc (2.5%)] However , treatment 7 [BPT5204+ (zinc 0.5%)] was found to be statistically at par with treatment 9 [BPT5204+(Zinc 2.5%)] (Table 2). The significant and higher test weight was recorded with the variety (BPT5204) might be due to better growth and translocation of photosynthates to reproductive parts, which might have increased grain weight of the plant. Similar results were reported by **Singh et al (2017)**. Further increased in test weight (g) with application of zinc (2.5 %) may be due to it is involved in several physiological processes including protein synthesis, enzyme activation, carbohydrates metabolism, auxins, lipids, and nucleic acids , resulted increased in test weight . Similar results were reported by **Sayed et al. (2022)**.

Grain Yield (t/ha):

The data revealed that significantly highest grain yield (5.81 t/ha) was recorded in the treatment 9 [BPT5204 + zinc (2.5%)] (Table 2). The significant and higher grain yield was recorded with the variety (BPT5204) might be due to greater vegetative growth and better light interception and higher dry matter partitioning towards economic part, yield variability among rice cultivars also attributed to genetic characters and environmental effects. Similar results were reported by **Kumar et al. (2017)**. Further increased in grain yield with application of zinc (2.5%) may due to fertilization is attributed to its involvement in many metabolic enzyme system, regulatory functions and auxin production enhanced synthesis of carbohydrates and their transport to the site of grain production. Similar results were reported by **Muthukumararaja et al (2012)**.

Straw Yield (t/ha):

The data showed that significantly Maximum straw yield (5.92 t/ha) was recorded in the treatment 9 [BPT5204 + zinc (2.5%)] (Table 2). The significant and Maximum straw yield was recorded with the variety (BPT5204) might be due to higher photosynthetic rate as a result of enhanced LAI, which in turn boosted dry matter formation, resulting in higher grain yield and straw yields. Similar results were reported by **Maurya et al. (2021)**. Further increased in straw yield with application of zinc (2.5%) may due to it is attributed to adequate supply of zinc that might have increased the availability and uptake of other essential nutrients resulting in improvement in metabolic activities and also due to the effect of zinc on the proliferation of roots. Similar results were reported by **Zinzala et al. (2019)**.

Harvest Index (%):

Data recorded that significant and higher harvest index (50.19 %) was recorded in treatment 9 [BPT5204 + zinc (2.5)]. However, treatment 6 [KNM118 + (zinc 2.5%)] was found to be statistically at par with treatment 9 [BPT5204 +(zinc 2.5%)] (Table 2). The significant and higher harvest index was recorded with the variety (BPT5204) might be due to adequate and available water at reproductive stages where irrigation system is flooded until ripening stage, resulted increased in harvest index. Similar results were reported by **Soriano et al.(2017)**. Further increased in harvest index with application of zinc (2.5) may be due to it is indicated the greater translocation of photosynthates from source to sink and better partitioning towards reproductive growth. Similar results were reported by **Saikh et al. (2022)**.

ECONOMIC ANALYSIS

Economics:

The result showed that maximum gross returns (1,63,120.00 INR/ha), Maximum net return (1,10,893.50 INR/ha) and highest benefit cost ratio (2.12) was recorded in treatment 9 [BPT5204 + zinc (2.5%)]. As compared to other treatments (Table 3). Maximum gross returns, net returns and benefit cost ratio was recorded with the application of zinc (2.5%) might be due to it is essential plant nutrient and its involvement in the physiological process is well pronounced , therefore increase in both grain and straw yield and ultimately increased benefit cost ratio. Similar results were reported by **Bareddy *et al.*(2020)**

CONCLUSION:

Based on above findings it can be concluded that combination of BPT5204 alongwith Zinc (2.5%) has performed better in growth parameters and yield attributes of rice and also proven profitable .

REFERENCES:

1. **Achin Kumar and AP Singh (2018)** . Direct and residual effect of zinc and boron on growth parameters of rice and wheat grown in sequence in red and alluvial soils of eastern Uttar Pradesh . International Journal of Chemical Studies. **6**(1): 587-592
2. **Arun Pratap Singh, Ashok Rai, Vikas Singh and AK Dube (2020)**. Comparative study of growth and yield attributes of different paddy cultivars under Tarai region of Uttar Pradesh . International Journal of Chemical Studies . **8**(2): 129-132
3. **B Vibhajam Sagal Kiran, VRK Murthy, M Sree Rekha, Y Ashoka Rani and PRK Prasad (2019)** . Growth, yield and nitrogen uptake of direct seeded rice varieties under variable nitrogen rates . International Journal of Chemical Studies . **7**(4): 198-201
4. **Bareddy Narendranath Reddy, C. Umesh, Pole Shiva kiran and Dhanush Reddy. (2020)**. Effect of Zinc and Boron Levels on Growth, Yield and Economics of Rice (*Oryza sativa* L.) var. Shiats Dhan-1. International Journal of Current Microbiology and Applied Sciences.**9**(12):2319-7706.
5. **D.K. Singh, Shilpi Gupta, Gangadhar Nanda, Yogesh Sharma, V.V. Singh and Dipti Bisarya (2017)** . Evaluation of Rice Varieties for Yield under Organic Farming in Tarai Region of Uttarakhand, India . International Journal of Current Microbiology and Applied Sciences . **6** (4) :2319-7706
6. **GOI , (2020)** . Agricultural statistics at a Glance , Agricultural statistics division , Directorate of Economics and statistics , Department of Agriculture and cooperation Ministry of Agriculture.
7. **Gomez , K, A and Gomez ,A .A(1976)** . Three or more factor experiment .in : Statistical Procedure for agricultural Research 2nd edition.P.139-141
8. **Junel B. Soriano1, Suhas P. Wani, Adusumilli N. Rao, Gajanan L. Sawargaonkar, and Jnanasha A.C. Gowda (2018)** . Comparative Evaluation of Direct Dry-Seeded and Transplanted Rice in the Dry Zone of Karnataka, India . **147** (1) : 0031 - 7683
9. **Kuldeep Dangi , S.K. Singh, D.K. Malviya1 , A. Gautam , N. Kanapuriya and Balvendra Kumar (2017)** . Effect of Rice Varieties on Growth, Yield and Economics at Varying Levels of Nitrogen under Direct Seeded Upland Condition Rewa Region .**6** (9): 2319-7706
10. **Muthukumararaja, T.M . and M.V. Sriramachandrasekharan (2012)** . Effect of zinc on yield, zinc nutrition and zinc use efficiency of lowland rice . Journal of Agricultural Technology . **8**(2): 551-561
11. **M. Umar Khan, M. Qasim and Israrullah Khan (2007)** . Effect of zn fertilizer on rice grown in different soils of dera ismail khan . Sarhad J. Agric. **23**(4)

12. **N. Maniraj, P. Revathi , K.B. Suneetha Devi and K. Chandra (2022)** . Growth and Yield Attributes of Rice as Influenced by Systems of Cultivation in different Varieties . Biological Forum – An International Journal . **14(2)** : 2249-323
13. **Praneeth kumar suvarna, s. Narender reddy, a. Siva sankar and y. Sharath kumar reddy (2015)**. Effect of zinc nutrition on dry matter production, yield and uptake of zinc by aromatic rice .The ecoscan . **9(1&2)**: 513-516
14. **Rahul saikh, kanu murmu , arup sarkar, ramyajit mondal, kalyan jana (2022)**. Effect of foliar zinc application on growth and yield of rice (Oryza sativa) in the Indo Gangetic Plains of India . *N u s a n t a r a b i o s c i e n c e* . **14(12)** :2087-3948
15. **Rajesh, S R B K Desai , Satyanarayana Rao ,B M Chittapur , K Narayana Rao , K T Vendan ,Benoit Clerget ,Tegegnetwork Gebremedhin (2016)** . Growth Yield and Nutrient Status of Rice Soil in Response to Genotypes and Levels of Nitrogen Under Different Cultivation Techniques . Journal of Biology, Agriculture and Healthcare **6(17)** 2224 3208
16. **Rakesh Maurya , Manoj Kumar Singh , Nikhil Kumar Singh , Manish Kumar Singh , Anurag Kumar Singh (2021)** . Effect of nitrogen levels on growth attributes, yield and nutrient uptake of different rice (oryza sativa l.) Varieties under the transplanted condition. Journal of Experimental Biology and Agricultural Sciences **9(3)** : 2320 – 8694
17. **S. Vikas Patil , G. Santhosh Kumar , U. Nagabhushanam , CH. Ramulu and M. Yakadri (2022)**. Evaluation of Rice Varieties under Different Systems of Cultivation for Central Telangana Zone .**12 (12)** : 2581-8627
18. **Suhail Sardar, UP Shahi, BP Dhyani, Ashok Kumar Ahluvalia, Ashwani K Tyagi and Waseem Ahma)2020)** . Effect of different sources and application mode of zinc on growth, yield attributing characters and yield of rice variety Pusa basmati 1121 in sandy loam soil . International Journal of Chemical Studies **8(6)** : 2321–4902
19. **Surendra Singh Jatav and Satish Kumar Singh (2018)**. Evaluation of different methods of Zinc application on growth and yield of hybrid rice (Oryza sativa L.) in inceptisol of Varanasi. Journal of Pharmacognosy and Phytochemistry **7(4)**: 2349-8234
20. **Sandeep Kumar , Sarabdeep Kour , Shivani Lalotra , Kartikeya Choudhary and Hari Singh(2017)** . Response of Different Rice Varieties and Fertility Levels on Relative Economics and Quality of Rice under Aerobic Conditions . Current Journal of Applied Science and Technology **23(6)** : 2231-0843
21. **USDA ,(2022)** . National Agricultural statistics Service .www.quickstats.nass.usda.gov
22. **Uma Shanker Ram, VK Srivastava, A Hemantaranjan, A Sen, RK Singh, JS Bohra and U Shukla (2013)** . Effect of Zn, Fe and FYM application on growth, yield and nutrient content

- 23. Veer Singh, Virendra Singh , Satybhan Singh and Richa Khanna (2020) .** Effect of Zinc and Silicon on Growth and Yield of Aromatic Rice (*Oryza Sativa*) in North-Western Plains of India . *Journal of Rice Research and Developments* **3**(1) : 2643-5705
- 24. V. K. Verma, deepak kumar verma, and ram kumar singh (2019).** Effect of zinc (zn) on growth, yield, and quality attributes of rice (*oryza sativa* l.) For improved rice production. Apple Academic Press, Inc. **13**: 978-0-429-48858-0
- 25. Victor Debbarma and Thomas Abraham (2015) .** Quality of organic rice [*Oryza sativa* (L.) sub sp. japonica] as influenced by planting methods and nutrient management .*New Agriculturist*, **26**(2) : 341–346
- 26. Vivek N. Zinzala and Ajay V. Narwade (2019).** Effect of Zinc applications on Grain Yield, Straw Yield and Harvest Index in kharif Rice (*Oryza sativa* L.) Genotypes . *International Journal of Current Microbiology and Applied Sciences* **8**(11) : 2319-7706
- 27. Waikhom Jiten Singh¹, Mahua Banerjee¹ and L. Nabachandra Singh (2018) .** Effect of Sulphur and Zinc on Yield Attributes, Yield and Economics of Rice . *International Journal of Current Microbiology and Applied Sciences* **7**(3) : 2319-7706
- 28. W.T. Abd El-Rahem and E.A. Abo-Marzoka (2020) .** Differantral response of some rice cultivars to seedling age and plant spacing and their effect on growth traits and yield. *Egypt. J. Plant Breed.* **24**(3):599– 616Ms_IJPSS_100044

UNDER PEER REVIEW

Table 1. Influence of Varieties and Foliar spray of Zinc on on growth parameters of rice.

Sl. No.	Treatments	Plant heights (cm)	Number of tillers/hill	Plant dry weight (g)	Crop growth Rate (g/m ² /day)	Relative growth rate (g/g/day)
		AT 100 DAT	AT 100 DAT	AT 100 DAT	80 – 100 DAT	80 – 100 DAT
1	RNR 15048 +Zinc 0.5%	106.07	11.50	47.95	6.05	0.0017
2	RNR 15048 +Zinc 1.5%	106.17	10.80	49.06	7.27	0.0020
3	RNR 15048 +Zinc 2.5%	106.97	11.20	48.69	7.78	0.0021
4	KNM118 +Zinc 0.5%	104.20	12.23	50.51	6.77	0.0018
5	KNM 118 +Zinc 1.5%	103.77	11.60	54.13	2.63	0.0006
6	KNM 118 +Zinc 2.5%	107.83	12.30	50.03	6.73	0.0018
7	BPT 5204 +Zinc 0.5%	108.70	12.12	49.30	7.95	0.0022
8	BPT 5204+Zinc 1.5%	110.77	12.56	53.51	6.17	0.0015
9	BPT 5204+Zinc 2.5%	113.77	12.80	53.14	5.12	0.0012
	F test	S	S	S	S	S
	SE(m)±	0.85	0.80	0.07	0.16	0.0038
	CD (p=0.05)	2.54	1.18	0.22	0.50	0.0012

Table 2. Influence of Varieties and Foliar spray of Zinc on yield attributes and yield of rice.

Sl No.	Treatments	Number of effective tillers/hill	Number of panicles/plant	Number of grains/panicle	Test weight (g)	Grain yield (t/ha)	Straw yield (t/ha)	Harvest Index (%)
1	RNR 15048 +Zinc 0.5%	11.0	8.00	96.50	13.77	4.46	5.34	45.48
2	RNR 15048 +Zinc 1.5%	10.29	8.67	97.50	13.60	4.56	5.44	45.63
3	RNR 15048 +Zinc 2.5%	10.12	8.67	96.17	13.73	5.11	5.47	48.30
4	KNM118 +Zinc 0.5%	11.54	7.67	94.73	15.30	4.85	5.49	46.94
5	KNM 118 +Zinc 1.5%	11.27	8.67	98.70	15.57	5.52	5.52	49.98
6	KNM 118 +Zinc 2.5%	10.57	8.33	98.47	15.60	5.59	5.65	49.73
7	BPT 5204 +Zinc 0.5%	11.58	8.78	99.10	18.60	5.67	5.63	49.55
8	BPT 5204+Zinc 1.5%	12.0	9.34	101.8	18.63	5.69	5.67	50.10
9	BPT 5204+Zinc 2.5%	12.50	10.20	104.70	18.70	5.81	5.92	50.19
	F test	S	S	S	S	S	S	S
	SE(m)±	0.45	0.36	0.26	0.14	0.07	0.14	0.27
	CD (p=0.05)	1.08	1.06	0.78	0.40	0.24	0.13	0.79

Table 3. Influence of Varieties and Foliar spray of Zinc on the economics of rice.

Sl.NO	Treatment combinations	Economics			
		Cost of cultivation	Gross return	Net return	B:C Ratio
1.	RNR 15048 +Zinc 0.5%	51,727.50	1,24,787	73,059	1.41
2.	RNR 15048 +Zinc 1.5%	51,727.50	1,27,773	76,046	1.47
3.	RNR 15048 +Zinc 2.5%	51,977.50	1,43,080	91,353	1.77
4.	KNM 118 +Zinc 0.5%	51,977.50	1,35,893	83,916	1.61
5.	KNM 118 +Zinc 1.5%	51,977.50	1,54,467	1,02,489	1.97
6.	KNM 118 +Zinc 2.5%	52,227.50	1,56,520	1,04,543	2.01
7.	BPT 5204 +Zinc 0.5%	52,227.50	1,58,760	1,06,533	2.04
8.	BPT 5204+Zinc 1.5%	52,227.50	1,59,320	1,07,093	2.05
9.	BPT 5204+Zinc 2.5%	51,707.50	1,62,773	1,10,546	2.12