

Original Research Article

The influence of vitamins, antibiotics and growth stimulators on L-lysine Production by *Bacillus subtilis* using Agricultural Products as Carbon and Nitrogen Sources

Abstract

L-lysine is one of the 9 amino acids which are essential for human and animal nutrition. It is a basic building block of all protein and cannot be synthesized biologically in the body. The influence of vitamins, antibiotics and growth stimulators on L-lysine production by *Bacillus subtilis* using agricultural products as carbon and nitrogen sources was examined. The L-lysine producing bacteria had already been isolated from Nigerian soil. They were purified and Identified as *B. subtilis* PR13 and *B. subtilis* PR9, using cultural, biochemical and molecular characteristics. The study was conducted to enhance lysine production by *Bacillus subtilis* through the evaluation of influence of vitamins, antibiotics and growth stimulators. The L-lysine was produced in 100 ml flasks containing 20 ml fermentation media (FM1 and FM2). The results obtained showed that, an enhanced lysine yield of 3.41 and 1.57 mg/ml by *Bacillus subtilis* PR13 and *Bacillus subtilis* PR9, was observed at in the presence of 1 µg/ml of biotin and 10 µg/ml respectively. The supplementation of 0.04 units/ml of penicillin, enhanced optimum L-lysine yield of 2.38 mg/ml for *B. subtilis* PR 13 and 1.64 mg/ml for *B. subtilis* PR9 respectively. the addition of 0.1% w/v peptone and yeast extract, enhanced optimum L-lysine yield of 2.66 mg/ml for *B. subtilis* PR 13 and 1.72 mg/ml for *B. subtilis* PR9 respectively. There was a positive correlation between peptone and lysine production by *B. subtilis* PR13 ($r= 0.85$) and yeast extract and lysine production by *B. subtilis* PR9 ($r= 0.54$). The results obtained in the study showed that the

Comment [D1]: Abstract should be in format provided by the guideline.

supplementation of vitamins, antibiotics and growth stimulators, could increase the L-lysine yield of by *B. subtilis* PR13 and *B. subtilis* PR9.

Keywords: *Bacillus subtilis*, L-lysine, Submerged fermentation, Vitamins, Antibiotics

1. Introduction

Amino acids are major industrial products derived by fermentation, covering a world market of more than 5 million tons per year, and among amino acids are the L-lysine that is one of the leading biotechnological products with a current production of 2.2 tons per year [1].

L-Lysine, is an amino acid primarily essential for human and animal nutrition, and is usually obtained in batch or fed-batch fermentation processes. Of that manufactured commercially, the largest amount of 80% is produced by fermentation and 20% by chemical synthesis [2]. Lysine is used as food (flavour) enhancer and also food preservation especially with ϵ -poly-L-lysine [3]. It is used as food supplements for humans (children have a high requirement of lysine, since it is needed for bone formation) [4, 5]. It is used to enrich feed stuff in order to provide an adequate diet for poultry, cattle and other live stock. Animal feed such as grain and defatted oil seeds, contain only a small quantity of lysine [6].

The use of L- Lysine is helpful to overcome angina pectoris. It is an essential ingredient used to clean arteries, important for cancer prevention [7]. Lysine supports bone health by ensuring adequate absorption of calcium and therefore prevents osteoporosis. It has an important role in production of antibodies for healthy immune system. It is the integral component of musculature [8]. In addition, it has pharmaceutical applications both in the formulation of diets with a balanced amino acid composition and in the infusion of amino acids [9]

A great variety of microorganisms have been discovered to produce L-lysine on industrial scale. They include *Corynebacterium glutamicum*, *Bacillus subtilis*, *Bacillus megaterium*, *Escherichia coli*, species of *Arthrobacter*, *Brevibacterium* and *Micrococcus* [10,11,12,13].

Extensive research has been made in order to improve the fermentation process not only from the point of lowering production costs but also of increasing productivity. Improvements have included for example, increased yield of desired metabolites, removal of unwanted co-metabolites, improved utilization of inexpensive carbon and nitrogen sources, or alteration of the morphology to a form better suited for separation of the organisms from the product [14].

Utilization of agricultural by-products as substrates for fermentation might offer an inexpensive alternative for microbial products such as amino acids [15,16,17].

Currently, Nigeria meets all her L-lysine needs only through importation. However, L-lysine can be made available and more economical if produced locally by fermentation using agricultural products. A huge amount of foreign exchange will be saved and this development will impact positively on the economy.

We had isolated three *Bacillus* species (which included *Bacillus subtilis* PR13, *Bacillus subtilis* PR9, and *Bacillus p umilus* SS16) from Nigerian soil, which produced various yields of L-lysine [18]. In another study, the *Bacillus* species were used for L-lysine production using carbohydrates as carbon and seed meals as nitrogen sources [19].

2.0 Materials and Methods

2.1 Microorganisms and culture maintenance conditions

B. subtilis PR13 and *B. subtilis* PR9 isolated from different soil [18] in Awka town, were used in the study. They were purified and Identified as *B. subtilis*

Comment [D2]: Her or their?

Comment [D3]: Is another study means citation number 19?

Comment [D4]: It would be better if citation number would be in sequence wise.

PR13 and *B. subtilis* PR9, using cultural, biochemical and molecular characteristics. The *Bacillus subtilis* were grown on nutrient agar slants for 24 h at 37°C. Thereafter, the cultures were then preserved at 4°C and transferring to new slants after 30 days in order to keep them viable for use in L-lysine production.

2.2 Seed culture preparation

The seed medium consisted of peptone, 10.0g; yeast extract, 10.0 g; NaCl, 5.0 g; water, 1litre; pH adjusted to 7.2. Two loopfuls of *B. subtilis* PR13 and PR9 were inoculated in an Erlenmeyer flask containing 50 ml of seed medium which had already been sterilized at 121 °C for 15 min. The inoculated flasks were incubated for 24 h on a rotary shaker at 120 rpm and 30 °C. Duplicate flasks were used.

Comment [D5]: Reference?

2.3 Fermentation Media Preparation

The submerged production of L-lysine by *Bacillus subtilis* PR13 and PR9, was conducted in two fermentation media namely fermentation medium 1 and 2 (FM1 and FM2). For *Bacillus subtilis* PR 13, L-lysine production was carried out in 100 ml Erlenmeyer flasks, containing 20ml of fermentation medium 1 (FM1). The fermentation medium 1, was composed of KH_2PO_4 , 0.5g; K_2HPO_4 , 0.5g; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.001g; $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 0.001g; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.001g; CaCO_3 , 50g, the carbon source (glucose) was replaced with millet starch hydrolysate 60g; the nitrogen source (ammonium sulphate) was replaced by soyabean meal 40g; water, 1 litre; pH adjusted to 7.2. For *Bacillus subtilis* PR 9, fermentation medium (FM2) was used for L-lysine production. FM2 is similar to FM1, except that the carbon source (glucose) was replaced with sorghum hydrolysates 60g, the nitrogen source (ammonium sulphate) was replaced by deffated peanut meal 40g. The carbon source substrates were prepared in the laboratory using the method of Umerie *et al.* [20].

Comment [D6]: Reference?

Comment [D7]: Please provide full name along with molecular formula

Comment [D8]: Are you suggesting that whole sentence is covered by reference number 20?

2.4 Optimization of culture conditions for L-lysine production

2.4.1 Effect of vitamins

The effect of vitamins, which included various concentrations (0.01-100µg/ml) of riboflavin, thiamine, biotin, nicotinic acid and cyanocobalamin on lysine production was determined. Fermentation was carried out in 100ml Erlenmeyer flasks containing 20 ml of fermentation media (FM1 and FM2) as was previously described. The vitamins were added to the fermentation media and sterilized at 121°C for 15 min. One milliliter volume (1.8×10^7 cfu/ml) of 24h cultures of *Bacillus* species was inoculated into the fermentation media. Uninoculated flasks served as control. The flasks were placed on a rotary shaker (at 160 rpm) and incubated at 30°C for 72 h. Following the termination of fermentation, the culture broth was subjected to centrifugation at 5,000 rpm for 15 min to obtain the cell-free supernatant which is the crude L- lysine. The cell-free supernatant was used for the determination of lysine. The experiments were conducted in triplicate.

Comment [D9]: Reference for the process?

2.4.2 Effect of antibiotics

The effect of antibiotics ,which included varying concentrations (0.01 – 0.10units/ml) of penicillin and (0.01 – 2.00µg/ml) of vancomycin, erythromycin, lincomycin, tetracycline and chloramphenicol on lysine production was studied. Fermentation was carried out in 100ml Erlenmeyer flasks containing the 20 ml of fermentation media (FM1 and FM2) as was previously described. The antibiotics were added to the fermentation media and sterilized at 121°C for 15 min. One milliliter volume (1.8×10^7 cfu/ml) of 24h cultures of *Bacillus* species was inoculated into the fermentation media. Uninoculated flasks served as control. The flasks were placed on a rotary shaker (at 160 rpm) and incubated at 30°C for 72 h. Following the termination of fermentation, the culture broth was subjected to centrifugation at 5,000 rpm for

15 min to obtain the cell-free supernatant which is the crude L- lysine. The cell-free supernatant was used for the determination of lysine. The experiments were conducted in triplicate.

Comment [D10]: Reference for the process?

Effect of growth stimulators

Comment [D11]: Does not it should be under section or subsection number?

The effect of growth stimulators ,which included various growth stimulators which included 0.1% w/v of gelatine, yeast extract, peptone, tryptone, casein and beef extract was carried out. Fermentation was carried out in 100ml Erlenmeyer flasks containing 20 ml of fermentation media (FM1 and FM2), as was previously described. The growth stimulators were added to the fermentation media and sterilized at 121°C for 15 min. One milliliter volume (1.8×10^7 cfu/ml) of 24h cultures of *Bacillus* species was inoculated into the fermentation media. Uninoculated flasks served as control. The flasks were placed on a rotary shaker (at 160 rpm) and incubated at 30°C for 72 h. Following the termination of fermentation, samples of the fermentation medium were aseptically dispensed into cuvettes using micropipettes. Thereafter, the cuvettes were placed in the spectrophotometer and the reading for bacteria growth was determined at 660nm. For the determination of L-lysine and residual sugar, the fermentation medium was subjected to centrifugation at 5,000 rpm for 15 min to obtain the cell free supernatant which is the crude L-lysine. The cell free supernatant was used for the determination of lysine and residual sugar. The experiments were conducted in triplicate.

Comment [D12]: Reference for the process?

2.5 Quantitative determination of lysine

L-lysine in the broth culture was determined by acidic ninhydrin method of Chinard [21]. A 5ml volume of the culture broth of the isolate was centrifuged at $5000 \times g$ for 20min, and the cell-free supernatant was collected and assayed for lysine production. One milliliter (1ml) of glacial acetic acid was added to 1ml of supernatant in a test tube. Thereafter, one ml of a reagent solution which contains an acid mixture, 0.4ml of 6M orthophosphoric acid, 0.6ml of glacial

acetic acid and 25mg of ninhydrin, was also added to the supernatant in the test tube. The blank contains 1ml of glacial acetic acid, 1ml of the acid mixture without ninhydrin and 1ml supernatant. Both tubes were capped and the contents mixed properly for 10min before heating at 100 °C in a water bath for 1h. The test tubes were cooled rapidly under tap water and 2ml of glacial acetic acid was added to each test tube to give a final volume of 5ml. The optical density of the reacting mixture was read against the blank at 515nm in a spectrophotometer. Results obtained with the test samples were extrapolated from a standard lysine curve.

Comment [D13]: Reference for the process?

2.6 Estimation of reducing sugar

The reducing sugar content was determined by dinitrosalicylic acid (DNS) method of Miller [22]. Reducing sugar was estimated by adding 1ml of DNS to 1ml of the supernatant. The mixture was heated in a water bath at 100 °C for 10min and allowed to cool. The volume of the mixture was thereafter increased to 12 ml with distilled water. After allowing the reaction mixture to stand for 15min at room temperature, the optical density was measured at 540 nm in a spectrophotometer against a blank prepared by substituting the supernatant with water. The reducing sugar content was subsequently determined by making reference to a standard curve of known glucose concentrations.

Comment [D14]: Reference for the process?

2.7 Statistical analysis

Data generated from this work were analyzed using correlation analysis with a software application SPSS version 14.

Comment [D15]: Why old version of SPSS is used since new version are available? Please provide full form before providing abbreviation.

3.0 Results


The results of the effect of growth stimulators on growth and lysine production by *Bacillus subtilis* PR13 and *Bacillus subtilis* PR9 is shown figures 1 and 2. The highest L- lysine production of 2.66 and 1.72 mg/ml by *Bacillus subtilis* PR13 and *Bacillus subtilis* PR9 were observed at the supplementation of 0.1 %

Comment [D16]: It would be appropriate if description part would be along with figure e.g first figure and its description.


w/v of peptone and yeast extract respectively. The highest yield corresponded with a final reducing sugar of 0.75 for *B. subtilis* PR13 and 0.55 mg/ml for *B. subtilis* PR9. There was a positive correlation between peptone and lysine production by *B. subtilis* PR13 ($r= 0.85$) and yeast extract and lysine production by *B. subtilis* PR9 ($r= 0.54$).

The results of the effect of vitamins on lysine production by *B. subtilis* PR13 and PR 9 are presented in Figures 3-4. The results showed that maximum lysine yields of 3.41 and 1.57 mg/ml by *Bacillus subtilis* PR13 and *B. subtilis* PR9, were observed at the addition of 1 and 10 $\mu\text{g/ml}$ of biotin and nicotinic acid respectively. There was a negative correlation between biotin and lysine production by *B. subtilis* PR13 ($r= -0.99$), while there was a positive correlation between nicotinic acid and lysine production by *B. subtilis* PR9 ($r= 0.38$).

The results of the effect of antibiotics on lysine production by *B. subtilis* PR13 and PR 9 are presented in Figures 5-6. The results showed that maximum lysine yields of 2.38 and 1.64 mg/ml by *Bacillus subtilis* PR13 and *B. subtilis* PR9, were observed at the addition of 0.04 units/ml of penicillin. There was a negative correlation between penicillin and lysine production by the *Bacillus subtilis* PR 13 and 9 ($r= -0.90$, and -0.90) respectively.



UNDER PEER



UNDER PEE






Figure 3: Effect of Antibiotics on Lysine Production by *Bacillus subtilis* PR13: A, Vancomycin; B, Erythromycin; C, Penicillin (units/ml); D, Lincomycin; E, Tetracycline; F, Chloramphenicol; G, Control(without antibiotics)

UNDER PLE




UNDER PEER REVIEW



Comment [D17]: It would be better if color is added to enhance the effect of figure and also for comparison.

UNDER PEER REVIEW



Comment [D18]: It would be better if color is added to enhance the effect of figure and also for comparison.

Discussion

Biotin stimulated lysine yield and biotin was found as a requirement for growth and lysine production. Shah *et al.* [23] studied the effects of biotin on lysine production by *Cornebacterium glutamicum* and observed that maximum lysine was produced at 5µg per 100ml of biotin. A gradual increase in the level of biotin in the medium stimulated growth and methionine production. Tosaka *et al.* [24] investigated the role of biotin levels on L-lysine formation in *B.lactofermentum*. They reported that accumulation of L-lysine was stimulated considerably by increasing the biotin level. Zaki *et al.* [25] worked on the effect of non-ionic detergents and vitamin on the production of amino acids by *Bacillus ammoniagenes*. They found that the presence of 20µg/l biotin induced the production of about 166mg% L-lysine. Also Banik and Majumdar [26] reported a maximum methionine production by an auxotrophic mutant of

Comment [D19]: Citation number should be at the end of the sentence. Please check and correct other's too.

Micrococcus glutamicus at a biotin level suboptimal for growth. A further increase in the biotin level stimulated growth, but not methionine production. The exact role of biotin in the amino acid production by microorganisms is not clear. While Tanaka *et al.* [27] believe that biotin functions by limiting growth and allowing the carbon and nitrogen sources to the formation of amino acids rather than to the synthesis of cell matter, Shiio *et al.*, [28], Veldkamp *et al.*, [29] and Otsuka *et al.*, [30] believed that a low biotin concentration makes the bacterial cells more permeable resulting in a higher leaching out of amino acids into the surrounding medium. Beker [31] developed threonine- methionine double auxotrophic mutant of *Brevibacterium flavum* which required biotin. He found that at low concentration of biotin, biosynthesis of glutamic acid takes place and intensive synthesis of L-lysine can be observed at the beginning of the stationary phase of growth. Young and Chipley [32] investigated the role of biotin in L-lysine production in *Brevibacterium lactofermentum*. They observed that biotin treated cell took up more glucose, than did the control one. Biotin apparently, caused some compositional changes in the cell wall membrane complex, allowing an increase in uptake of glucose. The result of uptake studies and fatty acid analysis suggested that biotin affected the cell surface, probably the bacterial membrane. It is well known that bacterial membrane plays an important role as a charged barrier. This mechanism might also regulate the amount of L-lysine released by the cells.

The antibiotics stimulated different degree of lysine production. Penicillin encouraged lysine production in all the strain. Sen and Chatterjee [33] tested the effects of different antibiotics on lysine production by *Arthrobacter globiformis* and observed that they stimulated growth and lysine production. Vancomycin stimulated lysine production in *Bacillus subtilis* PR9 and *Bacillus pumilus* SS16. This is in agreement with the findings of Ekwealor and Obeta [34] who

reported increase in lysine production in *Bacillus megaterium* when vancomycin was used.

Tetracycline stimulated lysine production in *Bacillus subtilis* PR13. Zaki *et al.*[35] reported that 22-24g/l lysine was produced by *Micrococcus glutamicum* when tetracycline was added to the fermentation medium. Demian and Brinbum [36] suggested a change in permeability of cell wall caused by antibiotics which may be responsible for improved amino acid yield. This change in permeability affects the intercellular accumulation of amino acids, with the result that the amino acid can no longer regulate its own synthesis by feed back control, thereby releasing high levels of amino acid into the medium. Chloramphenicol did not stimulate lysine production in *Bacillus subtilis* PR9. Israelides *et al.* [37] reported that In immobilized cell preparations, growth of cells outside the immobilization matrix, as free cells, is normally undesirable due to the appearance of cells in the product stream and clogging of such systems. Antibiotics could be used to arrest such free cell growth, while allowing the synthesis and excretion of the product into the medium. Chloramphenicol at 200 µg/ml effectively arrests free cell growth and hence the L-lysine being produced can be entirely attributed to the immobilized cells. Novobiocine on the hand at concentration of 100µg/ml, stopped free cell growth, and also prevented the production of L-lysine. Productivity and yields of L-lysine were adversely affected by chloramphenicol and novobiocin probably due to a great decrease in cell viability.

It was observed from the study that yeast extract stimulated the highest lysine yield in *Bacillus subtilis* PR9. This is in line with the report of Morinaga *et al.*[38], who observed an increase in lysine production by *Pseudomonas* species 518 when 7.5mg/ml of yeast extract was added. Ekwealor and Obeta [39]

reported that the growth promoting substances used retarded growth but stimulated lysine accumulation. [40] who studied the effect of selected nutrients on lysine production from whey permeate by *Brevibacterium lactofermentum* ATCC 210 reported a lysine production of 3.3g/l when 0.2% yeast extract was added. They believed that yeast extract contain certain components that stimulate lysine production. Also in a study done by Yamada *et al.*[41] to study the effects of organic nitrogen sources on L-methionine production by *Methylotroph* strain OM 33, reported that 5mg/ml yeast extract stimulated the optimum yield of L-methionine. Chan and Foster [42] and Tauro *et al.*[43] , reported a retardation in glutamic acid production by *Bacillus subtilis* and lysine accumulation in *Ustilago maydis*, respectively when growth promoters were used.

5. Conclusion

During the optimization study, which included influence of addition of vitamins, antibiotics, growth stimulators, it was observed that there was improved L-lysine production by *B. subtilis* PR13 and *B. subtilis* PR9. However, *B. subtilis* PR13 produced the maximum yield of L-lysine. The supplementation of 0.1% v/v of peptone, 1 µg/ml of biotin and 0.04 units/ml of penicillin were optimal for L-lysine production by *B. subtilis* PR13. The *Bacillus* species have shown potential for lysine production using readily available agricultural products. These products are good sources of carbon and nitrogen and are rich in fermentable substrates. This development indicates that large scale L-lysine production is feasible in Nigeria, this will help to meet present-day needs in its industrial sector.

Comment [D20]: Please rewrite conclusion part with appropriate concluding lines.

References

- [1] Eggeling L and Bott M. (2015). A giant market and a powerful metabolism: L-lysine provided by *Corynebacterium glutamicum*. *Appl Microbiol Biotechnol* 99(8):3387–3394.
- [2] Malothu R, Dowlathabad MR and Balanarasaiah T. (2012). Bioprocess Technology Development and Convalescing Production of L-Lysine in *Corynebacterium glutamicum*. International Conference on Chemical Engineering and its Applications , Bangkok (Thailand).
- [3] Kobayashi K. and Nishikawa M.(2006). Promotion of E-Poly-L-Lysine Production by Iron in *Kitasatospora kifune*. *World Journal of Microbiology and Biotechnology*, 10, 1033-1036.
- [4] Shah AH and Hameed A (2004). Nutritional and mutational aspects of indigenous lysine production by *Corynebacterium glutamicum*: comprehensive studies in shake flask fermentation-1. *J Chem Soci Pak*, 26: 416-428.
- [5] Hermann T. (2003). Industrial production of amino acids by Coryneform bacteria. *Journal of Biotechnology*, 104: 155-72.
- [6] Tosaka O. (1983). The Production of L-Lysine by Fermentation. *Trends in Biotechnol.* 1: 70-74.
- [7] Javaid M M, Haq IU, Sohail MI and Bokhari SAI (2012). Mutagenesis induced hyper- production of L- Lysine in shake flask and fermentor by *Brevibacterium flavum* IIBUV2. *Pakistan J. Bot.* 44: 347- 353
- [8] Haas EM (2006). *Staying healthy with nutrition*. 21 Century Edition, ISBN-13: 978- 980.
- [9] Shah, A. H. 1998. *Studies on the Microbial Production of Lysine*. Ph.D thesis. Quaid-i-Azam University, Islamabad.

- [10] Sen SK and Chatterjee SP. (1989). Influence of B-vitamins and trace elements on lysine production by *Micrococcus varians* 2fa, *Acta Biotechnol.* **9**: 63-67.
- [11] Leuchtenberger W. (1996). Amino acids – technical production and uses. In Rehm H.J., Reed, G., Phuler, A., Stadler, P., editors. Products of primary metabolism. *Biotechnology* vol. 6. Wiley- VCH, Weinheim, German. p 492.
- [12] Kircher M, and Pfeerle W. (2001).The fermentative production of L-lysine as an animal feed additive. *Chemosphere.* 43: 27-31.
- [13] Ekwealor, I.A. and Obeta, J.A.N.(2008). Antibiotics and surfactants effects of lysine accumulation by *Bacillus megaterium*. *African Journal of Biotechnology*, 7(10):1550-1553.
- [14] Nielsen J. (2001). Metabolic engineering. *Appl Microbiol Biotechnol.* 2001; **55**: 263-283.
- [15] Hermann T.(2003). Industrial production of amino acids by Coryneform bacteria. *J Biotechnol.* 104: 155-72.
- [16] Ikeda M. (2003). Amino acid production process. *Adv. Biochem. Eng. Biotechnol.* 79: 1–35.
- [17] Moosavi-Nasabi M, Ansari S. and Montazeri Z. (2007). Fermentative Production of Lysine by *Corynebacterium glutamicum* from Different Carbon Source. *Iran Agricultural Research*, Vol. 25, No. 2 99-106
- [18] Okpalla J. and Ekwealor IA. (2019a). Screening for lysine production by bacteria isolated from Nigerian soil. *World wide J Multidiscip Res Develop.* 5(4): 10-18.
- [19] Okpalla J and Ekwealor IA (2019b). Studies on lysine accumulation in broth culture of *Bacillus* species using carbohydrates as carbon sources

and seed meals as nitrogen sources. *Inter J Trend in Scientific Res Development*. 3(2): 760- 772.

- [20] Umerie SC, Ekwealor IA and Nwagbo IO (2000). Lysine production of *Bacillus laterosporus* from various carbohydrates and seed meals. *Bioresource Technology* **75**: 249-252.
- [21] Chinard FP. (1952). Photometric estimation of proline and ornithine. *J Biol Chem*, **199**: 91-95.
- [22] Miller GL. (1959). Use of dinitrosalicylic acid reagent for detection of reducing sugar. *Annals of Chem*. **31**: 427-431.
- [23] Shah AH, Hameed A. and Guj Madid K. (2002). Improved microbial production of lysine by developing a new Auxotrophic mutant of *Corneybacterium glutamicum*. *Pakistan Journal of Biological Sciences*, **5**(1): 80-83.
- [24] Tosaka, O., Hirakawo, H. and Takinami, K. (1979a). Effect of biotin level on lysine formation in *Brevibacterium lactofermentum*. *Agricultural and Biological Chemistry*, **43**: 491-5.
- [25] Zaki D, Aziz MA, Naguib M. and Shalbi A. (1987). Effect of non-ionic detergents and vitamins on the amino acid synthesis by *Brevibacterium ammoniagenes*. *Zentralel Mikrobiology*, **142**:333–339.
- [26] Banik AK. and Majundar SK. (1975). Effect of minerals on production of methionine by *Micrococcus glutamicum*. *Indian Journal of Experimental Biology* **13**: 510-2.
- [27] Tanaka K, Iwasaki T. and Kinoshita S. (1966). L-glutamic acid fermentation v. Biotin and L-glutamic accumulation by bacteria. *Nippon Noegi Kagaku Kaishi*, **34**: 593.
- [28] Shiio I, Otsuka S. and Takahashi M. (1962). Effect of biotin on the bacterial formation of glutamic acid I. Glutamate froamtion and cellular permeability of amino acids. *Journal of Biochemistry*, **51**: 56-62.

- [29] Veldkamp H, Vandenberg G and Zevenhuizen LPTM. (1963). Glutamic acid production by *Arthrobacter globiformis*. *Antonie Van Leeuwenhoek Journal of Microbiology and Serology*, **29**: 35-51.
- [30] Otsuka S, Miyajima R and Shiio I. (1965). Comparative studies on the mechanism of microbial glutamate formation II. Effect of biotin. *Journal of General and Applied Microbiology*, **11**: 295.
- [31] Beker MV, (1982). Product biosynthesis in continuous fermentation. *Folia Microbiology*, **27**:315–318.
- [32] Young TK. and Chipley JR (1984). Role of biotin in the production of lysine by *Brevibacterium lactofermentum*. *Microbiology*, **40**: 161-171.
- [33] Sen K. and Chatterjee SP (1983). Extracellular lysine production from hydrocarbon by *Arthrobacter globiformis*. *Folia microbiology*, **28**: 292–300.
- [34] Ekwealor IA. and Obeta JAN (2007). Effect of vitamins and bivalent metals on lysine yield in *Bacillus megaterium*. *African Journal Biotechnology*, **6**(11): 1348-1351.
- [35] Zaki D, Galal O, Hazino-Wahba SA, Morsi KI and Wakell EA.(1982). Microbiological production of lysine. *Nutrition Report International*, **26**: 537-546.
- [36] Demain A L and Brinbum J. (1965). Alteration of permeability for the release of metabolites from the microbial cell. *Current Topics in Microbiology*, **46**: 1-28.
- [37] Isrealides CJ, Weir A and Bull AT.(1989). Effect of antibiotic on lysine production in free and immobilized cells of *Bacillus subtilis*. *Applied Microbiology and Biotechnology*, **32**:134–136.
- [38] Morinaga Y, Tani Y and Yamada H. (1982a). L-methionine production by ethionine resistance mutant of facultative methylotroph, *Pseudomonas* FM 518. *Agricultural and Biological Chemistry* **46**: 473-80.
- [39] Ekwealor A and Obeta AN.(2005). Studies on lysine production by *Bacillus megaterium*. *African J. Biotechnol.* **4**: 633-638.

- [40] Young TK and Chipley JR. (1983). Microbial production of lysine and threonine from whey permeate. *Applied Environmental Microbiology*, **45**:610–615.
- [41] Yamada H, Morinaga Y and Yoshiki T. (1982). L-methionine over production by Ethionine-resistant mutants of obligate Methyloph strain OM 33. *Agriculture and Biological Chemistry*, **46**(1): 47-55.
- [42] Chao KC and Foster JW.(1959). A glutamic acid producing *Bacillus*. *Journal of Bacteriology*, **77**:715-725.
- [43] Tauro P, Ramachandra-Rao TN, Johar DS, Screenivasan A and Subrahmanyam V. (1963). L-lysine production by Ustilaginales fungi. *Agricultural and Biological Chemistry*, **27**: 227-235.

UNDER PEER REVIEW