

Original Research Article

Effect of different types of lights on growth, yield and quality of Kale” (*Brassica oleracea var. Acephala*)

ABSTRACT

A research project was conducted to investigate the impact of different types of lighting on the growth, yield, and quality of Kale (*Brassica oleracea var Acephala*), under the Prayagraj Agro Climatic conditions. The study was conducted at the Vegetable Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj, Uttar Pradesh, during the academic session of 2020-2022. The experiment utilized a statistical design of Randomized Block Design with two replications. According to the study's findings, among the various treatment levels, treatment T₂ (LED tube white) showed the highest values in growth parameters such as plant height (30.01 cm), petiole length (15.07 cm), and number of leaves per plant (14.33 leaves). It also demonstrated the highest yield parameters, fresh weight of leaves (24.40 gms), leaves yield per bag (50.54 g). Additionally, T₂ exhibited better quality parameters such as total soluble solids (3.23 °Brix), shelf-life (4.67 days), and chlorophyll content (56.40 nmol cm²), while the lowest values were observed in T₄ (LED FULL SPECTRUM).

Keywords: *Kale plant growth, yield and quality.*

INTRODUCTION

Kale, scientifically known as *Brassica oleracea*, has a long history of cultivation that spans over two millennia. Its resistance to frost has made it especially important in colder regions throughout history. As a member of the Brassicaceae family, kale is considered one of the most important leafy green vegetables. The word "kale" is derived from the Scottish term "coles" or "caulis," which the Greeks and Romans used to refer to cabbage-like plants. The edible portions of kale are highly curled, bluish-green leaves, and the plant does not form a solid head. Kale is one of the oldest forms of cabbage and originally comes from the eastern Mediterranean region of Europe. It has a similar appearance to the leafy canola plant. In the 19th century, kale was introduced as a minor temperate vegetable in India. Although it is

Comment [H1]: False information by authors

Comment [H2]: Mismatch with material methods means conducted without advise of statistician

Comment [H3]: var *Acephala*

Comment [H4]: , and never together

rarely found in India, it is commercially grown on a large scale in Kashmir and to a limited extent in Jammu, Assam, and Himachal Pradesh. Kale is a food that is low in calories and has a significant number of vitamins, particularly vitamin C, E, and K, as well as micronutrients such as iron, zinc, and manganese, and macronutrients like calcium and magnesium. It is also a source of dietary fibre, glutamine (an amino acid with anti-inflammatory properties), and plant phytochemicals including polyphenols, flavonoids, and carotenes. Consuming leafy vegetables on a daily basis can reduce the risk of cancer and heart disease, prevent fatigue, promote overall well-being, and delay the effects of aging. The use of various colours of light to demonstrate the relationship between light and plant growth is limited to three distinct colours: red, blue, and yellow. Photosynthesis is the food-making process in leaves that is powered by light. Green is the colour most leaves reflect rather than absorbing, which is why leaves appear green. A promising technological advancement to obtain high density growth and manipulate morphological traits and phytochemical composition of crops is the use of multilayer production under sole-source (SS) light-emitting diodes (LEDs) as an artificial lighting system in vertical farming operations. Light plays a crucial role in plant growth and morphology, and the recent LED technology has several advantages such as energy efficiency, low maintenance cost, longevity, and the possibility to control spectral composition and select high light intensity while maintaining low heat emission. Optimal management of light intensity and spectral composition is required to achieve optimal yield, appearance, and nutritional quality of plants. Green light can deeply penetrate plant canopies and affect plant growth and the synthesis of bioactive compounds. The effects of green light on plant physiology depend on its proportion in combination with red and blue LED lighting in controlled environmental agriculture. High proportions of green light (25-44%) can reverse the effects of blue or red light, leading to stem growth rate inhibition, reduced chloroplast gene expression, reduced stomatal opening, and decreased phytochemical accumulation. This can negatively affect the quality of green leafy vegetables.

Comment [H5]: low calorie food

Comment [H6]: delet

Comment [H7]: delet

Comment [H8]: delet

Comment [H9]: delete

Comment [H10]: delet

Comment [H11]: deelopment phase

Comment [H12]: delet

Comment [H13]: delet

Comment [H14]: delet

MATERIALAND METHODS

The present investigation entitled was done to understand the plant growth, fruit yield and quality of Kale crop variety curly leaf under influence of different lights. The details of the materials used, and the methods adopted in the investigation, which was carried out at Horticultural Research Farm (HRF), Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj during the *winter* season of 2022 are described under the following heads. The

Comment [H15]: fruit yield happens ?

Comment [H16]: Mislead as par abstract submit 2020-22 table , Without SOP of Kale

details of 8 different light used are mentioned in table 1 and replicated thrice. Observations were recorded at different stages of growth periods. The data were statistically analysed by the method suggested by Fisher and Yates, 1936.

Comment [H17]: Mismatch with Abstract confirm ANOVA

Comment [H18]: Not exist

Table 1 Treatment Details

Treatment Notation	Photosynthetically Active Radiation (PAR)
T ₀	Outdoor under sunlight
T ₁	LED bulb white
T ₂	LED tube white
T ₃	General bulb
T ₄	LED bulb full spectrum
T ₅	LED white bulb + LED bulb full spectrum
T ₆	LED tube white + LED bulb full spectrum
T ₇	General bulb + LED bulb full spectrum
T ₈	LED bulb full spectrum

RESULTS AND DISCUSSION

Plant height

The data pertaining to effect of application on height of plant is presented in Table.2. At 15 DAS treatment T₂ produced tallest plant of height (7.87) cm followed by treatment T₀ (7.43), treatment T₁ (7.33), treatment T₆ (7.07), and treatment T₄ significantly recorded the lowest height of plant (6.01). At 30 DAS treatment T₂ produced tallest plant of height (18.03 cm) followed by treatment T₀ (18.03), treatment T₁(17.29) treatment T₆ (17.17), treatment T₅ (16.93), T₄ significantly recorded the lowest height of plant (15.67 cm). At 60 DAS treatment T₂ produced tallest plant height (30.10cm) followed by treatment T₀ (27.78cm), treatment T₁ (27.00cm), treatment T₆ (26.76cm), treatment T₈ (25.03cm), and treatment T₄ significantly recorded the lowest height of plant (22.75cm). Similar findings were reported by Gerovac *et al*, (2016) and Ying *et al.*, (2020) in Brassicas; Yao *et al.*, (2017) and Yao *et al.*, (2020) in rapeseeds; Zhang *et al.*, (2020) in Kale.

Comment [H19]: Unit

Comment [H20]: T2 must be disclosed

Comment [H21]: unit always with data

Comment [H22]: treatment must be disclose uniformly

Comment [H23]: Not exist in References

Length of petiole (cm)

The data pertaining to effect of application on Length of petiole is presented in Table. 2. Treatment T₂ produced largest petiole (15.07 cm) followed by treatment T₀ (13.92cm), treatment T₃ (13.24), treatment T₆ (12.60cm), and treatment T₄ recorded significantly the lowest main branch (9.45cm). Similar findings were reported by Gerovac *et al*, (2016) and Ying *et al.*, (2020) in Brassicas; Yao *et al.*, (2017) and Yao *et al.*, (2020) in rapeseeds; Zhang *et al.*, (2020) in Kale.

Comment [H24]: lower case uniformly

Comment [H25]: Check T1 data also

Comment [H26]: Not exist in reference

Number of leaves/plants

The data pertaining to effect of application on Number of leaves/plants is presented in Table.2. Treatment T₂ produced highest Number of leaves/plant (14.33 leaves) followed by treatment T₀ (13.67 leaves), T₁ (13.00 leaves), treatment T₃ (12.67 leaves), treatment T₆ (12.33 leaves), and treatment T₄ significantly recorded the lowest Number of leaves/plant (11.00 leaves). Similar findings were reported by Gerovac *et al*, (2016) and Ying *et al.*, (2020) in Brassicas; Yao *et al.*, (2017) and Yao *et al.*, (2020) in rapeseeds; Zhang *et al.*, (2020) in Kale.

Comment [H27]: lower case

Comment [H28]: lower case

Comment [H29]: lower case in running

Days to Leaves picking

Treatment T₂ produced higher Days of leaves picking 62.67 followed by treatment T₀ (66.67), treatment T₁ (68.67) treatment T₃ (69.00), treatment T₆ (69.33), T₄ significantly recorded the lower Days of leaves picking 75.00. Similar findings were reported by Gerovac *et al.*, (2016) and Ying *et al.*, (2020) in Brassicas; Yao *et al.*, (2017) and Yao *et al.*, (2020) in rapeseeds; Zhang *et al.*, (2020) in Kale.

Comment [H30]: lower case

Comment [H31]: Missing

Fresh weight of leaves

Treatment T₂ produced highest Fresh weight of leaves 24.40 g followed by treatment T₀ (22.29 g), treatment T₁ (21.48 g), treatment T₃ (20.83 g), treatment T₆ (20.51 g), T₄ recorded significantly the lowest Fresh weight of leaves 14.48 g. Similar findings were reported by Gerovac *et al.*, (2016) and Ying *et al.*, (2020) in Brassicas; Yao *et al.*, (2017) and Yao *et al.*, (2020) in rapeseeds; Zhang *et al.*, (2020) in Kale.

Comment [H32]: lower case uniformly

Comment [H33]: lower case uniformly

Comment [H34]: Missing

Leaf yield/bag(g)

Treatment T₂ produced highest Leaf yield/bag 50.54g followed by treatment T₀ (45.57g), treatment T₁ (43.92g), treatment T₃ (42.96g), treatment T₆ (42.67g), T₄ significantly recorded the lowest Leaf yield/bag 23.22g. Similar findings were reported by Gerovac *et al.*, (2016) and Ying *et al.*, (2020) in Brassicas; Yao *et al.*, (2017) and Yao *et al.*, (2020) in rapeseeds; Zhang *et al.*, (2020) in Kale.

Comment [H35]: uniformly in running leaf yield

Leaves yield/hectare (Kg/ha)

Estimated leaves yield per hectare was observed in Treatment T₂ produced High Leaves yield/hectare 14.93 followed by treatment T₀ (14.91), treatment T₁ (14.38), treatment T₃ (14.30), treatment T₆ (13.80), T₄ significantly recorded the lowest Leaves yield/hectare 10.49. Similar findings were reported by Gerovac *et al.*, (2016) and Ying *et al.*, (2020) in Brassicas; Yao *et al.*, (2017) and Yao *et al.*, (2020) in rapeseeds; Zhang *et al.*, (2020) in Kale.

Comment [H36]: high leaves

Comment [H37]: Unit must be mentioned in kg ha⁻¹

Comment [H38]: lower case

Comment [H39]: Missing in Ref

TSS

Treatment T₂ produced high TSS 3.23 °Brix followed by treatment T₀ (2.19 °Brix), treatment T₁ (1.72 °Brix), treatment T₇ (1.58 °Brix), treatment T₆ (1.53 °Brix), T₄ recorded significantly the lowest TSS 0.51 °Brix. Similar findings were reported by Gerovac *et al.*,

Comment [H40]: Not match with table data

(2016) and Ying *et al.*, (2020) in Brassicas; Yao *et al.*, (2017) and Yao *et al.*, (2020) in rapeseeds; Zhang *et al.*, (2020) in Kale.

Shelf-life

Treatment T₂ produced high Shelf life 4.67 days followed by treatment T₀ (4.00 days), treatment T₁ (3.67 days), treatment T₇ (3.33 days), treatment T₆ (3.33 days), T₄ significantly recorded the lowest Shelf life 1.67 days.

Comment [H41]: Treatment must be disclose uniformly

Comment [H42]: lower case

Chlorophyll Content

Treatment T₂ produced high Chlorophyll content 56.40 nmol cm² followed by treatment T₀ (52.54 nmol cm²), treatment T₁ (50.17 nmol cm²), treatment T₇ (49.62 nmol cm²), treatment T₆ (48.23 nmol cm²), T₈ significantly recorded the lowest Chlorophyll content 40.39 nmol cm².

Comment [H43]: lower case

Comment [H44]: lower case

Conclusion

Based on the findings of present investigation, it is concluded that treatment T₂ (LED Tube White) performed best in respect to all other treatments for growth yield and quality of Kale under influence of light

Comment [H45]: Re-write

References

- Alrifai, O.; Hao, X.; Marcone, M.F.; Tsao, R. (2019). Current review of the modulatory effects of LED lights on photosynthesis of secondary metabolites and future perspectives of microgreen vegetables. *J. Agric. Food Chem.* 67, 6075–6090.
- Devlin, P.F.; Christie, J.M.; Terry, M.J. (2007). Many hands make light work. *J. Exp. Bot.* 58, 3071–3077.
- Di Gioia, F.; Santamaria, P. (2015). Microgreens, Agrobiodiversity and Food Security. In *Microgreens*; Di Gioia, F., Santamaria, P., Eds.; Ecologica: Bari, Italy, pp. 7–23.
- Dou, H.; Niu, G.; Gu, M. (2019). Photosynthesis, morphology, yield, and phytochemical accumulation in basil plants influenced by substituting green light for partial red and/or blue light. *HortScience.* 54, 1769–1776.

Comment [H46]: All References are not in formate of journal

- Dou, H.; Niu, G.; Gu, M.; Masabni, J. (2020).** Morphological and Physiological Responses in Basil and Brassica Species to Different Proportions of Red, Blue, and Green Wavelengths in Indoor Vertical Farming. *J. Am. Soc. Hortic. Sci.* 145, 267–278.
- Gerovac, J.R.; Craver, J.K.; Boldt, J.K.; Lopez, R.G. (2016).** Light intensity and quality from sole- source light-emitting diodes impact growth, morphology, and nutrient content of Brassica microgreens. *Hort Science.* 51,497–503
- Goto, E. (2012).** Plant production in a closed plant factory with artificial lighting. *ActaHortic.* 956,37–49.
- Hernández, R.; Kubota, C. (2016).** Physiological responses of cucumber seedlings under different blue and red photon flux ratios using LEDs *Environ. Exp. Bot.* 121, 66– 74.
- Jeong-Wook .H., Jae-Su. L ., Gong-In. L., Hyun-Hwan. K. (2017).** Growth of Kale seedlings affected by the control of light quality and intensity under smart greenhouse conditions with artificial lights. *Korean Journal of Environmental Agriculture*, Vol.36. I(3) : 193-200.
- Kyriacou, M.C.; Roupael, Y.; Di Gioia, F.; Kyratzisz, A.; Serio, F.; Renna, M.; De Pascale, S.; Santamaria, P. (2016).** Micro-scale vegetable production and the rise of microgreens. *Trends Food Sci. Technol.* 57, 103–115.
- Landi, M.; Zivcak, M.; Sytar, O.; Brestic, M.; Allakhverdiev, S.I. (2020).** Plasticity of photosynthetic processes and the accumulation of secondary metabolites in plants in response to monochromatic light environments: A review. *Biochim. Biophys. Acta Bioenerg.* 1861,148131.
- Olle Margit and Alsin Iva (2019),** Influence of wavelength of light on growth, yield and nutritional quality of greenhouse vegetable, *Proceedings of the Latvian academy of sciences*, Vol.73, No.1(718): 1-9.
- Paradiso, R.; Proietti, S. (2021).** Light-Quality Manipulation to Control Plant Growth and Photo morphogenesis in Greenhouse Horticulture: The State of the Art and the Opportunities of Modern Led Systems. *J. Plant Growth Regul.* 41, 742–780.

- Romeo, D.; Vea, E.B.; Thomsen, M. (2018).** Environmental impacts of urban hydroponics in Europe: A case study in Lyon. *Procedia CIRP* (2018), 69, 540–545.
- Samuolienė, G.; Brazaitytė, A. (2013); Jankauskienė, J.; Viršilė, A.; Sirtautas, R.; Novičkovas, A.; Sakalauskienė, S.; Sakalauskaitė, J.; Duchovskis, P.** LED irradiance level affects growth and nutritional quality of Brassica microgreens. *Cent. Eur. J. Biol.* 8, 1241–1249. *Horticulturae* (2022), 8, 217–18 of 20.
- Yao Xu-yang., Liu Xiao-ying., Xu Zhi-gang., Jiao Xue-lei. (2017),** Effects of light intensity on leaf micro structure and growth of rape seedlings cultivated under a combination of red and blue LEDs, *Journal of Integrative Agriculture*, Vol.16, I(1): 97-105.
- Ying, Q.; Jones-Baumgardt, C.; Zheng, Y.; Bozzo, G. (2021).** The Proportion of blue light from light-emitting diodes (RB light environment) alters microgreen phytochemical profiles in a species-specific manner. *HortScience* 56,13–20.
- Ying, Q.; Kong, Y.; Jones-baumgardt, C.; Zheng, Y. (2020).** Responses of yield and appearance quality of four Brassicaceae microgreens to varied blue light proportion in red and blue light-emitting diodes lighting. *Sci. Hortic.* 259,108857.
- Zhang Yiting, Ji Jazeng, Song Shiwei, Su Wei and Liu Houcheng (2020),** Growth, nutritional quality and health promoting compounds in Chinese kale grown under different ratios of red: blue LED lights, *Agronomy*, Vol.10, 1248.

Table 2 Performance of Kale on different light treatments for different parameters.

Treatment Notation	Photosynthetically Active Radiation (PAR)	Plant height (cm) 15DAS	Plant height (cm) 30 DAS	Plant height (cm) 60 DAS	Length of petiole (cm)	Number of leaves/plants	Days to leaves picking	Fresh weight of leaves (g)	Leaves yield/bag (g)	Yield per hectare (Kg/ha)	TSS [°Brix]	Shelf-life (days)	Chlorophyll content (nmol cm ²)
T ₀	Outdoor under sunlight	7.43	18.13	27.78	13.92	12.00	66.67	22.29	45.57	14.91	2.91	4	52.54
T ₁	LED bulb white	7.33	17.29	27	13.29	11.33	68.67	21.48	43.92	14.38	1.72	3.67	50.17
T ₂	LED tube white	7.87	18.03	30.01	15.07	14.33	62.67	24.4	50.54	14.93	3.23	4.67	56.4
T ₃	General bulb	6.48	16.01	23.08	13.24	12.33	69.00	20.83	42.96	14.30	1.37	2.67	44.39
T ₄	LED bulb full spectrum	6.01	15.67	22.75	9.45	12.33	75.00	15.78	23.22	10.49	0.51	1.67	40.41
T ₅	LED white bulb + LED bulb full spectrum	6.87	16.93	23.91	11.47	11.00	71.00	16.15	33.89	11.13	1.22	2.67	42.39
T ₆	LED tube white + LED bulb full spectrum	7.07	17.18	26.76	12.6	13.67	69.33	20.51	42.67	13.80	1.53	3.33	48.23
T ₇	General bulb + LED bulb full spectrum	6.81	17.78	23.13	10.92	12.67	72.67	14.48	33.63	11.45	1.58	3.33	49.62
T ₈	LED bulb full spectrum	6.67	16.43	25.03	12.35	13.00	69.67	17.98	39.45	12.52	1.39	2.67	45.74
'F' Test		S	S	S	S	S	S	S	S	S	S	S	S
S.E. (m) (±)		0.697	0.079	0.103	0.213	1.150	1.455	0.023	0.083	1.000	0.261	0.803	0.075
C.D. (5%)		0.144	0.168	0.219	0.455	1.310	3.111	0.049	0.178	2.138	0.557	1.126	0.035
C.V.		12.293	0.573	0.493	0.491	11.250	2.568	0.146	0.0258	9.346	19.511	13.212	0.123

Comment [H47]: Keep two digit after decimal