

**Effect of Spacing and Nitrogen Management on Yield and Economics of
Summer Sesame (*Sesamum indicum* L.)**

ABSTRACT

A field experiment was carried out in *Zaid 2022* at the Department of Agronomy's Agricultural Research Farm in Prayagraj (Uttar Pradesh). There are three planting geometries used in the experiment 30 cm x 10 cm, 40 cm x 10 cm, and 50 cm x 10 cm. There are three nitrogen management treatments used in the experiment 50% of the Recommended Dose of Nitrogen (RDN) through inorganic fertilizer, 50% of the RDN through inorganic fertilizer in combination with 50% of Nitrogen provided through Farm yard manure with *Azotobacter* seed inoculation, and 50% of the RDN through inorganic fertilizer. Ten treatments were duplicated three times in the randomized block design of the experiment. Findings showed that with 30x10cm spacing ,50% RDN + 50% N through Farm yard manure + biofertilizer (*Azotobacter*) highest grain yield (552.38 kg/ha), net return (30221.00 INR/ha), gross return (71809.00 INR/ha), and benefit: cost ratio (1.73).

Key words:

Spacing, Farm Yard Manure (FYM), Recommended Dose of Nitrogen (RDN), *Azotobacter*, and *Zaid*.

INTRODUCTION

Sesame (*Sesamum indicum* L.) is being cultivated from 3000 years, this is the oldest oilseed crop with rich nutty flavor. In India we can see sesame seed in many colors like black, red, brown, white. Oil content in sesame varies from 40 to 50%. Sesame is also known as Queen of oilseeds. These seeds are used in candy making, baking and other religious rituals. The sesame oil is used in making perfumes, soaps, paints etc. In India major sesame producing states are Gujarat, West Bengal, Tamil Nadu, Andhra Pradesh, Madhya Pradesh and Maharashtra.

Method of sowing determines plant population in the field. The most common method used for sowing summer sesame is broadcasting which results in uneven spacing in the field. Uneven plant spacing is responsible for low yield. Caliskan et al. (2004) reported positive effects of row planting.

Providing optimum nitrogen fertilizers to sesame crop also effect the intake of other nutrients, importantly P and K and some other micronutrients. Gebregergis Z. (2019) states that N P K fertilizers have been extensively researched and proven to significantly increase sesame yield. Particularly, growth and yield of sesame are greatly influenced by the application of N fertilizer.

Spacing and nitrogen are important factors in growth and yield of sesame. Therefore this study "Effect of spacing and nitrogen management on yield and economics was carried out in Department of Agronomy, Agriculture Research Farm in Prayagraj(Uttar Pradesh).

MATERIALS AND METHODS

The experiment was conducted in the *Zaid* of 2022 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agricultural, Technology and Sciences (SHUATS), Prayagraj (U.P.) This place is situated at 25° 57'12" N latitude and 87° 50'12" E longitude, 98 meters above mean sea level (MSL). This region is situated on the *Yamuna River's* right bank along the Prayagraj Rewa road, some 12 kilometers from the city. Ten treatments, each replicated three times and categorized as having three levels, were used in an experiment to see how spacing and nitrogen management affected the development and output of summer sesame (*Sesamum indicum* L.). The farm is situated at 25 degrees, 98 meters above mean sea level. 87° 50'12" East, 25° 57'12" North latitude; sea level (MSL). This region is situated on the *Yamuna River's* right bank along the Prayagraj, Rewa path, about 12 kilometres from the city.

Experimental Design:

The experimental design was set up using a randomized block design with 10 treatments that are duplicated three times to examine the effects of spacing and nitrogen management on summer sesame (*Sesamum indicum* L.) growth and yield.

When employed in combinations the treatments was separated into 3 levels of plant geometry and 3 for nitrogen as follows : T₁: Spacing 30cm × 10cm + 50% RDN + 50% N through FYM, T₂: Spacing 30cm × 10cm + 50% RDN + Biofertilizer (*Azotobacter*) , T₃: Spacing 30cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer(*Azotobacter*), T₄: Spacing 40cm × 10cm + 50% RDN + 50% N through FYM, T₅: Spacing 40cm × 10cm + 50% RDN + Biofertilizer (*Azotobacter*), T₆: Spacing 40cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (*Azotobacter*), T₇: Spacing 50cm × 10cm + 50% RDN + 50% N through FYM, T₈: Spacing 50cm × 10cm + 50% RDN + Biofertilizer (*Azotobacter*), T₉: Spacing 50cm × 10cm + 50% RDN 50% N through FYM + Biofertilizer (*Azotobacter*), T₁₀: 10. Spacing 30cm × 10 cm + 50 : 40 : 30 NPK Kg/ha (Control)

When the oil harvest was ready, it was carefully handled. After harvesting, seeds were extracted from each net plot and dried for three days under the sun. Five randomly selected sample plants from each plot of each replication were manually measured for growth characteristics such plant height (cm), dry matter accumulation (g/plant), and more.

After cleaning and winnowing, grain yield per hectare was estimated and expressed in kilograms per hectare. Each net plot's leftover yield was measured and expressed, after full drying under the sun for 10-days, in tons per acre.

Statistical Analysis:

The data was computed and analyzed by following statistical method of Gomez and Gomez (1984). and benefit cost ratio was worked out after price value of seed with stover, and total cost included in crop cultivation

RESULTS AND DISCUSSION

Effect of Spacing and nitrogen management on Yield and Yield Attributes

Number of capsules/plants

The statistical analysis of number of capsules produced by each plant found a significant influence. At a spacing of 30 cm × 10 cm + (50% RDN + 50% N via FYM + Biofertilizer (*Azotobacter*)), the plant produced largest number of capsules (78.69). Nonetheless, it was shown that the spacing of 30 cm × 10 cm + (50% RDN + Biofertilizer (*Azotobacter*)) and 40 cm × 10 cm + (50% RDN + 50% N via FYM + Biofertilizer (*Azotobacter*)), were statistically at par with spacing of 30 cm × 10 cm + (50% RDN + 50% N via FYM + Biofertilizer (*Azotobacter*)). A sufficient quantity of sunlight absorption promotes efficient photosynthesis. With restricted spacing and dense plant population lower yield attribute values were attained. Ample application of nitrogenous fertilizers will not only enhance the yield but also support soil N status encourage productivity. Several records show that in sesame, use of Nitrogenous fertilizers effect in remarkable rise in growth and yield variables and seed yield.

Number of seeds/ capsules

Statistically, treatment T3 (spacing 30 cm × 10 cm + 50% RDN + 50% N via FYM + biofertilizer (*Azotobacter*)) outperformed the other treatments and had the maximum number of capsules per plant (36.39). Both the T2 treatment (spacing 30 cm × 10 cm + 50% RDN + biofertilizer (*Azotobacter*)) and the T6 treatment (spacing 40 cm × 10 cm + 50% RDN + 50% N via FYM + biofertilizer (*Azotobacter*)). These results are in line with those of Yadav and similar (2007). Since it improves plant photosynthesis when handled appropriately, nitrogen is an essential part of chlorophyll. Higher nitrogen concentrations have been shown to increase dry matter because increased photosynthetic activity leads to the production of more photosynthate. In sesame, Shinde et al. (2011) discovered that increasing the quantity of nitrogen led to the plant accumulation of dry matter.

Seed yield (kg/ha)

The grain yield showed that treatment T3 (spacing 30 cm × 10 cm + 50% RDN + 50% N via FYM + Biofertilizer (*Azotobacter*)) outperformed the other treatments and provided the maximum quantity of seed yield (552.38 kg/ha). While treatment T2 (Spacing 30 cm × 10 cm + 50% RDN + Biofertilizer (*Azotobacter*)) was observed to be statistically equivalent to treatment T3 (spacing 30 cm × 10 cm + 50% RDN + 50% N via FYM + Biofertilizer (*Azotobacter*)). Perhaps less competition exists for nutrients, moisture, and light. Adequate sunlight absorption promotes efficient photosynthesis, which causes more photosynthates to accumulate across a larger region. With restricted spacing and a dense plant population, lower yield attribute values were attained. Ogundare (2015), Patra, and Mishra found findings that were comparable to these (2000).

Stover yield (kg/ha)

The yield of sesame stover was also influenced by the usage of spacing and nitrogen management. The treatment T10 (Control with specified spacing and RDN) produced the greatest recorded stover yield (2602.6 kg/ha), whereas the treatment T1 (Spacing 30 cm 10 cm + 50% RDN + 50% N via FYM) produced the lowest reported yield (1920.6 kg/ha). Perhaps less competition exists for nutrients, moisture, and light. A sufficient quantity of sunlight absorption promotes efficient photosynthesis, which causes more photosynthesis to accumulate across a larger region. With restricted spacing and a dense plant population, loweryield attribute values were attained. Potassium is involved in a variety of physiological processes, such a s protein synthesis and enzyme activation. Similar findings were reported by Nayek et al. (2014) and Preeti (2010).

CONCLUSION

It can be concluded that with spacing of 30 cm × 10 cm and 50% recommended dose of nitrogen along with 50% nitrogen through FYM and seeds treated with Biofertilizer (*Azotobacter*) was found more productive (552.38 kg/ha) seed yield, as well as economically viable with net return (30,221.00 INR/ha), gross return (71809.00 INR/ha), and benefit : cost ratio (1.73). therefore it is profitable and can be suggested to farmers.

REFERENCES

- Alwal, M. A. and Aktar, L. (2015).** Effect of row spacing on the growth and yield of peanut (*Arachis hypogaea* L.) stands. *International Journal of Agriculture, Forestry and Fisheries*. **3**(1): 7-11.
- Anusha, O., Reddy, G. K., Sumathi, V., Reddy, P. V. and Reddy, A. P. K. (2018).** Organic Production for Sustained Productivity of Rabi Groundnut. *Andhrapradesh Journal of Agricultural Sciences*. **4**(1):56-61.
- Arif, M., Shehzad, M. A. and Mushtaq, S. (2012).** Inter and Intra Row Spacing on Growth, Seed yield and Oil contents of White Mustard (*Sinapis alba* L.) under Rainfed conditions. *Pakistan Journal of Agricultural Sciences*. **49**(1):21-25.
- Babar, S. and Dongale, J. H. (2011).** Soil fertility status as influenced by integrated nutrient management in mustard-cowpea-rice cropping sequence in lateritic soils of Konkan. *Asian Journal of Soil Science* **6**(1):33-38.
- Beenish, O., Ahmad, L., Hussain, A. and Lal, E. P. (2018).** Organic Manure & Biofertilizers: Effect on the Growth and Yield of Indian Mustard (*Brassica juncea* L.) Varieties. *Current Journal of Applied Science and Technology*. **30**(4): 1-7.
- Bedigian, D. and Vander Maesen L.J.G. (2003).** Slimy leaves and oily seeds; Distribution and Use of *Sesamum* spp. And *Ceratotheca sesamoides* (Pedaliaceae) in Africa. International Workshop, Nairobi, Prota Foundation Wageningen, The Netherlands: 271-274.
- Bouajila, K. and Sanaa, M. (2011).** Effects of organic ammendments on soil physico- chemical and biological properties. *J. Mater. Environ. Sci*. **2**(1): 485-49.
- Brar, Z.S., Bal, D.S. and John, A.S. (1998).** Influences of sowing dates, nitrogen and planting geometry on the performance of Gobhi Saron (*Brassica napus* spp. *olifera* var. *annua*). *Indian J. Agron*. **43**(1):133-137.
- Chandrasekaran, R., Somasundaram, E., Amanullah, M. M., Thirukumaran, K. and Sathyamoorthi, K. (2007).** Influence of Varieties and Plant Spacing on the Growth and Yield of Confectionery Groundnut (*Arachis hypogaea* L.). *Research Journal of Agriculture and Biological Sciences*. **3**(5): 525-528.

- Chakraborty, P., Bairagya, M. D., Sarkar, S., Gulati, J. M. L., Santra, G. H., Nayak, N. and Sahoo, B. K. (2021).** Effects of Irrigation and Nutrient Management on Summer Sesame (*Sesamum indicum* L.). *International Journal of Current Microbiology and Applied Sciences*. **10**(10): 212-220.
- Chetan, K. K., Sagar, M., Tanmoy, S., Monalisha, P. and Lalchetti, S. (2022).** Growth and productivity of sesame (*Sesamum indicum* L.) as influenced by spacing and nitrogen levels. *Crop research*. **57**(3): 190-194.
- Dalei, B. B., Panda, S., Kheroar, S. and Deshmukh, M. R. (2014).** Effect of Integrated nutrient management on growth, yield and economics of niger (*Guizotia abyssinica* L.). *Journal of Oilseeds Research*. **31**(1):46-48.
- Dinesh Kumar, R.B. Ardesna, B.R. Verma and Patel, A.K. (2017).** Growth and Yield Performance of Summer Sesamum Based Intercropping Systems. *Int.J.Curr.Microbiol.App.Sci*. **6**(6): 3341-334.
- Eshanullah., Akbar, N., Iqbal, A. and Mohsin, M. R. (2007).** Effect of different planting patterns on the growth and yield of two varieties of sesame. *Pakistan Journal of Agricultural Sciences*. **44**(4).
- Ganvit, J. B., Sharma, S., Surve, V. H. and Ganvit, V. C. (2019).** Effect of sowing dates and crop spacing on growth, yield and quality of linseed under south Gujarat condition. *Journal of Pharmacognosy and Phytochemistry*. **8**(1): 388-392.
- Ghosh, A. K., Duary, B. and Ghosh, D. C. (2013).** Nutrient Management in Summer Sesame (*Sesamum indicum* L.) and its Residual Effect on Black Gram (*Vigna mungo* L.). *International Journal of Bio-resource and Stress Management*. **4**(4):541-546.
- Gora, R., Kumar, S., Singh, P. K., Pooniyan, S., Choudhary., Dhaka, P. S. and Gora, M. K. (2022).** Effect of integrated nutrient management on productivity and profitability of mustard (*Brassica juncea* L.). *The Pharma Innovation Journal*. **11**(2): 2239-2242.
- Haritha. G., Singh, R. and Indu, T. (2022).** Studies on growth and yield of sesame (*Sesamum indicum* L.) as influenced by spacing and foliar application of zinc. *The Pharma Innovation Journal*. **11**(3): 2366-2369.
- Helen, J. Leopoldo, G. Rene, J. and Gerry, G. (2006).** Nature Farming Manual, Resource Efficient Agricultural Production-Canada, pp1-36.
- Ali GM, Yasumoto S, Seki-Katsuta M (2007).** et al. Assesment of genetic diversity in sesame (*sesamum indicum*). Detected by amplified fragment length polymorphism markers *Electronic J. Biotechnology*. 2007;10(1):0717-3458.
- Patil S G, Leva R L, Patil H R, Chaudhari (2017).** N N . Effect of spacing and nutrient management on summer sesame (*sesamum indicum*) under south Gujarat conditions. *Indian J Agricultural sciences*. 2017;88(4):00-70867
- Ghose, D.C. and Sen, J.C. (1980).** Analysis of yield components of sesamum as influenced by nitrogen and phosphorus fertilization. *Madras Agric. J.*, **67**(10) : 9

Ogundare, S.K., Aydele, F.G. and Oloniha, J.A. (2015). Effect of time of sowing and urea application rates on the growth and yield of two varieties of sesame (*Sesamum indicum* L.) in Ejiba Kogi state, Nigeria. Nigerian Journal of Agricultural, food and Environment. 11(4) : 118-123.

Nayek SS, Brahmachari K, Choudhary MR(2014). Integrated approach in nutrient management of sesame with special reference to its yield, quality and nutrient uptake. The Bioscan. 2014; 9(1):101-105.

Preeti (2010). Development of organic nutrient management for sesame *Sesamum indicum* L. M. Sc (Ag.) Thesis, 2010.

Shinde SD, Raskar BS, Tamboili BD. Effect of spacing and sulphur levels on productivity of sesame (*Sesamum indicum* L.) under summer condition. Journal of Maharashtra Agriculture University. 2011;36(1):28-31

H. Abou-Gharbia (1997) et al. Effects of processing on oxidative stability of sesame oil extracted from intact and dehulled seeds J. Am. Oil Chem. Soc. 74,215-221 (1997)

Patra, A. K. and Mishra,(2000). Effect of variety, nitrogen and spacing on yield attributes and yields of sesame (*Sesamum indicum* L.) during post-rainy season. J. Oilseeds Res., 17(1): 113-116.

Caliskan, S., M. Arslan, H. Arioglu, and Isler, N. 2004. Effect of planting method and plant population on growth and yield of sesame (*Sesamum indicum* L.) in a Mediterranean type of environment, Asian J. Plant Sci.3, 610-613.

Z.Gebregergis and M Amare, Effect of nitrogen fertilization on the growth and seed yield of Sesame (*Sesamum indicum* L.) Hindawi, International journal of Agronomy., 5027254 (2019).

Table 1. Effect of spacing and nitrogen management on yield attributes of summer sesame.(*sesamum indicum*)

| Treatment combinations | Yield and Yield Attributes | | | | | |
|--|----------------------------|--------------------------|-----------------|--------------------|----------------------|-------------------|
| | No.of capsules/ Plant | No. of seeds/ capsule | Test Weight (g) | Seed Yield (kg/ha) | Stover Yield (kg/ha) | Harvest Index (%) |
| T1. Spacing 30cm × 10cm + 50% RDN + 50% N through FYM | 70.89 | 32.72 | 3.34 | 409.65 | 1920.6 | 15.03 |
| T2. Spacing 30cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>) | 75.61 | 35.28 | 3.38 | 451.64 | 1980 | 14.86 |
| T3. Spacing 30cm × 10cm + (50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>)) | 78.69 | 36.39 | 3.38 | 552.38 | 2052.3 | 17.51 |
| T4. Spacing 40cm × 10cm + 50% RDN + 50% N through FYM | 68.93 | 29.68 | 3.31 | 401.82 | 2250.6 | 15.15 |
| T5. Spacing 40cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>) | 72.54 | 34.12 | 3.34 | 438.69 | 2271 | 14.92 |
| T6. Spacing 40cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>) | 74.69 | 34.54 | 3.35 | 443.26 | 2501.9 | 14.76 |
| T7. Spacing 50cm × 10cm + 50% RDN + 50% N through FYM | 66.73 | 27.86 | 3.30 | 396.89 | 2560.8 | 16.70 |
| T8. Spacing 50cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>) | 68.61 | 28.73 | 3.30 | 399.51 | 2315.6 | 16.29 |
| T9. Spacing 50cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>) | 69.76 | 31.33 | 3.32 | 419.36 | 2587.9 | 15.59 |
| T10. Spacing 30cm × 10 cm + 50: 40: 30 NPK Kg/ha (Control) | 65.35 | 25.56 | 3.29 | 387.69 | 2602.6 | 16.80 |
| F test | S | S | NS | S | S | NS |
| SEm(±) | 0.81 | 0.48 | 0.04 | 8.10 | 2.60 | 0.28 |
| CD (P=0.05) | 2.40 | 1.43 | - | 24.06 | 7.72 | - |

Table 2. Effect of spacing and nitrogen management on Economics of summer sesame.(*sesamum indicum*)

| S.no | Treatments | Cost of cultivation (INR/ha) Indian rupees | Gross returns (INR/ha) | Net returns (INR/ha) | B:C ratio (Benefit cost ratio) |
|------|--|--|------------------------|----------------------|--------------------------------|
| T1. | Spacing 30cm×10cm+50% RDN+50% N through FYM | 30878.00 | 53255.00 | 22376.00 | 1.72 |
| T2. | Spacing 30×10cm+50% RDN+Biofertilizer(<i>Azotobacter</i>) | 36273.00 | 58713.00 | 22440.00 | 1.62 |
| T3. | Spacing 30cm×10cm+50% RDN+50% N through FYM+Biofertilizer (<i>Azotobacter</i>) | 41588.00 | 71809.00 | 30221.00 | 1.73 |
| T4. | Spacing 40×10cm+50% RDN+50% N through FYM | 30788.00 | 52237.00 | 21458.00 | 1.70 |
| T5. | Spacing 40×10cm+50% RDN+Biofertilizer(<i>Azotobacter</i>) | 36173.00 | 57030.00 | 20857.00 | 1.58 |
| T6. | Spacing 40cm×10cm+50% RDN+50% N through FYM+Biofertilizer (<i>Azotobacter</i>) | 41488.00 | 57624.00 | 16136.00 | 1.39 |
| T7. | Spacing 50×10cm+50% RDN+50% N through FYM | 30678.00 | 51596.00 | 20917.00 | 1.68 |
| T8. | Spacing 50×10cm+50% RDN+Biofertilizer(<i>Azotobacter</i>) | 36073.00 | 51936.00 | 15863.00 | 1.44 |
| T9. | Spacing 50×10cm+50% RDN+50% N through FYM+Biofertilizer (<i>Azotobacter</i>) | 41388.00 | 54517.00 | 13129.00 | 1.32 |
| T10 | Spacing 30cm × 10cm + 50 : 40 : 30 NPK kg/ha | 30227.00 | 50400.00 | 20713.00 | 1.67 |