

Correlation between Vitamin D and Severity of Periodontitis in CKD patient

Abstract

Introduction: As the disease progresses, CKD patients become susceptible to a broad range of adverse health outcomes, including periodontitis. Both CKD and periodontitis have been associated with vitamin D deficiency. Vitamin D, known for its role in bone health, also plays a significant role in regulating the immune system, which is why it has been postulated that vitamin D is linked to the severity of periodontitis in CKD patients. The aim of this review is to evaluate the relationship between vitamin D levels and the severity of periodontitis in CKD patients.

Study design: A prospective case control study

Place and duration of study: Department of Medicine SwaroopRani Nehru Hospital, Prayagraj, India between July 2021 to June 2022

Material and Method: A total 50 chronic kidney disease patients were included on the basis of inclusion and exclusion criteria. Periodontal evaluation was done and classified it by using CDC-AAP classification. Cases and control were defined after periodontal evaluation. Blood sample was taken for investigation like serum vitamin D, serum creatinine and other markers.

Result: Out of 50 chronic kidney disease patients, 40 CKD patient (80%) had periodontitis changes considered as CASES and 10 (20%) CKD patient did not show any periodontitis changes considered as CONTROL. Cases had lower mean \pm Standard deviation (SD) Serum vitamin D levels than controls (13.90 ± 8.16 versus 25.71 ± 12.68 ng/mL, $p < 0.002$). Mean Serum Vitamin D of the study population decrease significantly within mild, moderate and severe periodontitis (p value = 0.006).

Conclusion: Therefore, it is evident with our study that serum vitamin D levels were significantly lower in cases as compared to control. It is also observed that serum vitamin D levels decrease as the severity of periodontitis increases in patients of chronic kidney disease.

Keywords: chronic kidney disease, periodontitis, stage, severity, serum vitamin D

1. Introduction

Chronic kidney disease (CKD) is a progressive condition characterized by the gradual loss of kidney function, eventual kidney failure, and the need for renal replacement therapy. CKD patients often experience numerous complications, including an increased risk of periodontitis – a common inflammatory disease affecting the gum and bone around the tooth [1]. Low serum vitamin D levels have been identified as a potential risk factor for both CKD and periodontitis. Vitamin D is a fat-soluble vitamin that plays a crucial role in maintaining bone health and regulating calcium metabolism. Additionally, vitamin D has potent immunomodulatory properties, and its deficiency has been linked to a range of chronic diseases, including those affecting the oral cavity [2,3]. Periodontitis is a chronic inflammatory disease of the supporting tissues of teeth, caused by the interaction of bacterial plaque and the host immune response to gingival tissue. CKD patients have a higher risk of periodontitis development, progression and severity, possibly due to an impaired immune response, decreased healing capacity, or some other complex mechanisms related to the primary kidney disease. Several potential underlying mechanisms explain how vitamin D may modulate periodontitis in CKD patients. One possible factor is related to the immune system, a well-known role of Vitamin D, which may help reduce inflammation

40 and stimulate tissue healing. Additionally, vitamin D plays a role in maintaining the integrity of connective
41 tissues and promotes bone development and mineralization[4]. Some authors hypothesized that vitamin D may
42 play a role in improving osteoporotic changes of periodontal bone during periodontitis development in CKD
43 patients[5]. A better understanding of the relationship between vitamin D deficiency, CKD, and periodontitis
44 may lead to the development of targeted interventions aimed at improving oral health for CKD patients,
45 improving their quality of life, and reducing the risks of systemic complications.

46 We postulate that an inadequate level of vitamin D promotes the occurrence of periodontitis in patients with
47 CKD. Therefore, the objective of this study was to correlate between serum vitamin D and periodontitis in CKD
48 patients.

49 **2. Material and Method**

50 **2.1 Study design**

51 This is a prospective case-control study has been approved by the ethics committee of MLN Medical College
52 and Associate Hospitals Prayagraj with registration no. ECR/922/inst./UP/2017 issued under rule 122DD/of the
53 drugs and cosmetics rule 1945. The study was carried out in the Department of Nephrology MLN Medical
54 College Prayagraj India between July 2021 to June 2022. A total of 50 patients suffering from chronic kidney
55 disease participated in this study. The nature of the study was explained to the patients, and a written informed
56 consent was obtained from them. The inclusion criteria comprised of all adults (age > 18 years), male and female
57 patients. Patients pre-diagnosed CKD evaluated and recorded. Patient not on dialysis. Exclusion Criteria
58 comprised of: Individual who underwent periodontal therapy, malignancy, HIV, hepatitis, upper respiratory tract
59 infections, pregnant And lactating women , Other systemic Infection like AKI and pancreatitis and intake of
60 vitamin d in previous 3 month .

61 **2.2 Study Procedure**

62 Medical history and clinical examination was taken and recorded the CKD & its stages by measuring eGFR and
63 USG abdomen. Laboratory assays of serum BUN, serum creatinine and serum vitamin D were measured. Dental
64 examinations were performed using a Williams-USA probe and a dental mirror, including examination of
65 clinical attachment loss (CAL), pocket probing depth (PD) were measured and recorded. All four aspects of the
66 gums (facial, mesial, distal, and lingual) were evaluated for inflammation. To check the probing depth (PD), a
67 probe was inserted parallel to the longitudinal axis of the teeth and moved around tooth and then the numbers
68 obtained for each tooth are added. The sum was divided by the number of measured areas and to check for the
69 presence or absence of CAL, a probe was inserted into the gingival sulcus and the CEJ distance to the depth of
70 the pocket . By recording the pocket depth and clinical loss of attachment define periodontal disease by using
71 Centers for Disease Control and Prevention (CDC- AAP).

72 **2.3 Statistical analysis**

73 The data was collected on a semi-structured questionnaire and records all test reports were maintained. All
74 observation was made under direct supervisor. Categorical variables were presented in number and percentage
75 (%) and continuous variables were presented as mean and SD. Quantitative variables were compared using
76 Mann Whitney U test / unpaired T Test as appropriate between two groups. Qualitative variables were compared

77 using chi-square test and fisher,exact test as appropriate. To measure the strength of association between two
 78 scale parameters using spearman correlation coefficient as appropriate. A P value of < 0.05 was considerably
 79 statistically significant. The Data was entered in MS Excel spreadsheet and analysis was done using statistical
 80 package for Social Science version 23.0.

81 **Result**


82 Total 50 CKD patient included in the study by using inclusion and exclusion criteria

83 **Table 1: Distribution of study population**

Groups	Description	Frequency	Percent
Cases	CKD (Chronic Kidney Disease) +Periodontitis	40	80.0
Controls	CKD (Chronic Kidney Disease)	10	20.0
Total		50	100.0

84

85 Table 1 reveals that distribution of study population in which out of 50 CKD patients,40 patients showed
 86 periodontitis changes considered as case and 10 patient did not showed periodontitis considered as control.
 87 Figure 1 illustrated below showing the percentage of different stages of CKD in cases and control
 88



89

90 **Fig. 1: Bar graph showing the percentage of different stages**
 91 **of CKD in cases and control.**

92 Figure 1 reveals that in cases the percentage of Stage 5, Stage 4 and Stage 3 are 80.0%, 15.0% and
 93 5.0% respectively. In controls the percentage of Stage 5, Stage 4 and Stage 3 are 80.0%, 0.0% and 20.0%
 94 respectively. It shows an insignificant difference between the groups (p value = 0.153). Table 2 and figure 2
 95 depicted below showing the distribution of CKD stages based on severity of periodontitis in cases.


96

97

Table 2: Distribution of CKD stages based on severity of periodontitis in cases

CKD stage	Periodontal status						χ^2 value (df)	p-value
	mild		moderate		severe			
	N	%	N	%	N	%		
stage 3	2	10.5%	0	.0%	0	.0%	5.12 (4)	0.272
stage 4	1	5.3%	2	18.2%	3	30.0%		
stage 5	16	84.2%	9	81.8%	7	70.0%		

99



100

101

102

Figure 02: Bar graph showing the percentage of CKD stages with respect to the severity of periodontitis.

103

Table 2 and figure 2 reveals that the CKD stage 5 more associated with periodontitis irrespective of their severity in study population. In mild cases of periodontitis, the percentage of Stage 5, Stage 4, and Stage 3 are 84.2%, 5.3%, and 10.5% respectively. In moderate and severe cases of periodontitis; the percentage of Stage 5 was 81.8% and 70.0% respectively. Table 3 given below showing the distribution of serum vitamin d between the cases and control.

108


109

110

Table 3: Distribution of Serum vitamin D between the groups

Marker	Groups			Mann Whitney U test	
	Cases (Mean ±SD)	Control (Mean ±SD)	Total (Mean±SD)	Z value	P value
Vitamin D	13.90±8.16	25.71±12.68	16.26±10.26	-3.094	0.002

111



112

Fig. 3:Box & Whisker diagram showing the mean & standard deviation of Serum vitamin D in cases &control

115 Serum vitamin D of both groups based on their mean(±SD) is summarized in Table 3. The mean(±SD) serum
 116 vitamin D of cases is found to be 13.90±8.16 and controls is 25.71±12.68, the overall mean(±SD) of serum
 117 vitamin D is found to be 16.26±10.26. There is a significant difference found in both groups (p value = 0.002).
 118 (Fig. 3). Table 4 illustration below showing the relationship between serum vitamin d and severity of
 119 periodontitis in CKD patients.

120

121

122

Table 4: Relation between Serum vitamin D and severity of periodontitis

Serum Marker	Periodontal status			ANOVA test	
	Mild (Mean ±SD)	Moderate (Mean ±SD)	Severe (Mean ±SD)	F value	P Value
Serum vitamin D	20.02±11.20	12.12±5.53	9.92±1.55	5.66	0.006

123

124 The above table shows that the mean Vitamin D of the study population decreases significantly within mild,
125 moderate, and severe periodontitis patients (p value = 0.006).

126 Discussion

127 Oral health of CKD patients is often poor due to poor oral hygiene because of their poor condition. Due to the
128 role of periodontal disease in the development of systemic inflammation and the effectiveness of serum vitamin
129 D between CKD and periodontitis in this regard there are very few literatures with different materials and
130 method in different countries which may not be applicable in our country so we designed this study which is the
131 first from our country. The aim of this study to find out correlation between serum Vitamin D and severity
132 of periodontitis in CKD patients. The present study included 50 patients in which 80% of CKD patient showed the
133 periodontitis changes. The findings of this study revealed that the mean (±SD) of Vitamin D of cases were found
134 statistically significant with respect to control (p value = 0.002) which was consistent with Bastos et al who
135 studied in smaller size with comparison to our study but serum vitamin d level found to be significant in cases
136 with respect to control p value < 0.01. Similarly, [Rabienejad et al., \(2022\)](#) found that the difference in mean serum
137 vitamin D levels between the case and control groups was 33.66, which was statistically significant (P-value
138 < 0.001) (Rabienejad et al., 2022) but this study done in dialysis patients and our study specially on pre-
139 dialysis patient. Current study also showed that the serum vitamin D level of the cases decreases significantly as
140 the severity of periodontitis increases.

141 Conclusion

142 Therefore our study was evident despite excluding several confounding factors such as smoking, use of Anti-
143 inflammatory medications, recent antibiotic Use, pregnancy, cancer, infection with human immunodeficiency
144 virus, uncompensated diabetes, and fever of unknown origin or other infections, periodontal treatment within the
145 last 6 months, and cases of aggressive or acute periodontal disease. In summary, our findings suggest that
146 vitamin D deficiency predisposes patients with CKD to develop CP, possibly by limiting the patient's
147 inflammatory and immune response against bacterial invasion of the periodontium. A sufficient level of vitamin
148 D is important in maintaining a healthy periodontium and reducing the consequences of periodontitis.

149

150

151 **DISCLAIMER**

152 The products used for this research are commonly and predominantly use products in our area of research and
153 country. There is absolutely no conflict of interest between the authors and producers of the products because
154 we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge.
155 Also, the research was not funded by the producing company rather it was funded by personal efforts of the
156 authors.

157 **CONSENT**

158 As per international standard or university standard, patient's consent has been collected and preserved by the
159 authors.

160 **ETHICAL APPROVAL**

161 Ethical approval was sought and obtained from the Ethics Committee of MLN Medical College, Prayagraj.

162 **Abbreviations**

163 CKD – chronic kidney disease

164 P- periodontitis

165 PD – pocket depth

166 Mild p – mild periodontitis

167 Moderate p – moderate periodontitis

168 Severe P – severe periodontitis

169 CAL- clinical loss of attachment

170 Serum vit d – Serum vitamin d

171 **References**

- 172 1. Bastos JA, Diniz CG, Bastos MG, Vilela EM, Silva VL, ChaoubahASouza-Costa DC, et al.
173 Identification of periodontal pathogens and severity of periodontitis in patients with and without
174 chronic kidney disease. Arch Oral Biol 2011;56:804-11.
- 175 2. Dietrich T, Joshipura KJ, Dawson-Hughes B, Bischoff-Ferrari HA. Association between serum
176 concentrations of 25-hydroxyvitamin D3 and periodontal disease in the US population. Am J Clin Nutr
177 2004;80:108-13.
- 178 3. Nishida M, Grossi SG, Dunford RG, Ho AW, Trevisan M, Genco RJ. Calcium and the risk for
179 periodontal disease. J Periodontol 2000;71:1057-66.

180
181

4. Hildebolt CF, Pilgram TK, Dotson M, Armamento-Villareal R, Hauser J, Cohen S, et al. Estrogen and/or calcium plus vitamin D increase mandibular bone mass. *J Periodontol* 2004;75:811-6.

182
183
184

5. Miley DD, Garcia MN, Hildebolt CF, Shannon WD, Couture RA, Anderson Spearie CL, et al. Cross-sectional study of vitamin D and calcium supplementation effects on chronic periodontitis. *J Periodontol* 2009;80:1433-9.

185

UNDER PEER REVIEW