

Original Research Article

Rapid method for genomic DNA isolation of mungbean [*Vigna radiata* (L.) Wilczek]

Abstract

The most significant pulse crop in India is the mungbean [*Vigna radiata* (L.) Wilczek], which constitutes almost 16 % of the national pulse area. For biotechnological interventions, it is prerequisite to develop a rapid and cost-effective method for genomic DNA isolation suitable for PCR. In present investigation, modified CTAB method was used to extract the genomic DNA from the leaf tissues of mungbean. By crushing the leaves of several mungbean genotypes in a pre-heated CTAB extraction buffer, genomic DNA was extracted. Further, SSR markers were used to assess the efficiency of the extracted DNA. The SSR marker-based PCR amplification findings showed that the DNA extracted using this approach was of good quality and suitable for SSR analysis.

Keywords: Mungbean [*Vigna radiata* (L.) Wilczek], Genomic DNA, Cetyl Trimethyl Ammonium Bromide (CTAB), SSR Marker, PCR Amplification.

Introduction

Mungbean [*Vigna radiata* (L.) Wilczek] is one of the most important pulse crops in the Asiatic region. Mungbean, often known as greengram, is a crucial source of protein for humans and the health of the soil since it fixes atmospheric nitrogen. It is well-liked by people all over the world because of its high protein content, simple digestion, and lack of flatulence (Anwar et al., 2007). The grains, whether split or whole, are made into dal or flour. Cattle are given straw and husk as feed. Mungbean's susceptibility to biotic stresses is the main reason for the crop's low yield (Mishra et al., 2020). The advent of DNA markers resulted in an inevitable revolution in plant genetics and breeding (Amiteye 2021). Majority of DNA extraction protocols suggests the use of liquid nitrogen which is expensive and difficult to obtain and when required.

The recent deployment of molecular markers has tremendously increased the requirement of rapid and PCR quality DNA in very short period of time for crop improvement (Prasad et al., 2018). While the availability of DNA markers associated with qualities of interest has expanded, current DNA extraction technologies have limited the number of samples that can be processed. (Prasad et al., 2018; Rehman et al., 2007). The DNA extraction methods from mungbean are not time efficient, since they require several steps, like the tissues to be grounded in liquid nitrogen, followed by precipitation of the DNA pellet in ethanol, washing and drying the pellet, etc. However, a plethora of DNA extraction methodologies have been reported and although some of these are undoubtedly rapid, many require the use of expensive, often environmentally hazardous chemicals, and specialized laboratory equipment (Kang and Yang, 2004; Amani et al., 2011). The need for a rapid and simple procedure is urgent, especially when hundreds of samples need to be analyzed. Here, we describe a rapid, cheaper and efficient genomic DNA extraction method for PCR amplification from leaf tissue

Materials and Methods

Materials used

Leaves samples (young immature leaves) of mungbean plant were used for DNA extraction. The seeds were germinated under favourable conditions with adequate water and sunlight. After 10-12 days, leaves (0.5 g) were taken for genomic DNA extraction.

Reagents used

- 2X CTAB Extraction Buffer: The extraction buffer consisted of 100 mM Tris-HCl (pH 8.0), 20 mM EDTA (pH 8.0), 1.4 M NaCl, 1% (w/v) PVP, and 2% CTAB. Warm CTAB buffer in a water bath for 5 minutes just before using it.
- 5X CTAB Buffer
- 70% Ethanol
- Chloroform: Isoamyl alcohol (24:1)
- 0.5 M EDTA (pH-8.00)
- Isopropanol
- Ethanol

- 1M Tris-Cl (pH-8.00)
- 0.1X TE Buffer: 1 mM Tris-HCl (pH 8.0), and 0.1 mM EDTA (pH 8.0)
- 10X TBE Buffer

Genomic DNA Extraction

The leaf tissues are ground in a clean mortar and pestle with the addition of 2ml of warm 2X CTAB extraction buffer. The ground sample was transferred into a nucleases free microcentrifuge tube(2 ml) and incubated in a water bath for 5 minutes at 65°C. After incubation, equal amount of chloroform: isoamylalcohol mixture (24:1) were added, and mixed by gentle inversion followed by centrifugation at 12000 rpm for 5 minutes. The supernatant (uppermost aqueous layer) was carefully transferred to a new microcentrifuge tube using a pipette. One fifth volume of 5X CTAB Buffer was added to the supernatant and mixed well followed by extraction through chloroform: isoamylalcohol as described above. DNA was precipitated by addition of either ethanol (2.5 vol) or isopropanol (0.6 vol) and incubated on ice for 10-15 min. DNA pellet was obtained by centrifugation at 12,000 rpm for 5 min. Supernatant was discarded and DNA pellet was washed with 70% ethanol (1 ml) to remove excess salt. The washed DNA pellet was air dried and dissolved in 30 μ l of nucleases free water and stored at 4°C. The quality of extracted genomic DNA from mungbean genotypes were analysed using agarose gel electrophoresis (0.8 %) percent,

Comment [KKC1]: A nuclease or nucleases




Figure 1. Overview of genomic DNA extraction steps from the leaves of mungbean. (a) Grown mungbean seedlings; (b) grinding of leaf tissue in warm DNA extraction buffer in mortar with pestal; (c) Three layers separation after Chloroform : isoamyl alcohol treatment, (d) carefully transferred upper or aqueous phase into new micro centrifuge tubes, e) precipitation of thread like structure of DNA after addition of either ethanol or isopropanol, f) visualization of genomic DNA band on agarose gel

SSR (Full form) analysis

The extracted genomic DNA was diluted in a 1:10 ratio and 1 μ l of diluted DNA was used for SSR marker analysis. The SSR amplification was carried out as described previously (Mishra et al., 2020). In brief, SSR marker amplification was set up in total 10 μ l system volume containing 5 μ l All-in-One PCR supermix (Helix), 0.2 μ M forward and reverse primer and 50 ng of genomic DNA in a thermal cycler. The amplification reaction involved an initial 940C for 2 min for denaturation followed by 30cycles of 1 min at 94°C, 30 sec at 51°C, 1 min at 72 °C and final extension at 72°C for 3 min. Amplified products were electrophoresized on 2.5 % agarose gel containing 0.5 μ l ethidium bromide and the gels were visualized under UV light. The list of SSR markers used in this study is mentioned in Table 1.

Comment [KKC2]: ?

Table 1 : List of SSR markers used in this study

S. No	Primer Name	Forward Primer Sequence (5 to 3')	Reverse Primer Sequence (5' to 3')
1.	MBM00389	GCAAGTGTTTCGACGAGGTT	GCTACACGCCATTGTTTTGA
2.	MBM0003	GGAAGGGAAGGAAAAAGGAA	ATCCAGAAACCGAATCGTTG
3.	MR7322B	TCAGTCAGTGTCGATAGCATA GC	GACACAGAGAGAGAGAGAGA G

Results and Discussion

In recent year use of SSR marker in diversity as well as marker assisted breeding (MAB) programme in Mungbean have tremendously increased (Mishra et al., 2020; Tabasum et al., 2020). In the present investigation, we aimed to develop a rapid and high throughput DNA isolation protocol for molecular marker analysis in mungbean. The developed protocol was found to be suitable for the rapid isolation of DNA and the suitability of isolated DNA for downstream marker analysis was checked using an SSR marker. The rapid method of DNA

isolation has also been developed in other crop plants (Prasad et al., 2018; Ahmadikhah, 2009; Liang et al., 2016).

In present investigation four mungbean genotypes were used for SSR marker analyses. The amplification were observed in all the four genotypes in all the three SSR marker used (Figure 2).




Figure 2. PCR amplification for SSR markers in mungbean genotypes.

The reproducibility of this method was checked by repeating the extraction of DNA and PCR several times.

Numerous protocols for DNA isolation are available in literature but most of them either uses liquid nitrogen and other carcinogenic reagents (Sharma et al., 2002). Our modified protocol doesn't require use of liquid nitrogen and phenol. Additionally, this modified protocol doesn't require any specialized apparatus (e.g. the matrix mill) (Hill-Ambroz et al., 2002). The commercially available kits are one of the alternatives to traditional methods of DNA isolation, but their higher cost is the limiting factor (Kang and Yang, 2004; Ahmed et al., 2009). The major

advantage of our modified protocol is its rapidity without the use of liquid nitrogen and phenol, a dangerous organic solvent.

Conclusion

Our results clearly demonstrated that the quality of extracted DNA is good enough to perform SSR marker analysis. This method has provided a convenient, rapid, low-cost and safe DNA extraction method for SSR analysis in mungbean.

References

- Ahmadikhah, A. (2009) A rapid mini-prep DNA extraction method in rice (*Oryza sativa*). African Journal of Biotechnology. 8 (2): 323-327.
- Ahmed I, Islam M, Arshad W, Mannan A, Ahmad W, Mirza1 B. (2009) High-quality plant DNA extraction for PCR: an easy approach. J Appl Genet. 50:105-107.
- Amani J, Kazemi R, Ali Reza Abbasi AR, Salmanian AH. (2011) A simple and rapid leaf genomic DNA extraction method for polymerase chain reaction analysis. Iranian J Biotech. 9(1):69-71.
- Amiteye S. (2021) Basic concepts and methodologies of DNA marker systems in plant molecular breeding. Heliyon. 30: 7(10):e08093.
- Anwar F, Latif S, Przybylski R, Sultana B, Ashraf M. (2007) Chemical composition and antioxidant activity of seeds of different cultivars of mung bean. J Food Sci. 72(7): S503–10.
- Handayani F, Wulandari RA, Murti RH. (2016) Genomic DNA extraction method from mature leaf of lai (*Durio kutejensis* Becc.). Agrivita. 38(1): 73-9.
- Hill-Ambroz KL, Brown-Guedira GL, Fellers JP. (2002) Modified rapid DNA extraction protocol for high throughput microsatellite analysis in wheat. Crop Sci. 42:2088-2091.
- Kang TJ, Yang MS. (2004) Rapid and reliable extraction of genomic DNA from various wild-type and transgenic plants. BMC Biotechnol. 4:20.
- Liang H, Deng Y, Wang C., Xu X. (2016) A high-throughput DNA extraction method from rice seeds. Biotechnology & Biotechnological Equipment, 30(1): 32-35
- Mishra A., Sahni S., Prasad BD., Kumar B., Kumar S. (2020) Molecular characterization of Mungbean [*Vigna radiata* (L.) Wilczek] genotypes for resistance against Mungbean Yellow Mosaic Virus (MYMV). Res. J. Biotech. 15(6): :123-134.
- Prasad BD, Sahni S., Kumar P (2018) Instant genomic DNA extraction method from safflower leaf for PCR Analysis. Journal of Pharmacognosy and Phytochemistry 2018; 7(2): 3500-3502.

Rehman A-U, Stodart B, Raman H. (2007) A high throughput, non-organic method for plant genomic DNA isolation. Pak J Bot. 2007; 39(3):831-840.

Sharma AD, Gill PK, Singh P (2002) DNA isolation from dry and fresh samples of polysaccharide-rich plants. Plant Mol Biol Rep. 20:415a-415f.

Tabasum A, Hameed A, Asghar MJ. (2020) Exploring the Genetic Divergence in Mungbean (*Vigna radiata* L.) Germplasm Using Multiple Molecular Marker Systems. Mol Biotechnol. 62(11-12):547-556.

UNDER PEER REVIEW