

Review Article

EMERGING WEED PROBLEMS UNDER CHANGING CLIMATIC CONDITION – A REVIEW

Abstract

Climate is a major force in Earth's environmental system, and even minor changes in climate can have complex and serious effects on the environment and nature. Atmospheric CO₂, temperature and precipitation are major important abiotic variables that directly affect weed physiology and growth. Weeds are more genetically diverse than crops, hence show enhanced growth and reproductive stability than crops. As with crops, weeds with C3 or C4 photosynthetic pathways exhibit differences in their response to changing climate. The C3 weeds (*Argemone mexicana*, *Chenopodium album* and *Phalaris minor* etc.) shows more response in photosynthesis to increased CO₂ level in atmosphere. Elevated CO₂ under sufficient water condition will lead to higher C3 weed competitiveness in C4 crops. So weed growth is favoured as CO₂ is increased.

Temperature changes resulted in an expansion of weeds, with some species shifting to higher latitude and altitude. Based on the differences in temperature optima for physiological processes, C4 *spp.* able to tolerate high temperature than C3 *spp.* Therefore, C4 weeds (*Digitaria ciliaris*, *Cyperus spp.*, *Echinochloa colona* etc.) benefit more than the C3 crops from any temperature increases that accompany elevated CO₂ levels. Normally, the mean temperatures increases over a long period, weeds expand their range into new areas. Precipitation is also important climate variable that influences the growth and distribution of weed vegetation. Although CO₂ concentrations and temperature rise in future are predictable, distribution and variation in the intensity of precipitation patterns remains much more uncertain. Moisture is a key factor required for weed seed germination and establishment. Climate change can affect the frequency and intensity of rainfall and result in occurrences of extreme events such as floods and droughts; consequently, weeds adapted to these conditions will have a higher competitive advantage.

Key words: *Climate Change, elevated CO₂, Temperature, Precipitation, C3 and C4 weeds*

Introduction:

The difficult of climate change is increasingly more receiving attention from scientists, public peoples, and policy makers. Climate change is now a broadly accepted term (Petit et al. 1999; Loss et al. 2011). Climate Change defines “a change of climate which is credited directly or indirectly to human activity by alters the compositions in the atmosphere negatively” (Robinson and Gross 2010). The atmospheric CO₂ concentration haverisen from about 280 ppm (pre-industrial period) to today’s about 390 ppm and it is expectedthat by the end of 21st century, it will reach to the levels of 600-700 ppm, if current emissiontrends continue (IPCC, 2001; Leegood, 2002). Climate models projected that the earth surface temperature is likely to rise in a range of 1.1 to 6.4°C during the 21st century due to the risingof carbon dioxide concentration in the atmosphere (IPCC, 2004; Houghton, 2001). There are concerns that global CO₂ enrichment will affectthe drymatter of weeds and crop yields directly or indirectly through global warming and its associated changes in climate such as alteration in precipitation, wind pattern, rise in sea level and moreflood and drought. It has been reported that elevated CO₂concentration cause an increases inphotosynthesis (Kendall *et al.* 1985; Ghannoum et al. 2000; Dass *et al.*, 2017), decrease in photorespiration and respiration (Ziska, 2000), different crop and weed species differ in their response toelevated CO₂ concentration levels (Poorter, 1993).

The various responses of C3 and C4 plants (weeds & crops) to elevated CO₂ and air temperature have important implications on crop/weed competition as most of the weeds are C4. But this basic idea that the most crops are C3 photosynthesis and most weeds are C4 photosynthesis, and hence weeds are mature and decay as earlier (Ziska, 2003; Pautasso *et al.* 2010; Singh *et al.* 2011).Evidently, there are four major C4 crops of economic importance (Maize, Pearlmillet, Sorghum, and Sugarcane), and many important C3 weeds (e.g. *Chenopodium album*, *Avena ludoviciana*, *Phalaris minor*, *Eclipta prostrate*, *Ammania baccifera* etc.).Generally, it is reported that plants with C3 photosynthetic pathways are expected to benefit more than C4 from CO₂ enrichment but inverse is true with rising temperature (Jinger *et al.*, 2016).

Climate change may bring changes in weed population and in their phenology (Patterson, 1995). Many weed species may expand their range and spread to new areas. In addition to its impact on agricultural weed, literature suggest that invasive species may

Comment [HUR1]: That depends on the scenario. It would be better to put the whole range from the most optimistic scenario to the most pessimistic.

Comment [HUR2]: Review the most current report of the IPCC 2022

Comment [HUR3]: Review the most current report of the IPCC 2022

become more of a threat in changing climate because of their strong response to elevated CO₂ and changing climate compared to other native species.

Moreover, some weed species which directly impact human health through allergic reactions, skin irritations, or internal poisoning have shown positive response to changing climate particularly to elevated CO₂ concentration by producing higher plant biomass, pollens or poisonous compounds (Dukes *et al.*, 2009; Singer *et al.*, 2013). Weeds may show a greater range of responses to increasing temperatures because of their wider gene pool compared with crops, which enables them to adapt to diverse environment conditions (Treharne, 1989). The degree of photosynthetic stimulation and growth response varies between C₃ and C₄ plants as the temperature increases (Morgan *et al.*, 2001).

Precipitation is another important climate variable that influences the growth and distribution of vegetation (Lobell and Burke, 2008; Robinson and Gross, 2010). Although CO₂ concentrations and temperature rise in future are predictable, distribution and variation in the intensity of precipitation patterns remains much more uncertain. Moisture is a key factor required for weed seed germination and establishment (Skov and Svenning 2004; Bergmann *et al.* 2010). Climate change can affect the frequency and intensity of rainfall and result in occurrences of extreme events such as floods and droughts; consequently, weeds adapted to these conditions will have a higher competitive advantage (Rodenburg *et al.*, 2011). Finally, the present reviews covered a weed emergence problem under changing climatic condition. Further investigations and research areas are needed in this concern.

Weeds – Characteristics

In this world totally 46 major crops are there but, over 410 “serious” weed species.

1. Produces larger number of seeds compare to crops
2. Most of the weed seeds are small in size and contribute enormously to the seed reserves.
3. Weed seeds germinate earlier and their seedlings grow faster than crop seeds.
4. The viability of seeds remains intact, even if they are buried deep in the soil.
5. Weed seeds do not lose their viability for years even under adverse conditions.
6. They flower earlier and mature ahead of the crop they infest.

7. They have the capacity to germinate under varied conditions, but very characteristically, season bound. The peak period of germination always takes place in certain seasons in regular succession year after year.
8. Weed seeds possess the phenomenon of dormancy
9. Most of the weeds possess C4 type of photosynthesis, which is an added advantage during moisture stress.
10. They possess extensive root system, which go deeper as well as of creeping type.
11. Quick response to available soil moisture and nutrients.
12. Tolerance to shading effects by the crops at the time of establishment
13. Mimicry: Resembles the crop plants, morphological characters are similar to the crop Plants.
14. Weed dispersal has no geographical boundaries. Common agents of weed dispersal are wind, water, animals, birds, organic manures, agriculture implements and human beings.

Climate Change on Weeds

The main parameters for climate change impacts on weeds include increased atmospheric CO₂ levels, temperatures, changed rainfall, more extreme weather, more frequent frosts, changed phenology and changed land use (Patterson, 1995; Ziska, 2016). The rate of response of invasive plants and weeds is expected to be faster than for other plants, including native species and crops. Secondly, climate change is likely to foster the appearance of a new set of weed species (Dukes *et al.* 2009; Singer *et al.* 2013).

Change in atmospheric CO₂, temperature, rainfall and wind will affect on Weed species distribution, growth and physiology, prevalence within weed and crop communities, affect crop-weed balance and lead to weed invasion (Patterson, 1995; Ramesh *et al.*, 2017; Anwar *et al.*, 2021).

Impact of climate change on weed biology (Ziska, 2000)

- Weed shifts and spread of invasive species
- Elevated CO₂ and its effects on weeds
- Effect of increased temperature on weeds
- Effect of change in precipitation on weeds

- Other environmental stresses and weed competitiveness.

Table 1 : Impact of climate change on risks of invasiveness (Bradley *et al.*, 2010)

Climate change factor	Risks of invasiveness
Elevated CO ₂	Increase
Rising temperature	Might increase or decrease
Changing precipitation regime	Might increase or decrease

Comment [HUR4]: There is no reference to this table in the text.

Weeds response to increasing of CO₂ levels

Elevated atmospheric CO₂

Weeds have a greater genetic diversity than crops. Consequently, if a resource (light, water, nutrients or carbon dioxide) changes within the environment, it is more likely that weeds will show a greater growth and reproductive response. Elevated atmospheric CO₂ improves the growth of plants due to increased efficiency (Ziska, 2008). The interaction between eCO₂ and growth is strongly influenced by the mode of energy capture (photosynthesis), with some plants (C₄ and CAM), including many grasses and succulents, being more efficient than others (C₃ plants) (Patterson, 1995; Battset *al.* 1997; Morison and Lawlor, 1999).

Comment [HUR5]: CO₂

Many of the weed species have the C₄ photosynthetic pathway. However, this argument does not consider the range of available C₃ and C₄ weeds present in any agronomic environment. Hence, if a C₄ weed species does not respond, it is likely that a C₃ weed species will (Morison and Lawlor, 1999). In addition, many growers recognize that the worst weeds for a given crop are similar in growth habit or photosynthetic pathway; indeed, they are often the same uncultivated or “wild” species, e.g. oat and wild oat, sorghum, rice and red rice. To date, for all weed/crop competition studies where the photosynthetic pathway is the same, weed growth is favoured as CO₂ is increased (Carter and Peterson 1983; Ziska, 2003).

Direct physiological effects of elevated CO₂ on photosynthesis and plant growth have been well documented. Plants with C₃ photosynthetic pathways are expected to benefit more than C₄ from CO₂ enrichment (Patterson and Flint, 1980). This differential response of C₃

and C4 plants to elevated CO₂ can have important implications on crop/weed competition as most of the weeds are C4. Therefore, it can be argued that because of C4 photosynthetic pathway of many weed species, they will show smaller response to elevated CO₂ relative to crops which are mostly C3 (Zangerl and Bazzaz, 1984; Ziska, 2003; Rogers et al. 2008). Therefore, it is important to examine crop/weed competition cropping system based to develop effective weed management practice for the emerging species in the changing climate scenario.

Impacts(Reddy, *et al.*, 2010)

1. Higher CO₂ : Stimulate Photosynthesis and Growth, Reduce ET and Increase WUE
2. Many C3 weeds have shown significant increases in growth, with substantial decreases in the yields of competing crops.
3. Rising CO₂ levels - Positive effect on the establishment and persistence of invasive species. (eg. *Bromus tectorum*).

Comment [HUR6]: define the terms ET and WUE.

INDIA: CO₂ VS Weeds

Common weeds species found in India such as *Ageratum conyzoides*, *Digitaria ciliaris*, *Cyperus spp.*, *Echinochloa colona*, *Paspalum orbiculare* and *Setaria glauca* and having C4 photosynthetic pathway will show smaller response in photosynthesis to increased CO₂ level in atmosphere (Singh, *et al.*, 2011).

Comment [HUR7]: The acronym for Versus is vs.

Comment [HUR8]: Cursive

Whereas weed species with C3 photosynthetic pathways like *Agropyron repens*, *Argemone mexicana*, *Chenopodium album*, *Phalaris minor*, *Poa annua* and *Rumex acetosella*, may show enhanced photosynthesis to increased CO₂ level in atmosphere, in addition, many of the worst weed for given crop are similar in growth habit or photosynthetic pathways for example, oat and wild oat, wheat and little seed canary grass, rice and wild rice, and hence weed growth is more favoured due to increased CO₂ level in the atmosphere (Patterson, 1985; Patterson, 1995; Mishra, 2003).

Table 2 : Important C3 and C4 weed species of rice and wheat crops in India

Crop	C3 weeds	C4 weeds
Rice	Weedy rice, <i>Scirpus spp.</i> , <i>Monochoria</i> , <i>Eclipta prostrata</i> , <i>Ammania baccifera</i>	<i>Echinochloa spp.</i> , <i>Cyperus spp.</i> , <i>Leptochloa chinensis</i> , <i>Bracharia</i>
Wheat	<i>Chenopodium album</i> , <i>Phalaris minor</i> , <i>Avena fatua</i> , <i>Convolvulus arvensis</i> , <i>Canada thistle</i>	<i>Cynadon dactylon</i>

Comment [HUR9]: There is no reference to this table in the text.

Table 3 : Response of C3 and C4 weeds to double atmospheric CO₂ levels

(Chandrasena, 2009)

Sn.	C3	Range of response biomass	C4	Range of response biomass
1	<i>Abutilon theophratsii</i>	1.0-1.52	<i>Amaranthus retroflexus</i>	0.9-1.41
2	<i>Bromus mollis</i>	1.37	<i>Andropogon virginicus</i>	0.8-1.17
3	<i>Bromus tectorum</i>	1.54	<i>Cyperus rotundus</i>	1.02
4	<i>Cassia obtusifolia</i>	1.4-1.8	<i>Digitaria ciliaris</i>	1.06-1.6
5	<i>Chenopodium album</i>	1.0-1.6	<i>Echinochloa crusgalli</i>	0.95-1.6
6	<i>Datura stramonium</i>	1.7-2.27	<i>Eleusine indica</i>	1.02-1.8
7	<i>Phalaris aquatica</i>	1.48	<i>Sorghum halepense</i>	0.56-1.1

Comment [HUR10]: There is no reference to this table in the text.

Table 4 : Shift in weed flora in India

Crop	Original weed flora	New weed flora
Wheat (first phase)	<i>Chenopodium spp.</i> , <i>Spergula arvensis</i> , <i>Anagallis arvensis</i>	<i>Phalaris minor</i> , <i>Avena fatua</i> , <i>Lolium temulentum</i>
Wheat (Second phase)	<i>Phalaris minor</i> , <i>Avena fatua</i>	<i>Lathyrus aphaca</i> , <i>Convolvulus arvensis</i> , <i>Medicago spp.</i> , <i>Cirsium arvense</i>
Rice	<i>Echinochloa spp.</i>	<i>Cyperus iria</i> , <i>Fimbristylis miliacea</i> , <i>Sphenoclea zeylanica</i>

Comment [HUR11]: There is no reference to this table in the text.

Sugarcane	Broadleaved weeds	Grassy weeds
------------------	-------------------	--------------

Higher photosynthetic rates at elevated CO₂ in C3 crops (Rice, Wheat, Soybean, etc.) mean they will respond more favourably to higher CO₂ levels than the C4 weeds (palmer amaranth (*A. palmeri*), water hemp (*Amaranthus rudis*), kochia (*K. scoparia*) (Elmore and Paul, 1983). In contrast, C3 weeds (lambsquarters, velvetleaf (*Abutilon theophrasti*), common ragweed (*Ambrosia artemisiifolia*) and giant ragweed (*Ambrosia tridca*)) will respond more favourably to increased CO₂ levels and offer stiffer competition to C4 crops (Maize, Sorghum, Sugarcane, etc.) (Tang *et al.* 2009)

C3 weeds in C4 crops

- Elevated CO₂ under sufficient water condition will lead to higher C3 weed competitiveness in C4 crops. At elevated CO₂ - Seed yield or total above ground biomass of sorghum was significantly reduced by C3 weeds (Ziska 2003).
- C3 weed dandelion (*Taraxacum officinale*) produced more fertile seeds and larger seedlings under elevated CO₂ (McPeck & Wang, 2007).

CO₂ VS C4 weeds and C4 Crops

Ziska and Bunce (1997) compared the effect of elevated CO₂ levels on the growth and biomass production of six C4 weeds (*Amaranthus retroflexus*, *Echinochloa crusgalli*, *Panicum dichotomiflorum* Michaux, *Setaria faberi* Herrm, *Setaria viridis*, *Sorghum halapense*) and four C4 crop species (*Amaranthus hypochondriacus*, *Saccharum officinarum*, *Sorghum bicolor*, and *Zea mays*). In these C4 species showed a significant increase in photosynthesis. The largest and smallest increases observed were for *A. retroflexus* (+30%) and *Z. mays* (+5%), respectively.

CO₂ VS Soybean - C3 weeds

Ziska (2000) evaluated the outcome of competition between 'Round-up Ready' Soybean (*Glycine max.*) and a C3 weed (lambsquarter, *Chenopodium album*) and a C4 weed (Redroot pigweed, *Amaranthus retroflexus*), grown at ambient and enhanced CO₂ (ambient + 250 μL L⁻¹). In a weed-free environment, elevated CO₂ resulted in a significant increase in soybean growth and seed yield, compared to the ambient CO₂ condition. However, soybean growth and seed yield were significantly reduced by the presence of either weed species at either level of CO₂. With lambsquarter, at elevated CO₂, the reduction in soybean

Comment [HUR12]: The acronym for Versus is vs.

Comment [HUR13]: The acronym for Versus is vs.

seed yield relative to the weed-free control increased from 28 to 39%; concomitantly, the dry weight of lambsquarter increased by 65%.

Conversely, for pigweed, soybean seed yield losses diminished with increasing CO₂ from 45 to 30%, with no change in the dry weight of the weed. This study suggests that rising CO₂ could alter yield losses due to competition from weeds, and that weed control will be crucial in realizing any potential increase in the yield of crops, such as soybean, as climate change occurs (Chandrasena, 2008).

Table 5 : Effect of elevated CO₂ on crop-weed interactions (Ziska, 2000)

Crop-weed interaction	Ambient CO ₂	Elevated CO ₂	Percentage change
Above ground biomass			
Soybean (C3)	340±13	448±14	+31.8
+C3 weed	261±18	297±29	+14.0
+C4 weed	204±17	329±27	+61.3
Seed yield			
Soybean (C3)	187±8	228±8	+21.9
+C3 weed	135±9	141±15	+4.4
+C4 weed	103±13	158±14	+53.4

(Total aboveground biomass and seed yield (±standard error) at maturity for soybean (g/m row) at ambient and elevated CO₂ (ambient +250 µl / l CO₂) when grown with or without the presence of a C3 weed (*Chenopodium album*) or a C4 weed (*Amaranthus retroflexus*).

CO₂ VS Redroot pigweed

The elevated CO₂ effect on Redroot pigweed (C4 weed), no change in biomass and yield loss in soybean decreased from 30% to 45% in competition with this weed. Growth of redroot pigweed in a sorghum field was favoured by increasing atmospheric CO₂ levels (Ziska *et al.*, 2003). *A. retroflexus* is established well under lack of soil moisture with good root system and may take advantages of elevated CO₂ for increasing the dry matter production of Redroot pigweed (Weller *et al.*, 2021). Redroot pigweed was grown with very well germination, growth, and CO₂ exchange rates at high temperatures, whereas the common lambsquarters grown well under low temperature (Chu, *et al.*, 1978; Egley, 1989).

CO₂ with Temperature

The competitiveness could be enhanced in C3 crop (Rice) relative to a C4 weed (*Echinochloa glabrescens*) with elevated CO₂ alone but simultaneous increases in CO₂ and temperature still favour C4 weed *spp.*

Comment [HUR14]: There is no reference to this table in the text.

Comment [HUR15]: The acronym for Versus is vs.

The competition outcomes between rice and a C4 weed- *Echinochloa glabrescens* L. using replacement series mixtures at two different CO₂ concentrations (393 and 594 µL L⁻¹) under day/night temperatures of 27/21°C and 37/29°C. Increasing the CO₂ concentration, at 27/21°C, resulted in a significant increase in aboveground biomass (+47%) and seed yield (+55%) of rice when averaged over all mixtures. For the C4 weed, higher CO₂ concentration did not produce a significant effect on biomass or yield.

When grown in mixture, the proportion of rice biomass increased significantly relative to that of the C4 weed in all mixtures at elevated CO₂ indicating increased 'competitiveness' of rice. However, under elevated CO₂ level and the higher temperature regime, competitiveness and reproductive stimulation of rice was reduced compared to the lower growth temperature, suggesting that while a C3 crop like rice may compete better against a C4 weed at elevated CO₂ alone (Alberto *et al.*, 1996).

CO₂ VS Leaf area and Biomass

Elevated CO₂

- In C3 weeds, leaf area generally responds less than biomass to CO₂ enrichment.
- However, in C4 weeds, leaf area and biomass more responses to CO₂ doubling (Fuhrer, 2003).

Elevated CO₂

- C3 grass species - Increase in tillering and C4 grass species - Increase in leaf area. Increases in CO₂ and temperature could still favour a C4 species (Yin and Struik 2008).

Table 6 : Effects of doubling CO₂ concentration on biomass and leaf area of C3 weeds (Patterson, 1995)

Range of response (% of growth at ambient)				
Scientific name	Common name	Biomass	Leaf area	No. of reports
<i>Abutilon theophrasti Medicus</i>	Velvet leaf	100-152	87-117	6
<i>Cassia obtusifolia L.</i>	Sicklepod	138-160	104-134	2
<i>Chenopodium album L</i>	lambquarters	100-155	122	2
<i>Cirsium arvense (L.) Scop</i>	Canada thistle	121	92	1
<i>Crotalaria spectabilis Roth</i>	Showy crotalaria	167	154	1

Comment [HUR16]: The acronym for Versus is vs.

Comment [HUR17]: There is no reference to this table in the text.

<i>Datura stramonium L.</i>	Jimson weed	174-272	146	1
<i>Elytrigia repens (L.) Neveski</i>	Quack grass	164	130	1
<i>Lolium perene L.</i>	Perennial rye grass	134-143	-	2
<i>Phalaris aquatica L.</i>	Harding grass	143	131	1
<i>Plantago lanceolata L.</i>	Buckhorn plantain	100-133	133	2
<i>Plantago major L.</i>	Broad leaf plantain	155	-	1
<i>Poa annua L.</i>	Annual blue grass	100	-	1
<i>Poa trivialis L.</i>	Rough stalk blue grass	103	-	1
<i>Rumex acetosella L.</i>	Red sorrel	131	-	1
<i>Rumex crispus L.</i>	Curly dock	118	96	1

Table 7 : Effects of doubling CO₂ concentration on biomass and leaf area of C4 weeds (Patterson, 1995)

Range of response (% of growth at ambient)				
Scientific name	Common name	Biomass	Leaf area	No. of reports
<i>Amaranthus retroflexus L.</i>	Pigweed, red rot	96-141	94-125	4
<i>Andropogon virginicus L.</i>	Broom sedge	81-117	88-129	2
<i>Cyperus rotundus L.</i>	Nut sedge, purple	102	92	1
<i>Digitaria ciliaris (Retz.)</i>	Crab grass	106-161	104-166	2
<i>Echinochloa crusgalli (L.) Beav.</i>	Barnyard grass	95-159	95-177	3
<i>Eleusine indica (L.) Gaertn.</i>	Goose grass	102-121	95-132	3
<i>Rottboellia cochinchinensis</i>	Itch grass	121	113	1
<i>Setaria faberi</i>	Foxtail, giant	93-135	101-140	3
<i>Sorghum halepense (L.)</i>	Johnson grass	56-110	99-103	3

Comment [HUR18]: There is no reference to this table in the text.

CO₂ VS Weedy rice

Ziska *et al.*, (2010) & Ziska *et al.*, (2013) found that in case of rice, rice biomass increased with increase in CO₂ from 300 to 400 ppm but did not increase further with increase in CO₂ to 500 ppm, whereas rice yield did not respond to elevated CO₂. Red rice responded linearly in terms of biomass as well as seed production. The elevated CO₂ (700 ± 50 μmol mol⁻¹) had increased the weedy rice germination, growth and biomass (Balbinot *et al.*, 2022)

Comment [HUR19]: The acronym for Versus is vs.

CO₂ VS Cocklebur

Ziska (2001) reported the effects of elevated CO₂ on the growth of grain sorghum (C₄) with and without common cocklebur (*X. strumarium*). At elevated CO₂, biomass and leaf area were higher for common cocklebur and significantly lower for sorghum in competitive mixtures, indicating that growth of C₄ crops could be reduced in the presence of C₃ weeds as future CO₂ levels increase (Kato and Esashi, 1975; Esashi, *et al.*, 1988). The elevated CO₂ enhances the imbibed cocklebur seeds (Esashi and Kato, 1975; Esashi *et al.*, 1978, Ishizawa *et al.*, 1988).

Comment [HUR20]: The acronym for Versus is vs.

CO₂ VS Ragweed and Quack grass

Increased CO₂ stimulated ragweed (*Ambrosia artemisiifolia*) pollen production several times more than that it stimulates overall growth (Singer *et al.* 2005). Tremmel and Patterson (1993) studied that increased CO₂ ameliorated the high temperature effects on quack grass (*Elytrigia repens*) and increase the growth and reproduction. C₃ weeds like lambsquarters or quackgrass are well grown in an elevated CO₂ environment (Tremmel and Patterson, 1994; Ziska *et al.* 1999; Ziska & Teasdale 2000).

Comment [HUR21]: The acronym for Versus is vs.

CO₂ VS Crops and weeds

Increasing CO₂ levels can favour either the crop or the weed in the same field. Rising atmospheric CO₂ concentration altered the competition between rice and barnyard grass in favour of rice (Zeng *et al.*, 2011). Nitrogen was found to be the limiting factor for biomass increase in rice at elevated CO₂ levels compared with barnyard grass (Zhu *et al.*, 2008).

Comment [HUR22]: The acronym for Versus is vs.

CO₂ VS Rice and Wheat

The effect of CO₂ enrichment on weed species at the Directorate of Weed Science, Jabalpur revealed that a few weed species such as *Dactyloctenium aegyptium* and *E. colona* responded to elevated CO₂, but *Cyperus rotundus* and *Eleusine indica* did not respond to CO₂ enrichment. Weedy rice (*Oryza sativa* L.) responds more strongly than cultivated rice to rising CO₂ level with greater competitive ability, suggesting that weedy rice may become a more problematic weed in the future. The growth and biomass accumulation was increased of the C₄ weed (*Amaranthus viridis*) under elevated CO₂ (Naidu and Paroha, 2008).

Comment [HUR23]: The acronym for Versus is vs.

The C3 crop like rice, elevated CO₂ levels may have positive effects on crop competitiveness with C4 weeds. The problems of *P. minor* and *Artemisia ludoviciana* in wheat would aggravate with increase in CO₂ due to climate change. Further, the problem could be aggravated with water scarcity. Due to CO₂ enrichment, the wheat plant could gain biomass against *P. minor*. Under water stress conditions, *P. minor* had advantage over wheat with CO₂ enrichment (Fuhrer, 2003).

CO₂ VS Wild oat

The CO₂ enrichment/elevated CO₂ hastened the seed maturity in *Avena fatua* (wild oat) in wheat. The seeds were matured 13 days in advance compared to the plants grown under ambient CO₂ conditions and enrichment of soil seed bank as the wild oat seeds shatter well before the harvest of the crop. The wild oat plants grown at 480 ppm CO₂ produced 44% more seed than 357 ppm (O'Donnell and Adkins, 2001; Johannessen et al., 2005; Granger et al., 2012; Momtazi and Miri, 2015; Ziska, et al., 2015; Oraki et al., 2016; Ziska, 2017; Balbinot, 2020)

Comment [HUR24]: The acronym for Versus is vs.

CO₂ VS Noxious or Invasive weeds

The elevated CO₂ increases the growth and development of Noxious weeds (Dukes, 2000; Polley et al., 2003; Weltzin et al., 2003; Salo, 2005). *Cynodon dactylon* in rice and *Convolvulus arvensis* in wheat. These weeds may show a strong response in growth with increase in atmospheric CO₂ (Ziska et al., 2004). In India, *Parthenium hysterophorus* had shown tremendous growth response to elevated CO₂. Increased flower production of *Parthenium hysterophorus* (C3) and *Amaranthus viridis* (C4) under elevated CO₂ (Naidu, 2013; Waryszak et al., 2018). The elevated CO₂ recorded in significant changes in dry matter production and morphology of invasive *M. micrantha* and *W. trilobata* (Song et al., 2009). Many invasive plants have been shown to respond positively to elevated (CO₂) when grown individually or in monoculture (Weltzin et al., 2003). Examples of these are species that have invaded North America, such as cheatgrass (*Bromus tectorum*), kudzu (*Pueraria lobata*) and Japanese honeysuckle (*Lonicera japonica*, which has also invaded New Zealand and parts of Europe) (Ziska, 2003).

Comment [HUR25]: The acronym for Versus is vs.

CO₂ VS Worst weed

Many growers recognize that the worst weeds for a given crop are similar in growth habit. There are 14 of the world's worst weeds comes under C4 plants. Eg: Oat - wild oat,

Comment [HUR26]: The acronym for Versus is vs.

Wheat - little seed canary grass, Rice -wild rice growth is more favoured due to increased CO₂ level in the atmosphere (Patterson, 1995; Ziska, 2003).

Weed Response to Increasing Temperature

Climate change involves rising temperatures (Tubiello *et al.* 2007). Temperature is a primary factors influencing the distribution of weeds. Increasing temperatures on weeds shift into higher latitudes or higher altitudes. Under high temperature, plants with C4 carbon fixation pathway (mostly weeds) have a competitive advantage over crop plants. Increased atmospheric CO₂ levels are likely to be accompanied by high temperature favouring C4 weeds over C3 crops. A similar shift in weed species composition can also be expected under erratic rainfall because of climate change. Due to sudden change in climate, environmental stress on a crop may increase and as a result the crop could become more vulnerable to attack by insect and pathogens, and less competitive with weeds.

The aberrations in weather conditions not only affect crop-weed competition, but also trigger weed seed germination in several flushes causing serious weed management issues. Three flushes of *P. minor* are not uncommon in the wheat fields in northwest India, which are not controlled by a single application of herbicide. Weeds are more responsive to increasing temperature because of their wider gene pool compared with crops, which enables them to adapt to diverse environment conditions. Normally the mean temperatures increases over a long period, weeds will expand their range into new areas (Tubiello *et al.* 2007).


Warmer seasonal temperatures and milder winters will extend the distribution of invasive weeds. (Kudzu and Ragweed) (Garrett, 2012). Temperature rises will give competitive advantage to C4 plants than to C3 plants. Hanzlik and Gerowitt (2012) studied that Soybean was more adversely affected by rising temperature than prickly sida (*Sida spinosa*) and sicklepod (*Senna obtusifolia*).

Temperature VS Weed Phenology

Climate change may bring changes in weed population and in their phenology. The increased temperature had a greater effect on plant Phenological development than elevated CO₂. The increasing temperature by 4°C had advanced the emergence timing of *C. album* and *S. viridis* by 26 and 35 days, and flowering time by 50 and 31.5 days. Increased temperatures strongly affected the biomass accumulation by annual grass species during their reproductive phase as compared with the vegetative phase (Lee, 2011).

Comment [HUR27]: The acronym for Versus is vs.

UNDER PEER REVIEW



INDIA

Sing *et al.* (1991) reported that most of the weeds in rice are of C4 type. *Ischaemum rugosum* more in Tropical areas. *Rumex spinosus* in wheat of North West India has increased.

AUSTRALIA

The Buffel grass (*Cenchrus ciliaris*) is one of the few weeds in Australia to be extensively assessed for growth response to climate change (growth at 35°C versus 25°C).

The warm temperature increases many C4 weeds, such as *Amaranthus retroflexus* L., *Setaria* sp., *Digitaria* sp., *Sorghum halepense* L., *Paspalum dichotomiflorum* (L.) are expected to expand further north side of Australia (Clements and DiTommaso, 2011). *Hieracium aurantiacum* L. (Brinkley and Bomford, 2002).

CALIFORNIA

Temperature can allow sleeper weeds to become invasive. Very aggressive weeds that are currently found in the lower latitudes are limited in the higher latitudes of California. The Itch grass has profusely tillering at increasing temperature (Bunce and Ziska 2000). The Robust grass weed could have been invaded the central Midwest and California with a 3°C warming trend (Patterson, 1995).

European

Due to warmer the three important Central European C4 weeds in maize like, *Amaranthus retroflexus*, *Echinochloa crusgalli* and *Setaria viridis*. Most of these weeds are late germinators and emerge from early summer to early autumn (Walther *et al.*, 2002). O'Donnell and Adkins (2001) reported that wild oat plants grown at high temperature 23/19°C (day/night) completed their development faster than those grown at normal temperature 20/16°C.

Temperature VS *Striga*

Atmospheric temperature is regarded as an important indicator of weed species distribution in a geographical area. *Striga* species are highly diverse. *Striga asiatica* is relatively insensitive to temperature. Its distribution may be more affected by changes in the geographical range of the host crop than directly by temperature (Mohamed *et al.*, 2007).

Comment [HUR28]: The acronym for Versus is vs.

Temperature VS *Parthenium*

Comment [HUR29]: The acronym for Versus is vs.

A higher percentage of *Parthenium* seed germination was observed during summer. Nguyen et al. (2010) have reported that the highest number of *Parthenium* seeds were produced under warm conditions, with a maximum number (60%) of unfilled seeds under cool condition. Warm dry conditions allow reproduction after only 50 days of growth, while cool and wet conditions delay this to after 75 days. Increased temperature enhances growth of *Parthenium*, enlarges canopy size and structure and accelerates population growth rate because of the shortened life cycle (Singh 2011; Toh et al., 2011).

Weed Response to Precipitation

Lantana camara, for example, could expand if rainfall increased in some areas (McFadyen 2008). The drought-tolerant C4 weeds, parasitic weeds that thrive in erratic and low rainfall environments (e.g. *Striga hermonthica*) or temporarily flooded conditions (e.g. *Rhamphicarpa fistulosa*) could benefit. *Striga* spp. problems are also associated with low soil fertility area (Kroschel, 2000).

Competition of cotton with *Abutilon theophrasti* and *Anoda cristata* (L.) increased under drought conditions (Patterson and Highsmith, 1989). A yield reduction due to *Xanthium strumarium* L. was more pronounced in well-watered soybeans compared with water-stressed soybeans (Mortensen and Coble, 1989). An increase in rainfall provided greater competition to wheat growth and yield against *Cirsium arvense* (Donald and Khan, 1992). *Amaranthus spinosus* and *Leptochloa chinensis* (L) survived under water stress conditions and produced a significant number of tillers/branches.

Cheat grass and Yellow star thistle (*Centaurea solstitialis*) are annual weeds survive under drought conditions (Vollmer and Vollmer, 2006). While cheat grass completes its life cycle based on available moisture and yellow star thistle competes with native plants by developing deeper root system (Hatfield et al., 2011). Drought and dry soil conditions prolong the weed seed bank longevity.

Flooding

Submergence protects rice plants from severe competition with C4 weeds. On the other hand upland rice and rainfed lowland rice with limited precipitation face severe competition with C4 weeds. Alternate wetting and drying in puddled as well as dry-seeded

rice may encourage weeds such as *Livistona chinensis*, *Eleusine indica* and *Eleusine prostrata*. Tamarisk (*Tamarix aphylla*) Willows (*Salix spp.*) and *Rhamphicarpa fistulosa* are benefit (increase growth) to flooding events (Rodenburg *et al.*, 2010).

Wind

The increasing CO₂ levels may also increase wind dispersal of weed seeds by either increasing the height of the weed plant or by increasing the plant size. Wind dispersed invasive weed species are *Cirsium arvense*, *Sonchus arvensis* L., *Sonchus oleraceus* L., and *Carduus nutans* L (Ziska and Dukes 2011).

Fire

Gamba grass (*Andropogon gayanus*) is reason for fire under extreme weather condition. Fire is a common extreme event in Australia and its frequency and severity is often influenced by extreme weather.

Climate Change: Crop-Weed Interactions

Crop-weed interactions vary significantly in various climatic regions, depending on temperature, precipitation, soil, etc.

- A. C3 crop competing with both C3 and C4 weeds where C3 weeds are dominant, whereas in another situation C4 weeds are dominant.
- B. C4 crop competing with both C3 and C4 weeds where C3 weeds are dominant, whereas in another situation C4 weeds are dominant.

Table 8: Crop/weed competition outcome at elevated CO₂ conditions (Bunce and Ziska 2000)

Weed species	Crop	Favoured under elevated CO ₂	References
<i>Amaranthus retroflexus</i> (C4)	Soybean (C3)	Crop	(Ziska, 2000)
<i>Amaranthus retroflexus</i> (C4)	Sorghum (C4)	Weed	(Ziska, 2003)
<i>Chenopodium album</i> (C3)	Soybean (C3)	Weed	(Ziska, 2000)
<i>Taraxacum officinale</i> (C3)	Lucern (C3)	Weed	(Bunce, 1995)
<i>Albutilon theophrasti</i> (C3)	Sorghum (C4)	Weed	(Ziska, 2003)
<i>Taraxacum and Plantago</i> (C3)	Grasses (C3)	Weed	(Potvin and Vasseur, 1997)

Comment [HUR30]: There is no reference to this table in the text.

Red rice (C3)	Rice (C3)	Weed	(Ziska <i>et al.</i> , 2010)
<i>Echinochloa glabrescens</i> (C4)	Rice (C3)	Weed	(Alberto <i>et al.</i> 1996)

Climate Change on Weed Management

Herbicide Efficacy

I. Carbon Dioxide and Temperature

Reduced stomatal conductance can alter the efficacy of both foliar- and soil-applied herbicides. Change in temperature and CO₂ concentration, may alter transpiration, the number of leaf stomata, or the thickness of the leaf, which may affect the absorption and translocation of herbicides.

II. Relative Humidity

Relative humidity could influence the efficacy of foliar-applied herbicides through interactions between the herbicide droplet, leaf cuticle, and availability of water (Devine *et al.*, 1993).

III. Wind

Wind may have a less pronounced influence on herbicide performance. Nonetheless, windy conditions can interfere with surface application and cause spray drift, thereby reducing spray application efficiency

IV. Rainfall

Increased rainfall frequency and intensity will have an adverse effect on uptake, retention, and activity of soil-applied herbicides

Weed management under climate change

- Continue current control options
- Integrated Weed Management (IWM)
- Use control options suitable for current extreme events.
- Develop new control methods adapted for climate change.

Summary

Changes in temperature and carbon dioxide are likely to have significant direct (CO₂ stimulation of weed growth) and indirect effects (climatic variability) on weed biology. In spite of the importance of weed biology in both the environment and in farms, very little is known regarding the impact of these environmental changes on both the reproductive success of agronomic or invasive weeds, and the potential consequences for their management. The growth and reproduction of C₃ weeds will increase under changing climatic conditions.

Conclusion

Global climate change factors have serious implications for not only crop and weed growth and also herbicide performance and the effectiveness of chemical weed management. Weeds tend to show better survival mechanisms under changing climate because of their greater inter-specific genetic variation and physiological plasticity.

Reference

- Alberto, A.M.P., L.H. Ziska, C.R. Cervancia, and P.A. Manalo. 1996. The Influence of Increasing Carbon Dioxide and Temperature on Competitive Interactions Between a C₃ Crop, Rice (*Oryza sativa*) and a C₄ Weed (*Echinochloa glabrescens*). *Australian Journal of Plant Physiology*, 23, 795-802.
- Anwar, M. P., Islam, A. M., Yeasmin, S., Rashid, M. H., Juraimi, A. S., Ahmed, S., & Shrestha, A. (2021). Weeds and Their Responses to Management Efforts in a Changing Climate. *Agronomy*, 11(10), 19-21.
- Balbinot, A. (2020). *Effect of enhanced CO₂ atmospheric concentration on rice and weedy rice* (Master's thesis, Universidade Federal de Pelotas).
- Balbinot, A., da Rosa Feijó, A., Fipke, M. V., Gehrke, V. R., Agostinetto, D., Kruse, N. D., and de Avila, L.A. (2022). Rising atmospheric CO₂ concentration affect weedy rice growth, seed shattering and seedbank longevity. *Weed Research*.
- Batts G.R, Morison JIL, Ellis RH et al (1997) Effects of CO₂ and temperature on growth and yield of crops of winter wheat over four seasons. *Eur J Agron* 7:43-52.
- Bergmann J, Pompe S, Ohlemüller R (2010) The Iberian Peninsula as a potential source for the plant species pool in Germany under projected climate change. *Plant Ecol* 207:191–201.
- Bradley, B.A., D.M. Blumenthal, D.S. Wilcove, and L.H. Ziska. 2010. Predicting plant invasions in an era of global change. *Trends in Ecology and Evolution*, 25, 310-318.
- Brinkley, T. R., and M. Bomford. 2002. Agricultural Sleeper Weeds: What is the Potential Threat? Kingston, ACT: *Bureau of Rural Sciences*.

- Bunce, J. A., and L.H. Ziska, 2000. "Crop ecosystem responses to climatic change: crop/weed interactions," in *Climate Change and Global Crop Productivity*, eds K. R. Reddy and H. F. Hodges (New York, NY: CABI), 333–348.
- Carter DR, Peterson KM (1983) Effects of a CO₂-enriched atmosphere on the growth and competitive interaction of a C3 and a C4 grass. *Oecologia* 58:188-193.
- Chandrasena, N. (2008). Weeds and weed management under a changing climate. In *Conference: Biennial Conference of Indian Weed Science Society*.
- Chandrasena, N. (2009). How will weed management change under climate change? Some perspectives. *Journal of Crop and Weed*, 5(2), 95-105.
- Chu, C.C., Ludford, P.M., Ozbun, J.L., & Sweet, R.D. (1978). Effects of Temperature and Competition on the Establishment and Growth of Redroot Pigweed and Common Lambsquarters 1. *Crop Science*, 18(2), 308-310.
- Clements, D. and A. Ditommaso. 2011. Climate change and weed adaptation: Can evolution of invasive plants lead to greater range expansion than forecasted? *Weed Res.*, 51, 227-240.
- Dass, A., Shekhawat, K., Choudhary, A.K., Sepat, S., Rathore, S.S., Mahajan, G. and Chouhan, B.S. 2017. Weed management in rice using crop competition- a review. *Crop Protection* 95: 45-52.
- Devine, M.D., S.O. Duke, and C. Fedtke. 1993. *Foliar Absorption of Herbicides*. Prentice-Hall, Englewood Cliffs, NJ. 29–52.
- Donald, W.W., and M. Khan, 1992. Yield loss assessment for spring wheat (*Triticum aestivum*) infested with Canada thistle (*Cirsium arvense*). *Weed Sci.*, 40, 590–598.
- Dukes JS, Pontius J, Orwig D (2009) Responses of insect pests, pathogens, and invasive plant species to climate change in the forests of north eastern North America: what can we predict? *Can J For Res.*, 39:231–248.
- Dukes, J.S., 2000. Will the increasing atmospheric CO₂ concentration affect the success of invasive species? In: Mooney, H.A., Hobbs, R.J. (Eds.), *Invasive Species in a Changing World*. Island Press, Washington.
- Dukes, J.S., J. Pontius, D. Orwig. 2009. Responses of insect pests, pathogens, and invasive plant species to climate change in the forests of northeastern North America: what can we predict? *Can J For Res.*, 39, 231–248.
- Egley, G. H. (1989). Some effects of nitrate-treated soil upon the sensitivity of buried redroot pigweed (*Amaranthus retroflexus* L.) seeds to ethylene, temperature, light and carbon dioxide. *Plant, Cell & Environment*, 12(5), 581-588.
- Elmore, C.D., & Paul, R.N. (1983). Composite list of C4 weeds. *Weed Science*, 31(5), 686-692.

- Esashi, Y., & Katoh, H. (1975). Dormancy and impotency of cocklebur seeds III. CO₂-and C₂H₄-dependent growth of the embryonic axis and cotyledon segments. *Plant and cell physiology*, 16(4), 707-718.
- Esashi, Y., Kawabe, K., Isuzugawa, K., & Ishizawa, K. (1988). Interrelations between carbon dioxide and ethylene on the stimulation of cocklebur seed germination. *Plant Physiology*, 86(1), 39-43.
- Esashi, Y., Okazaki, M., Yanai, N., & Hishinuma, K. (1978). Control of the germination of secondary dormant cocklebur seeds by various germination stimulants. *Plant and Cell Physiology*, 19(8), 1497-1506.
- Fuhrer, J. (2003). Agro-ecosystem responses to combination of elevated CO₂, ozone and global climate change. *Agric. Ecosys. Environ.*, 97, 1–20.
- Garrett, K. (2012). *Climate Change Impacts on Insect Pests, Weeds, and Disease (PowerPoint)* (No. 1456-2016-120261).
- Ghannoum O, von Caemmerer S, Ziska LH, Conroy JP (2000). The growth response of C4 plants to rising atmospheric CO₂ partial pressure: a reassessment. *Plant, Cell and Environment*, in press.
- Granger, K. L. (2012). *Mitigation of drought-induced reductions in wild oat seed quality by elevated carbon dioxide*. The Pennsylvania State University.
- Hanzlik, K., and B. Gerowitt. 2012. Occurrence and distribution of important weed species in German winter oilseed rape fields. *J. Plant Dis. Prot.*, 119, 107–120.
- Hatfield, J.L., K.J. Boote, B.A. Kimball, L.H. Ziska, R.C. Izaurralde, D. Ort, A.M. Thomson, and D. Wolfe. 2011. Climate impacts on agriculture: implications for crop production. *Agron. J.*, 103, 351–370.
- Houghton, J. (2001). The science of global warming. *Interdiscip. Sci. Rev.* 26, 247–257.
- IPCC, 2001. The Intergovernmental Panel on Climate Change. <http://www.ipcc.ch/>
- IPCC, 2004. Intergovernmental Panel on Climate Change, IPCC Fourth Assessment Report: Climate Change 2007 (AR4). <http://www.ipcc.ch>
- Ishizawa, K., Hoshina, M., Kawabe, K., & Esashi, Y. (1988). Effects of 2, 5-norbornadiene on cocklebur seed germination and rice coleoptile elongation in response to CO₂ and C₂H₄. *Journal of Plant Growth Regulation*, 7(1), 45-58.
- Jinger, D., Dass, A., Kumar, V., Kaur, R., & Kumari, K. (2016). Weed Management Strategies Weed Management Strategies Weed Management Strategies Weed Management Strategies. *Indian Farming*, 66(9), 09-13.
- Johannessen, M. M., Mikkelsen, T. N., Nersting, L. G., Gullord, M., Von Bothmer, R., & Jørgensen, R. B. (2005). Effects of increased atmospheric CO₂ on varieties of oat. *Plant Breeding*, 124(3), 253-256.
- Katoh, H., & Esashi, Y. (1975). Dormancy and impotency of cocklebur seeds I. CO₂, C₂H₄ O₂ and high temperature. *Plant and Cell Physiology*, 16(4), 687-696.

- Kendal, A.C., J.C. Turner, and S.M. Thomas. 1985. "Effects of CO₂ enrichment at different irradiances on growth and yield of wheat", I. Effects of cultivar and duration of CO₂ enrichment. *Journal of Experimental Botany*, 36, 252-260.
- Kroschel J. 2000. *Striga* - How will it affect African agriculture in the future? — An ecological perspective. 137–158.
- Lee, J. 2011. Combined effect of elevated CO₂ and temperature on the growth and phenology of two annual C3 and C4 weedy species. *Agric. Ecosys. Environ.*, 140, 484–491.
- Leegood, R.C. (2002). C4 photosynthesis: principles of CO₂ concentration and prospects for its introduction into C3 plants. *J. Exp. Bot.* 53, 581–590
- Lobell DB, Burke MB (2008) Why are agricultural impacts of climate change so uncertain? The importance of temperature relative to precipitation. *Environ Res Lett* 3:1–8.
- Loss SR, Terwilliger LA, Peterson AC (2011) Assisted colonization: integrating conservation strategies in the face of climate change. *Biol Conserv* 144:92–100
- Mahajan, G., S.Singh and B.S. Chauhan. 2012. Impact of climate change on weeds in the rice–wheat cropping system. *Current Science*, 102, 1254-1255.
- McFadyen, R. 2008. *Invasive Plants and Climate Change. Weeds* CRC Briefing Notes. CRC for Australian Weed Management.
- McPeck, T.M., & Wang, X. (2007). Reproduction of dandelion (*Taraxacum officinale*) in a higher CO₂ environment. *Weed Science*, 55(4), 334-340.
- Mishra, G. N. 2003. Weeds and their management in upland rice. *Ind. Farming*. 18-21.
- Mohamed, K.I., J.F. Bolin, L.J. Musselman, and A. Townsend Peterson. 2007. Genetic diversity of *Striga* and implications for control and modelling future distributions. 71–84. In: Ejeta G and Gressel J eds. *Integrating New Technologies for Striga Control - Towards ending the witch-hunt*. World Scientific, Singapore.
- Momtazi, F., & Miri, H. (2015). Growth and competitiveness of wheat against C3 and C4 plant in CO₂ enrichment and drought condition. *Journal of Plant Ecophysiology*, 7(20), 112-127.
- Morgan, J.A., D.R. LeCain, A.R. Mosier, and D.G. Milchunas. 2001. Elevated CO₂ enhances water relations and productivity and affects gas exchange in C3 and C4 grasses of the Colorado short grass steppe. *Global Change Biol.*, 7, 451–466.
- Morison JIL and Lawlor DW (1999) Interactions between increasing CO₂ concentration and temperature on plant growth. *Plant Cell Environ* 22: 659-682.
- Mortensen, D.A., and H.D. Coble. 1989. The influence of soil water content on common cocklebur (*Xanthium strumarium*) interference in soybeans (*Glycine max*). *Weed Sci.*, 37, 76–83.
- Naidu V.S. and S. Paroha. 2008. Growth and biomass partitioning in two weed species, *Parthenium hysterophorus* (C3) and *Amaranthus viridis* (C4) under elevated CO₂. *Ecology Environment and Conservation*. 14, 9-12.

- Naidu, V.S.G.R., 2013. Invasive potential of C3-C4 intermediate *Parthenium hysterophorus* under elevated CO₂. *Indian Journal of Agricultural sciences*. 83, 176-179.
- Nguyen, T.L.T., S.C. Navie, and S.W. Adkins. 2010. The reproductive capacity of *Parthenium* weed under different climatic conditions. In Proceedings 17th Australasian Weeds Conference, Published by New Zealand Plant Protection Society (Ed. S. M. Zydenbos). 26 - 30th September. 124-127
- O'Donnell, C.C. and S.W. Adkins. 2001. Wild oat and climate change: the effect of CO₂ concentration, temperature, and water deficit on the growth and development of wild oat in monoculture. *Weed Sci.*, 49, 694–702.
- O'Donnell, C.C., & Adkins, S.W. (2001). Wild oat and climate change: The effect of CO₂ concentration, temperature, and water deficit on the growth and development of wild oat in monoculture. *Weed Science*, 49(5), 694-702.
- Oraki, H., Modares-Sanavi, S. A. M., Zand, E., & Mokhtassi-Bidgoli, A. (2016). Effect of carbon dioxide enrichment on yield and yield components of wheat under weed competitive pressure of winter wild oat (*Avena ludoviciana* L.). *Iranian Journal of Field Crop Science*, 47(2).
- Patterson , D.T. 1985. Comparative ecophysiology of weeds. In : *Weed Physiology : Reproduction and Ecophysiology*, S. O. Duke (ed.). CRC Press, Inc. Boca Raton, Florida. pp. 101-132.
- Patterson, D.T. (1995). Effects of environmental stress on weed/crop interactions. *Weed Science*, 43(3), 483-490.
- Patterson, D.T. 1995. Weeds in a changing climate. *Weed Science*, 43, 685–701.
- Patterson, D.T., and M.T. Highsmith. 1989. Competition of spurred anoda (*Anoda cristata*) and velvet leaf (*Albutilon theophrasti*) with cotton (*Gossypium hirsutum*) during simulated drought and recovery. *Weed Sci.*, 37, 658–664.
- Pautasso M, Dehnen-Schmutz K, Holdenrieder O et al (2010) Plant health and global change—some implications for landscape management. *Biol Rev* 85:729–755.
- Petit JR, Jouzel J, Raynaud D et al (1999) Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica. *Nature* 399:429–436.
- Polley, H.W., Johnson, H.B., Tischler, C.R., 2003. Woody invasion of grasslands: evidence that CO₂ enrichment indirectly promotes establishment of *Prosopis glandulosa*. *Plant Ecol* 164, 85-94
- Poorter, H., 1993. "Interspecific variation in the growth response of plants to an elevated ambient CO₂ concentration", *Vegetation*, 104/105, 77–97.
- Ramesh, K., Matloob, A., Aslam, F., Florentine, S. K., & Chauhan, B. S. (2017). Weeds in a changing climate: vulnerabilities, consequences, and implications for future weed management. *Frontiers in plant science*, 8, 95.

- Reddy, A.R., Rasineni, G.K., & Raghavendra, A.S. (2010). The impact of global elevated CO₂ concentration on photosynthesis and plant productivity. *Current Science*, 46-57.
- Robinson TMP, Gross KL (2010) The impact of altered precipitation variability on annual weed species. *Am J Bot* 97:1625–1629.
- Robinson TMP, Gross KL (2010) The impact of altered precipitation variability on annual weed species. *Am J Bot* 97:1625–1629.
- Rodenburg, J., C.R. Riches, and J.M. Kayeke. 2010. Addressing current and future problems of parasitic weeds in rice. *Crop Prot.*, 29, 210–221.
- Rodenburg, J., H. Meinke, and D.E. Johnson. 2011. Challenges for weed management in African rice systems in a changing climate. *J. Agric. Sci.*, 149, 427–435.
- Rogers HH, Runion GB, Prior SA et al (2008) Effects of elevated atmospheric CO₂ on invasive plants: comparison of purple and yellow nutsedge (*Cyperus rotundus* L. and *C. esculentus* L.). *J Environ Qual* 37:395–400.
- Salo, L.F., 2005. Red brome (*Bromus rubens* subsp. *madritensis*) in North America: possible modes for early introductions, subsequent spread. *Biol. Invasions* 7, 165–180
- Singer A, Travis JMJ, Johst K (2013) Interspecific interactions affect species and community responses to climate shifts. *Oikos* 122:358–366.
- Singer, A., J.M.J. Travis, K. Johst. 2013. Interspecific interactions affect species and community responses to climate shifts. *Oikos*, 122, 358–366.
- Singer, B.D., L.H. Ziska, D.A. Frenz, D.E. Gebhard, and J.G. Straka. 2005. Increasing Amb a1 content in common ragweed (*Ambrosia artemisiifolia*) pollen as a function of rising atmospheric CO₂ concentration. *Functional Plant Biology*, 32: 667-670.
- Singh T., J.S Kolar, and R.K. Bhatia. 1991. Emergence behaviours of *Ischaemum rugosum* (Wrinkle Grass) as affected by variable temperature, submergence period and depth of seed placement in the soil. *Indian Journal of Weed Science*, 23: 56–60
- Singh, R.P., Singh, R.K., & Singh, M.K. (2011). Impact of climate and carbon dioxide change on weeds and their management-a review. *Indian Journal of Weed Science*, 43(1&2), 1-11.
- Skov F, Svenning J.C. (2004) Potential impact of climatic change on the distribution of forest herbs in Europe. *Ecography (Cop)* 27:366– 380.
- Song, L., Wu, J., Li, C., Li, F., Peng, S., & Chen, B. (2009). Different responses of invasive and native species to elevated CO₂ concentration. *Acta Oecologica*, 35(1), 128-135.
- Tang, J., Xu, L., Chen, X. and Hu, S. 2009. Interaction between C₄ barnyard grass and C₃ upland rice under elevated CO₂ : Impact of mycorrhizae. *Acta Oecologica*, 35: 227-235.
- Toh R, J. Hanan, K. Dhileepan, R.G. Shivas and S.W. Adkins. 2011. Stimulation of *Parthenium* weed canopy under changing climate using L - systems. 19th International Congress on Modelling & Stimulation, Perth, Australia. 12-16th December. 1009-1015

- Treharne, K., 1989. The implications of the 'greenhouse effect' for fertilizers and agrochemicals. In: Benner, R.D. (Ed.), *The Greenhouse Effect and UK Agriculture*. Ministry of Agriculture, Fisheries and Food, London, pp. 67–78.
- Tremmel, D.C., & Patterson, D.T. (1994). Effects of elevated CO₂ and temperature on development in soybean and five weeds. *Canadian journal of plant science*, 74(1), 43–50.
- Tremmel, D.C., and D.T. Patterson. 1993. Responses of soybean and five weeds to CO₂ enrichment under two temperature regimes. *Can. J. Plant Sci.*, 73, 1249–1260.
- Tubiello, F.N., J.F. Soussana, and S.M. Howden. 2007. Crop and pasture response to climate change. *Proc. Natl. Acad. Sci. U.S.A.* 104, 19686–19690.
- Vollmer, J.L., J.G. Vollmer. 2006. Controlling cheat grass in winter range to restore habitat and endemic fire. In: Kitchen, S.G., Pendleton, R.L., Monaco, T.A., Vernon, J. (Comps.). *Proc. Shrublands under fire: disturbance and recovery in a changing world*, 6–8.
- Walther, G.R., E. Post, P. Convey, A. Menzel, C. Parmesan, T.J.C. Beebee, J.M. Fromentin, O. Hoegh-guldberg, and F. Bairlein. 2002. Ecological responses to recent climate change. *Nature*. 416, 389–395.
- Waryszak, P., Lenz, T. I., Leishman, M. R., & Downey, P. O. (2018). Herbicide effectiveness in controlling invasive plants under elevated CO₂: sufficient evidence to rethink weeds management. *Journal of environmental management*, 226, 400–407.
- Weller, S., Florentine, S., Javaid, M. M., Welgama, A., Chadha, A., Chauhan, B. S., & Turville, C. (2021). *Amaranthus retroflexus* L.(Redroot Pigweed): Effects of elevated CO₂ and soil moisture on growth and biomass and the effect of radiant heat on seed germination. *Agronomy*, 11(4), 728.
- Weltzin, J.F., Belote, R.T., & Sanders, N.J. (2003). Biological invaders in a greenhouse world: will elevated CO₂ fuel plant invasions?. *Frontiers in Ecology and the Environment*, 1(3), 146–153.
- Weltzin, J.F., Belote, R.T., Sanders, N.J., 2003. Biological invaders in a greenhouse world: will elevated CO₂ fuel plant invasions? *Front. Ecol. Environ* 1, 146–153.
- Yin, X., and P.C. Struik. 2008. Applying modelling experiences from the past to shape crop. *New Phytologist*. 179: 629–642.
- Zangerl AR, Bazzaz FA (1984) The response of plants to elevated CO₂— II. Competitive interactions among annual plants under varying light and nutrients. *Oecologia* 62:412–417
- Zeng, Q., B. Liu, B. Gilna, Y. Zhang, C. Zhu, H. Ma, J. Pang, G. Chen, and J. Zhu. 2011. Elevated CO₂ effects on nutrient competition between a C₃ crop (*Oryza sativa* L.) and a C₄ weed (*Echinochloa crus-galli* L.) nutrient cycling. *Agroecosystems*. 89, 93–104.

- Zhu, C., Q. Zeng, L.H. Ziska, J. Zhu, Z. Xie, and G. Liu. 2008. Effect of nitrogen supply on carbon dioxide-induced changes in competition between rice and barnyard grass (*Echinochloa crusgalli*). *Weed Sci.*, 56, 66–71.
- Ziska LH, Teasdale JR (2000) Sustained growth, photosynthesis and increased tolerance to glyphosate observed in a C3 perennial weed, quackgrass, grown at elevated carbon dioxide. *Australian Journal of Plant Physiology*, 27, 157±164
- Ziska LH, Teasdale JR, Bunce JA (1999) Future atmospheric carbon dioxide concentrations may increase tolerance to glyphosate [N-(phosphonomethyl) glycine] in weedy species. *Weed Science*, 47, 608-615.
- Ziska, L. H. (2003). Evaluation of the growth response of six invasive species to past, present and future atmospheric carbon dioxide. *Journal of Experimental Botany*, 54 (381), 395-404.
- Ziska, L. H. and J.S. Dukes. 2011. *Weed Biology and Climate Change*. Ames, IA: Blackwell Publishing Ltd, 68-205.
- Ziska, L. H., Gealy, D. R., Burgos, N., Caicedo, A. L., Gressel, J., Lawton-Rauh, A. L., & Merotto Jr, A. (2015). Weedy (red) rice: an emerging constraint to global rice production. *Advances in agronomy*, 129, 181-228.
- Ziska, L. H., Tomecek, M. B., & Gealy, D. R. (2013). Assessment of cultivated and wild, weedy rice lines to concurrent changes in CO₂ concentration and air temperature: determining traits for enhanced seed yield with increasing atmospheric CO₂. *Functional Plant Biology*, 41(3), 236-243.
- Ziska, L.H. (2008). Rising atmospheric carbon dioxide and plant biology: the overlooked paradigm. *DNA and Cell Biology*, 27(4), 165-172.
- Ziska, L.H. (2016). The role of climate change and increasing atmospheric carbon dioxide on weed management: herbicide efficacy. *Agriculture, Ecosystems & Environment*, 231, 304-309.
- Ziska, L.H. (?). Could recent increases in atmospheric CO₂ have acted as a selection factor in *Avena fatua* populations? A case study of cultivated and wild oat competition. *Weed Research*, 57(6), 399-405.
- Ziska, L.H. 2000. The impact of elevated CO₂ on yield loss from a C3 and C4 weed in field-grown soybean. *Global Change Biology*, 6, 899-905.
- Ziska, L.H., D.E. Gebhard, D.A. Frenz, S. Faulkner, and B.D. Singer. 2003. Cities as harbingers of climate change: Common ragweed, urbanization, and public health. *Journal of Allergy and Clinical Immunology*, 111:290-295.
- Ziska, L.H., Faulkner, S., and Lydon, J. (2004). Changes in biomass and root: shoot ratio of field-grown Canada thistle (*Cirsium arvense*), a noxious, invasive weed, with elevated CO₂: implications for control with glyphosate. *Weed Science*, 52(4), 584-588.

- Ziska, L.H., M.B. Tomecek, and D.R. Gealy. 2010, "Evaluation of competitive ability between cultivated and red weedy rice as a function of recent and projected increases in atmospheric CO₂", *Agronomy Journal*, 102,118-123
- Ziska, L.W. and J. Bunce, 1993, "Inhibition of whole plant respiration by elevated CO₂ as modified by growth temperature", *Physiologia Plantarum*, 87, 459-466.
- Ziska, L.W., 2000, "The impact of elevated CO₂ on yield loss from a C3 and C4 weed in field-grown soybean", *Global Change Biology*, 6, 899-905.
- Ziska, L.W., 2001, "Changes in competitive ability between a C4 crop and a C3 weed with elevated carbon dioxide", *Weed Science*, 49, 622-627.
- Ziska, L.W., 2003, "Evaluation of yield losses in field sorghum from a C3 and C4 weeds with increasing CO₂", *Weed Science*, 51, 914-918.

UNDER PEER REVIEW