

Original Research Article

Typhoid Fever: Azithromycin vs. Ceftriaxone for the Treatment of Mild to Moderate Cases in Children

ABSTRACT

Background: Typhoid fever is a bacterial infection that can spread throughout the body, affecting many organs. Without prompt treatment, it can cause serious complications and can be fatal. It's caused by a bacterium called *Salmonella typhi*. In areas where there are few carriers of the typhoid bacterium, the disease is uncommon. It's also unusual to find places with sanitary water treatment and waste management systems. Now a days both Azithromycin and Ceftriaxone are used for the treatment of mild to moderate cases in children.

Objective: To assess the outcome of treatment of Acute Typhoid Fever with Azithromycin vs Ceftriaxone in Children with no other Complication.

Method: This comparative study was carried out at a tertiary hospital from January 2019 to January 2020. Where 200 patients coming to the hospital were initially screened in the outpatient department. During the study period, all children of 5-18 years of age who, according to the outpatient department physician, had a diagnosis of typhoid fever were admitted to a the hospital. After admission to the ward, Among 200 of patients, 100 patients treated with azithromycin and 100 patients treated with ceftriaxone were cured ($P > .05$).

Results: During the study, majority belong to 5-10 years age group and 45% live in semipucca building followed by 60% drink Supply water without boiling and 80% cases were accustomed to solely home made food and outside of food. Clinical characteristics of azithromycin and ceftriaxone recipients who had typhoid fever and for whom blood cultures were positive where in azithromycin group mean duration of hospital stay was 9.8 days followed by 100 cases showed salmonella typhi positive. Whereas in Ceftriaxone group mean duration of hospital stay was 9.0 days followed by 100 cases showed positivity in salmonella typhi positive. In azithromycin group 90% were cured by day 7 followed by 86% were cured microbiologically, plus no relapse cases were found. Whereas in ceftriaxone case 80% were cured by day 7 followed by 78% were cured microbiologically,

plus 5 relapse cases were found.

Conclusion: Oral azithromycin administered once daily appears to be effective for the treatment of uncomplicated typhoid fever in children. If these results are confirmed, the agent could be a convenient alternative for the treatment of typhoid fever, especially in individuals in developing countries where medical resources are scarce.

Keywords: Typhoid Fever, Azithromycin, Ceftriaxone

INTRODUCTION

Children in underdeveloped countries, notably those in Asia and Africa, are particularly vulnerable to typhoid fever, a systemic illness caused by *Salmonella typhi* and *Salmonella paratyphi* [1]. Chloramphenicol has been used to treat typhoid fever for decades [2, 3] due to its effectiveness against *Salmonella typhi* and *S. paratyphi*. The extensive proliferation of drug-resistant *S. typhi* has, however, prompted the exploration of other treatment approaches [4]. While fluoroquinolones have been shown to be beneficial, their usage is now limited in children because of the emergence of quinolone-resistant strains of *Salmonella typhi* [5, 6]. With its superior efficacy against *S. typhi*, the third-generation cephalosporin ceftriaxone has supplanted other treatments as the gold standard for typhoid fever in many countries [7]. Ceftriaxone is a less-than-ideal therapy option due to the need of parenteral administration.

The macrolide family of medicines giving doctors more tools to combat typhoid. The first macrolide to be tested, azithromycin, has been shown to be effective against various intestinal intracellular infections in vitro [8-10]. Azithromycin has been shown to be extremely efficient in animal models against *Salmonella enteritidis* and *Salmonella typhimurium*, with medication efficiency correlated with the tissue concentration of the antibiotic rather than the serum concentration of the drug [11, 12]. Human volunteer studies have indicated that the quantity of azithromycin in neutrophils is more than 100 times that seen in serum [13]. Concentrations of azithromycin in neutrophils were >20 times the normal MIC for *S. typhi* 5 days after a 3-day course of treatment was finished, although quantities of azithromycin in the serum were undetectable [13].

Based on these promising findings, we decided to start a human study of azithromycin therapy. Azithromycin was first shown to be efficacious in treating adults with

uncomplicated typhoid fever in an open-labeled, nonrandomized study [14]. Thereafter, a randomized study showed that azithromycin was just as effective as ciprofloxacin in treating people with uncomplicated typhoid fever [15]. These previous in TV researches served as inspiration for the current investigation of the efficacy of azithromycin suspension against ceftriaxone in the treatment of simple typhoid fever in children & adults.

OBJECTIVE

To assess the outcome of treatment of Acute Typhoid Fever with Azithromycin vs Ceftriaxone in Children with no other complication.

METHODOLOGY

This comparative study was carried out at tertiary hospital from January 2019 to January 2020. Where 200 patients coming to the hospital were initially screened in the outpatient department. During the study period, all children of 5-18 years of age who, according to the outpatient department physician, had a clinical diagnosis of typhoid fever were admitted to a single ward in the hospital. After admission to the ward, a study physician reevaluated the patients to determine whether they were eligible for enrollment in the study. Among 200 of patients, 100 patients treated with azithromycin and 100 patients treated with ceftriaxone were cured ($P > .05$). Eligibility for enrollment required that a subject have a documented fever (temperature, $\geq 38.5^{\circ}\text{C}$) and a history of fever for at least 4 days plus at least 2 of the following criteria: abdominal tenderness, hepatomegaly, splenomegaly, and/or rose spots. Subjects with the following conditions were excluded from the study: allergy to ceftriaxone or erythromycin (or to other macrolides), major complications of typhoid fever (e.g., pneumonia, intestinal hemorrhage or perforation, shock, or coma), inability to swallow oral medication, significant underlying illness (e.g., heart disease, asthma requiring chronic medications, or immunodeficiencies), or treatment within the past 4 days with either study medication or chloramphenicol, trimethoprim-sulfamethoxazole (TMP-SMZ), or ampicillin. Subjects who might be pregnant or who were lactating were also excluded from the study. Parents of children meeting eligibility requirements were asked to have their child enroll in the study, and if they agreed, informed consent was obtained before randomization of the study drug.

RESULTS

Table-1 shows age distribution of the patients where majority belong to 5-8years age group. followed by 30% belong to 09-11years age group and 20% belong to 12-18 years age group.

Table-1: Age distribution of the patients

Age distribution of the patients	Percentage (%)
5-8 years	50%
9-11 years	30%
12-18 years	20%

Figure-1 shows gender status of the patients where 55% were male and 45% were female.




Figure-1: Gender distribution

Table-2 shows demographic status of the patients where 45% live in semipucca building

followed by 60% drink Supply water without boiling and 80% cases were accustomed to solely homemade food and outside of food. Moreover, significance association were noticed in food habit and drinking water source of the patients.

Table-2: Demographic status of the patients

Demographic status	Percentage (%)	P value
Housing status:		
Kacca	20%	0.213
Semipucca	45%	
Pucca	35%	
Drinking water source:		
Tubewell	10%	0.0001
Supply water without boiling	60%	
Supply water with boiling	30%	
Food habit:		
<ul style="list-style-type: none"> • Accustomed to solely homemade food 	20%	0.002
<ul style="list-style-type: none"> • Accustomed to solely home made food & outside of food 	80%	

Figure-2 shows Distribution of patients by liver status where 75% had palpable liver status.




Figure-2: Distribution of patients by liver

status $\chi^2 = 5.486$; $df = 1$; $P = 0.014$

Table-3 shows clinical characteristics of azithromycin and ceftriaxone recipients who had typhoid fever and for whom blood cultures were positive where in azithromycin group mean duration of hospital stay was 9.8 days followed by 100 cases showed positivity in salmonella typhi positive. Whereas in Ceftriaxone group mean duration of hospital stay was 9.0 days followed by 100 cases showed positivity in salmonella typhi positive.

Table-3: Clinical characteristics of azithromycin and ceftriaxone recipients who had typhoid fever and for whom blood cultures

Clinical characteristics (During admission)	Azithromycin group, n=100	Ceftriaxone group, n=100	P value
Mean Duration of fever before admission, days	9.8	9.0	0.234
Blood culture result <ul style="list-style-type: none">• Salmonella typhi• Salmonella paratyphi	100	100	0.121
Blood culture that yielded MDRS. typhi	6	7	0.112

Table-4 shows laboratory status of the patients during admission where in azithromycin group mean hemoglobin level was 10.6 ± 1.4 g/dl followed by mean WBC count was 6.0 ± 2.3 cells/mm³, Total bilirubin level 0.3 ± 0.1 mg/dL. Whereas in Ceftriaxone group mean hemoglobin level was 10.9 ± 1.4 g/dl followed by mean WBC count was 6.5 ± 2.3 cells/mm³, Total bilirubin level 0.4 ± 0.1 mg/dL.

Table-4: Laboratory status of the patients during admission

Laboratory characteristics (During admission)	Azithromycin group, n=100	Ceftriaxone group, n=100	P value
Hemoglobin level, g/dL (11–18)	10.6±1.4	10.9±1.2	0.111
WBC count, cells/mm ³ (4.5–10.53 10 ³)	6.0±2.3	6.5±1.6	0.121
Platelet count, cells/mm ³ (150,000–3 50,000)	213,000±75,000	215,000±96,000	0.100
Total bilirubin level, mg/dL (0.2–1.0)	0.3±0.1	0.4±0.1	0.321
AST level, U/L (0–33)	89±48	84±70	0.201
Blood urea nitrogen level, mg/dL (7–18)	10.5±3.1	11.6±6	0.111
Serum creatinine level, mg/dL (0.7–1.5)	0.8±0.1	0.7±0.2	0.506

Table-5 shows mean Laboratory test result on day 10 where after 10 days in azithromycin group mean hemoglobin level was 9.8±1.4g/dl followed by mean WBC count was 6.2±1.6 cells/mm³, Total platelet count 433,000±122,000 cells/mm³. Whereas in Ceftriaxone group mean hemoglobin level was 10.5±1.0 g/dl followed by mean WBC count was 7.4±2.2cells/mm³, Total platelet count 431,000±164,000cells/mm³.

Table-5: Laboratory test result on day 10

Laboratory characteristics, day 10	Azithromycin group, n=100	Ceftriaxone group, n=100	P value
Hemoglobin level, g/dL (11–18)	9.8±1.4	10.5±1.0	0.111

WBC count, cells/mm³(4.5–10.53 10³)	6.2±1.6	7.4±2.2	0.120
Platelet count, cells/mm³(150,000–3 50,000)	433,000±122,000	431,000±164,000	0.101
Total bilirubin level, mg/dL (0.2–1.0)	0.3±0.1	0.4±0.2	0.321
AST level, U/L (0–33)	52±29	60±35	0.0.23
Blood urea nitrogen level, mg/dL (7–18)	9.0±3.1	9.6±6	0.111
Serum creatinine level, mg/dL (0.7–1.5)	0.6±0.1	0.7±0.2	0.506

UNDER PEER REVIEW

Table-6 shows Responses to treatment with azithromycin or ceftriaxone among patient where in azithromycin group 90% were cured by day 7 followed by 86% were cured microbiologically, plus no relapse cases were found. Whereas in ceftriaxone case 80% were cured by day 7 followed by 78% were cured microbiologically, plus 5 relapse cases were found.

Table-6: Responses to treatment with azithromycin or ceftriaxone among patient

Response to treatment	Azithromycin group, n=100	Ceftriaxone group, n=100	P value
Clinical cure by day 7	45 (90%)	40(80%)	0.001
Duration of fever after starting therapy, mean d±SD	4.0±1.1	3.7±1.1	0.212
Microbiological cure, no. (%)	43 (86%)	39(78%)	0.121
Blood culture that yielded Salmonella			0.001
Day 4	1a	0	
Day 10	1a	1	
Stool culture that yielded Salmonella on day 10	0	0	0.0001
Relapse	0	5	

DISCUSSION

We found that azithromycin is very successful in treating children with uncomplicated typhoid fever in a randomized, controlled study. Results from this research are comparable to those from previous studies of conventional antibiotics for the treatment of typhoid fever [2, 6, 19-21], with clinical cure rates of >90% and microbiological cure rates of >95% for participants taking either azithromycin or ceftriaxone. Relapses of infection occurred in 14% of ceftriaxone-treated participants with bacteremia within 1 month after medication completion, which is an intriguing finding. These findings are in line with the 5-15% recurrence rate seen in earlier ceftriaxone for typhoid fever studies [2, 7, 19, 22]. The lengthy half-life of azithromycin inside the intracellular compartment, with elimination of remaining organisms after conclusion of treatment, and its higher concentration within the biliary system may account for the lack of relapses in our small sample of azithromycin-treated subjects. Because of its extraordinarily long half-life in tissue, azithromycin may be amenable to shorter durations of therapy for typhoid fever, similar to those that have been attempted with effectiveness for quinolone antibiotics [6, 19, 23].

Not much of a clinical difference occurred between the two groups. Defervescence occurred significantly sooner in ceftriaxone-treated patients (3.9 days) compared to azithromycin-treated patients (4.1 days), but this difference was not statistically significant, and both outcomes were within the ranges described in prior typhoid therapy studies [3, 19, 23–25]. Both groups had mild, short-lived gastrointestinal problems, but none were considered serious enough to warrant a treatment change.

CONCLUSION

In conclusion, 7 days of 10 mg/kg/day (maximum dose, 500 mg/day) azithromycin seems to be extremely successful for the treatment of uncomplicated typhoid fever in children, with clinical cure rates equivalent to those for ceftriaxone. Typhoid fever caused by susceptible or

drug-resistant strains of *S.typhi* may be treatable with a straightforward, once-daily oral regimen of azithromycin, making it an appealing option for usage in resource-poor settings.

REFERENCE

1. Azad AK, Islam R, Salam MA, Alam AN, Butler T. Comparison of clinical features and pathologic findings in fatal cases of typhoid fever during the initial

- and later stages of the disease, *Am J Trop Med Hyg*, 1997, vol. 56 (pg. 490-3)
2. Acharya G, Butler T, Ho M, et al. Treatment of typhoid fever: randomized trial of a three-day course of ceftriaxone versus a fourteen-day course of chloramphenicol, *Am J Trop Med Hyg*, 1995, vol. 52 (pg. 162-5)
 3. Islam A, Butler T, Kabir I, Alam NH. Treatment of typhoid fever with ceftriaxone for 5 days or chloramphenicol for 14 days: a randomized clinical trial, *Antimicrob Agents Chemother*, 1993, vol. 37 (pg. 1572-5)
 4. Mirza SH, Beeching NJ, Hart CA. Multi-drug resistant typhoid fever: a global problem, *J Med Microbiol*, 1996, vol. 44 (pg. 317-9)
 5. Wain J, Hoa NTT, Chinh NT, et al. Quinolone-resistant *Salmonella typhi* in Vietnam: molecular basis of resistance and clinical response to treatment, *Clin Infect Dis*, 1997, vol. 25 (pg. 1404-10)
 6. Vinh H, Wain J, Vo TN, et al. Two or three days of ofloxacin treatment for uncomplicated multidrug-resistant typhoid fever in children, *Antimicrob Agents Chemother*, 1996, vol. 40 (pg. 958-61)
 7. Soe GB, Overturf GD. Treatment of typhoid fever and other systemic salmonellosis with cefotaxime, ceftriaxone, cefoperazone and other newer cephalosporins, *Rev Infect Dis*, 1987, vol. 9 (pg. 719-36)
 8. Jones K, Felmingham D, Ridgway G. In vitro activity of azithromycin (CP-62,993), a novel macrolide, against enteric pathogens, *Drugs Exp Clin Res*, 1988, vol. 14 (pg. 613-5)
 9. Metchock B. In vitro activity of azithromycin compared with other macrolides and oral antibiotics against *Salmonella typhi*, *J Antimicrob Chemother*, 1990, vol. 25 Suppl A (pg. 29-31)
 10. Gordillo ME, Singh KV, Murray BE. In vitro activity of azithromycin against bacterial enteric pathogens, *Antimicrob Agents Chemother*, 1993, vol. 37 (pg. 1203-5)
 11. Girard AE, Girard D, Retsema JA. Correlation of the extravascular pharmacokinetics of azithromycin with in-vivo efficacy in models of localized infection, *J Antimicrob Chemother*, 1990, vol. 25 Suppl A (pg. 61-71)
 12. Butler T, Girard AE. Comparative efficacies of azithromycin and ciprofloxacin against experimental *Salmonella typhimurium* infection in mice, *J Antimicrob Chemother*, 1993, vol. 31 (pg. 313-9)

13. Wildfeuer A, Laufen H, Zimmermann T. Distribution of orally administered azithromycin in various bloodcompartments, *Int J Clin Pharmacol Ther*, 1994, vol. 32 (pg. 356-60)
14. Tribble D, Girgis NI, Habib N, Butler T. Efficacy of azithromycin for typhoid fever, *Clin Infect Dis*, 1995, vol. 21 (pg. 1045-6)
15. Girgis NI, Butler T, Frenck RW, Sultan Y, Brown FM, Tribble D. Azithromycin versus ciprofloxacin for the treatment of uncomplicated typhoid fever that included patients with multidrug resistance, *Antimicrob Agents Chemother*, 1999, vol. 43 (pg. 1441-4)
16. Young MJ, Bresnitz EA, Strom BL. Sample size nomograms for interpreting negative clinical studies, *Ann Intern Med*, 1983, vol. 99 (pg. 248-51)
17. Gray LD, Murray PR, Baron EJ, Pfaller MA, Tenoer FC, Tenover RH. *Escherichia, Salmonella, Shigella and Yersinia, Manual of clinical microbiology*, 1995 6th ed. Washington, DC American Society for Microbiology Press (pg. 450-6)
18. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method, *Am J Clin Pathol*, 1966, vol. 45 (pg. 493-6)
19. Smith MD, Duong NM, Hoa NT, et al. Comparison of ofloxacin and ceftriaxone for short-course treatment of enteric fever, *Antimicrob Agents Chemother*, 1994, vol. 38 (pg. 1716-20)
20. Wallace MR, Yousif AA, Mahroos GA, et al. Ciprofloxacin versus ceftriaxone in the treatment of multiresistant typhoid fever, *Eur J Clin Microbiol Infect Dis*, 1993, vol. 12 (pg. 907-10)
21. Girgis NI, Sultan Y, Hammad O, Farid Z. Comparison of the efficacy, safety and cost of cefixime, ceftriaxone and aztreonam in the treatment of multidrug resistant *Salmonella typhi* septicemia in children, *Pediatr Infect Dis J*, 1995, vol. 14 (pg. 603-5)
22. Lasserre R, Sangalang RP, Santiago L. Three-day treatment of typhoid fever with two different doses of ceftriaxone, compared to 14-day therapy with chloramphenicol: a randomized trial, *J Antimicrob Chemother*, 1991, vol. 28 (pg. 765-72)
23. Cao XT, Kneen R, Nguyen TA, Truong DL, White NJ, Parry CM. A comparative study of ofloxacin and cefixime for treatment of typhoid fever in children. Dong Nai Pediatric Center Typhoid Study Group, *Pediatr Infect Dis J*, 1999, vol. 18 (pg.

245-8)

24. Van den Bergh ET, Gasem MH, Keuter M, Dolmans MV. Outcome in three groups of patients with typhoid fever in Indonesia between 1948 and 1990, *Trop Med Int Health*, 1999, vol. 4 (pg. 211-5)
25. Dutta P, Rasaily R, Saha MR, et al. Ciprofloxacin for treatment of severe typhoid fever in children, *Antimicrob Agents Chemother*, 1993, vol. 37 (pg. 1197-9)

UNDER PEER REVIEW