

STUDIES ON THE PRESERVATION OF KOKUM (*GARCINIA INDICA*) RIND JUICE AT AMBIENT STORAGE CONDITION

Abstract

The study aimed to determine the effect of different concentrations of sodium benzoate at ambient storage condition on kokum rind juice. For this experiment, Factorial Completely Randomized Design (F.C.R.D.) was used. During the investigation, the chemical parameter, microbial analysis and sensory qualities of kokum rind juice were studied. During the 3 months of ambient storage condition, analysis was conducted at the interval of one month. The chemical parameter like acidity, TSS, reducing sugar, total sugar, anthocyanins and microbial analysis, sensory qualities changed significantly. From the experiment it can be concluded that the kokum rind juice could be stored at 1500 ppm sodium benzoate concentration in a good acceptable condition at ambient storage condition.

Keywords: Kokum, Preservation, Sodium benzoate

Introduction

“Kokum (*Garcinia indica* Choisy) is a major indigenous tree spice crop that originated and is grown in India's Western Ghats”(Padhye et al. 2009). “Maharashtra's South Konkan region, Coorg, Wynad, and Goa, and is found in evergreen and semi-evergreen woods as well as as a home garden tree” (Subash Chandran, 2005). The tree grows extensively in the Konkan region of Maharashtra.

“As for a chemical composition of Kokam fruit is concerned, it contains 80% moisture, 2% protein, 2.8% tannin, 5% pectin, 14% crude fiber, 4.1% total sugars, 1.4% fat, 2.4% pigment, 22% hydroxy citric acid, 0.06% ascorbic acid 5.90% organic acid, 2.40% anthocyanins, 2.60% ash and 1% starch” (Krishnamurthy et al. 1982, Sayeed et al. 2009 and Nayak et al. 2010)

“Kokum fruits (*Garcinia indica*) contain the important bioactive compounds like anthocyanin (ACNs), hydroxycitric acid (HCA), garcinol (GL), and isogarcinol (IGL) at a significant quantity” (Nainegali 2021). “It exhibits various phytochemical properties like anti-ulcerogenic, cardio protective, anticancer, chemo preventive, free radical scavenging and anti-obesity effects. *Garcinia* known for medicinal properties is also used for curing piles, dysentery, tumour pains and cardiac problems. Even fat reducing tablets and capsules based on *Garcinia* are available in the market” (Rasha et al., 2015)

“Today, the preservation of fruit juice has become a business activity of great importance, and countries with plentiful fruit resources and a short harvest season place more emphasis on well-

established storage to preserve fruit quality, extend its shelf life and preserve fruit juice for off season use” (Tasnim et al. 2010) Fruit juice must be conserved in order to be used later in the creation of beverages with juice as the main ingredient. However, due to the high nutritional content of fruit juices, storage of fruit juices at room temperature without the addition of any chemical preservatives or pasteurization is not feasible.

Fruit juices are refreshing and retain characteristic taste and aroma even after a few months of their preparation into a beverage. The kokum rind juice has a very short storage life. Generally, after extraction of juice from kokum it starts deteriorating. Kokum juice is generally utilized for the preparation of different products like **Ready to serve**, syrup, jelly, solkadi etc. however, it has a very poor storagelife therefore a systematic research work is needed to increase the storage life of kokum rind juice conversion into a different value-added product. It is highly necessary to study the storage life of kokum fruit juice which would be useful for value addition. The storage life of kokum rind juice could be extended by using chemical preservatives as well as low-temperature storage. It is highly necessary to extend the storage life of kokum juice with preservatives under different storage conditions.

The chemical preservation technique is used primarily to prevent microbial spoilage of fruit juices during storage, both in retail stores and in the consumer's home. Benzoic acid (E210) and sodium benzoate (E211) are approved food preservatives with an acceptable daily intake of 5 mg/kg body weight by the FAO/WHO due to their long history of safe use. Sodium benzoate, with its broad antibacterial activity, non-volatility and water solubility, is widely used as a preservative especially in coloured fruit drink. The sodium benzoate is effective against yeast, mold and bacteria. It is most suitable for use as an antimicrobial agent in food and beverages which naturally are in the pH range below 4.5 per cent, compared to other preservatives (Kaddumukasa et al. 2017).

Materials and Methods


The fruits required for conducting research were procured from Dapoli. The fresh, ripe and mature fruits were selected and brought to laboratory for conducting the research

1.1 Preparation of fruits and juice extraction

Fully ripe, fresh and sound kokum fruit were selected for extraction of kokum juice. The selected fruits were washed with water and sodium hypo-chloride to remove dirt and dust. The kokum fruits were then cut with help of kokum fruit cutter machine. Then the fruit rind and pulp with seed were separated. The kokum rind juice was extracted by using basket press. The extracted kokum juice was then strained through double fold muslin cloth. The sodium benzoate was then added in the kokum rind juice as per the treatment. Kokum juice was then filled in pre-sterilized glass bottles and the bottles were then sealed with threaded cap. The product, thus prepared was labeled and stored at ambient storage conditions for further investigations.

FLOW CHART - I

Preparation of kokum rind juice



1.2 Chemical parameter of kokum rind juice

1.2.1 Total Soluble Solids

The total soluble solids were determined by using Hand Refractometer (Atago Japan, 0-320B) and the values were corrected at 20 °C with the help of temperature correction chart (AOAC, 2020).

1.2.2 Titratable acidity

The titratable acidity was assessed by the standard method of Ranganna (1986). 10ml of sample was diluted with 100 ml distilled water. Further, 10ml of this aliquot was titrated against 0.1N NaOH solution to a pink end point using phenolphthalein as an indicator. The acidity as citric acid was calculated by the following expression:

Titratable acidity(%)

$$= \frac{\text{Normality of alkali} \times \text{Titre reading} \times \text{Volume made} \times \text{Equivalent weight of acid}}{\text{Weight of sample taken} \times \text{Volume of sample taken for estimation} \times 1000} \times 100$$

1.2.3 Reducing sugars

The reducing sugars were determined by the method of Lane and Eynon (1923) as described by Ranganna (1986). "25 ml weight of sample was taken in 250 ml volumetric flask. To this, hundred ml of distilled water was added and the contents were neutralized by 1 N sodium hydroxide. Then, 2 ml of 45 percent lead acetate was added to it. The contents were mixed well and kept for 10 minutes. Two ml of 22 percent potassium oxalate was added to it to precipitate the excess of lead. The volume was made to 250 ml with distilled water and solution was filtered through Whatman No. 4 filter paper. This filtrate was used for determination of reducing sugars by titrating it against the boiling mixture of Fehling 'A' and Fehling 'B' solutions (5 ml each) using methylene blue as indicator to a brick red end point. The results were expressed on per cent basis". [21]

$$\text{Reducing sugars (\%)} = \frac{\text{factor} \times \text{Dilution}}{\text{Titre reading} \times \text{Weight of sample}} \times 100$$

1.2.4 Total sugar

"For inversion at room temperature, a 50 ml aliquot of clarified solution was transferred to 250 ml volumetric flask, to which, 10 ml of 50 percent HCl was added and then allowed to stand at room

temperature for 24 hrs. It was then neutralized with 40 percent NaOH solution. The volume of neutralized aliquot was made to 250 ml with distilled water". [21] This aliquot was used for determination of total sugars by titrating it against the boiling mixture of Fehling 'A' and Fehling 'B' (5ml each) using methylene blue as indicator to a brick red end point. The results were expressed on percent basis.

$$\text{Total sugars (\%)} = \frac{\text{factor} \times \text{Dilution}}{\text{Titre reading} \times \text{weight of sample}} \times 100$$

1.2.5 Anthocyanin

The anthocyanin content in juice was determined as per the procedure described by (Shriwastava and Kumar, 2015). Kokum juice was diluted with 0.1N HCl and allowed to equilibrate in the dark for 1hr. The Optical Density was recorded at 510 nm using spectrophotometer and total anthocyanin were estimated as given below,

$$\text{Total O.D. (Optical Density)/100ml} = \frac{\text{O.D. (Spectrophotometer reading)} \times \text{volume made up} \times 100}{\text{ml of juice taken}}$$

$$\text{Total Anthocyanins (mg/100 ml)} = \frac{\text{Total O.D} \times 100\text{ml}}{87.3}$$

1.3 Microbial analysis

The microbial analysis of kokum fruit juice was carried out at 0 day and after 90 days of storage as per the method described by Kiiyukia (2003). Nutrient agar media was prepared by weighing required quality of nutrient agar. The media was autoclaved at 121°C for 20 min. then temperature of media was lowered to 40°C and used for plating. The plating was carried out with 1 ml. sample in sterile petri plate under the Laminar Air Flow. The sample of each treatment was taken on a separate petri plate, followed by pouring of approximately 20 ml. of media (35 to 40°C) on the sample and mixing was done by tilting plate properly, the plates were sealed with parafilm and incubated at 37°C for 24 hrs.

1.4 Sensory evaluation

The sensory quality of kokum rind juice was studied by preparing an RTS (6 per cent juice, 12°B TSS and 0.28 per cent acidity and remaining water) from various treatments. The sensory evaluation was done by using 9-point Hedonic scale. The tasting panel was consisting of 10 members. They were asked to evaluate the color, flavor and overall acceptability by a scoring rate on a 9 scale. 9= Like extremely, 8=Like very much, 7= Like moderately, 6= Like slightly, 5= neither like nor dislike, 4=Dislike slightly, 3= Dislike moderately, 2=Dislike very much and 1=Dislike extremely. The different preferences as indicated by scores were evaluated by statistical methods.

Result and discussion

Total soluble solids (°B)

The data presented in Table 1 shows that TSS content of kokum rind juice increase significantly in all treatment during storage period of 90 days.. The TSS content of kokum rind juice was significantly influenced by the treatments throughout the ambient storage period of 90 days. The maximum (12.92 °B) mean value of TSS was observed in the T1 (500 ppm) followed by the treatment, T2(12.69) and T3(12.49). The treatment T5 (1500 ppm) showed a minimum (12.23°Brix) mean value for the TSS content and the treatment T4 was at par with the treatment T5. As regards the storage, an increasing trend in the TSS content of kokum rind juice was notice 90 days of ambient storage condition. At the initial stage, the mean TSS was 12.5 °Brix which was increased up to 12.94 °Brix at 90 days ambient storage condition. The interaction effect between different treatments and ambient storage condition on the TSS of kokum rind juice was found to be statistically significant. After 90 days of ambient storage condition, the maximum (13.60 °B) was recorded in the treatment T1.The minimum (12.47 °B)TSS was observed in the treatment of T5 at ambient storage condition, however, it was at par treatment, except the T3 and T4. This condition could be due to the fact that the higher-level sodium benzoate reduced the biochemical change in kokum rind juice during storage. Investigation reveals that increase in TSS is probably due to the hydrolysis of polysaccharides into monosaccharides and

the concentration of juice due to dehydration. The increase trend in TSS during storage was also reported by Shahnawaz et al. (2013) in orange juice and, Das (2009) in jamun processed products. Majumdar et al. (2011) “in bottle gourd-basil leaves juice, Majumdar et al. (2012) in ash gourd-mint leaves juice, by Kaur and Aggarwal (2014) in bitter gourd juice preserved with different chemical preservatives”.

Titrateable acidity (%)

It could be observed in Table 2 that different concentrations of sodium Benzoate showed a significant effect on the titrateable acidity of kokum rind juice during ambient storage conditions. It was observed from that on an average highest mean acidity(3.80%) was observed in the treatment T1 (500 ppm) which was at par with the treatment T2(3.71). The lowest mean titrateable acidity(3.53 %) was observed in the treatment T5 (1500 ppm) which was at par with the treatments T4 and T3.As regards the ambient storage period an increasing trend in titrateable acidity content in kokum rind juice was noticed and the mean acidity significantly increased from 3.44 per cent at the initial stage to 3.84 per cent after 90 days of storage. The increase in titrateable acidity might be due to the formation of organic acids by the degradation of ascorbic acid. An increase in acidity might be attributed due to the breakdown of pectin into pectinic acid or due to the formation of acid by the breakdown of polysaccharides or oxidation of reducing sugars. Similar findings of increasing acidity were observed by Hussain et al., (2011) in apple and apricot blend juice Desai (1991) in ber juice showed at ambient and low temperature, Bharwal and Shreera (2009)in hill lemon juice. Majumdar et al. (2012),Kaur and Aggarwal (2014) and Gajera and Joshi (2015) observed “increase in titrateable acidity during storage in ash gourd-mint leaves juice, bitter gourd juice and bottle gourd-based blend juice, respectively”.

Reducing sugars(%)

It was clearly seen from the data that in Table 3 there was a significant difference among the treatments with respect to reducing sugars content in kokum rind juice. The treatment T1 (500 ppm) showed the significantly highest (9.05%) mean reducing sugar, followed by the treatment T2. The minimum (8.40%) mean was observed in the treatment T5 which was at par with the treatments T4 (8.47%) and T3 (8.55%).During ambient storage, the lowest reducing sugar content (8.26%) was rerecorded at 0 days. A significant increase in reducing sugar from 8.26 to 9.12 per cent was noticed up to 90 days of ambient storage condition. As regards, it is observed that in reducing sugar content increased with the increase in storage period in all the treatments. It could be attributed to the gradual inversion of non-reducing sugars into reducing sugars in an acidic medium. . The

increase in reducing sugars during storage might be due to the rapid hydrolysis of polysaccharide and their subsequent conversion to reducing sugars. A similar result was reported by Palaniswamy and Muthukrishnan (1974), who observed an increase in reducing sugar content in lemon juice during storage, Shrestha and Bhatia (1982) in aonla juice. Identical results were also reported by Bharwal and Shreera (2009) in hill lemon juice and Mhapralkar (2007) in jamun juice.. Mehmood et al. (2008) observed an increase in reducing the sugars of apple juice preserved with chemical preservatives and stored at ambient temperature for three months. Babu (2015) and Kadam and Lele (2016) also reported “an increase in reducing sugars of ash gourd-mint leaves juice, ripe pumpkin pulp and ash gourd-carrot juice during storage, respectively”

Total sugars (%)

It is evident from the data Table 4 that the different concentrations of sodium benzoate exhibited a significant effect on total sugars during storage at ambient conditions. The treatment T1(500) showed the significantly highest (10.09%) mean of total sugar followed by treatments T2 and T3. The lowest (9.58%) total sugar content was observed in the treatment T5 (1500 ppm) which was at par with treatment T4 (1250 ppm). The sodium benzoate level of either 1250 ppm or 1500 ppm did not exhibit a significant increase in the total sugar content of kokum rind juice. As regards, ambient storage conditions, an increase in total sugar content was observed during 90 days of ambient storage condition. The total sugars significantly increased from 9.46 per cent at the initial stage to 10.07 per cent up to 90 days of ambient storage condition.. The increase in total sugars might be due to the hydrolysis of polysaccharides like pectin, cellulose, starch, etc. and its conversion into simple sugars. The results obtained were similar to the findings of Bharwal and Shreera (2009) in hill lemon juice. Chobe (1991) in pomegranate juice and Patil and Jadhav (2001) in sweet orange juice

Anthocyanins (mg/100g)

It could be observed from the Table 5 that there was a significant difference among the treatments with respect to anthocyanin content of kokum rind juice. The treatment T5 (1500 ppm) showed the highest (2.48 mg/100g) mean anthocyanin content which was at par with the treatment T4 . The minimum (1.85 mg/100g) mean was observed in the treatment T1. It is clear from the data presented data that the storage period had a significant effect on the anthocyanin content of kokum rind juice during 90 days of storage. At 0-day stage the anthocyanin, content was 2.52 mg/100g and it was decreased up to 1.92 mg/100g 90 days at storage period . After 90 days of storage period, the maximum (2.44 mg/g) anthocyanin was observed in the treatment T5 which was at par with treatment T4 . The minimum (1.21 mg/g) anthocyanin was observed on in the treatment T1 at 90

days storage. Anthocyanins were decreased during storage, this might be due to the hydrolysis of anthocyanins at higher temperature Rhim (2002).

Microbial analysis of kokum rind juice

Considering sodium benzoate concentrations, it was served that, the results were significant throughout the storage period with respect to yeast and bacterial count. From initial day of storage there was no growth of yeast and bacteria in treatment . At 3-month storage the lowest yeast and bacterial growth was found in treatment T5 which was significantly superior over rest of the treatments. While, highest yeast and bacterial load was recorded in T1 at 3-month storage, yeast and bacterial population was increased rapidly. From above findings it could be seen that, as the concentration of sodium benzoate increases, yeast and bacterial growth get decreased. Similar findings have been reported by, Oladipo et al. (2010) in some Nigerian fruit juices and Hussain et al.(2011) in apple and apricot blended juice. With respect to effect of storage conditions on the yeast and bacterial growth, it was observed that, as the storage period increases the yeast and bacterial load get increased at faster rate in ambient temperature as compared to cold storage . Similar findings have been reported by Emamifar (2010) in orange juice.

Sensory evaluation

The sensory score for changes in colour, flavour and overall acceptability of kokum rind juice during storage are presented in Table 7 to 9 . From the table it was observed that, mean score of colour, flavour and overall acceptability of kokum juice decreased with increased in storage period. Decrease in juice colour, flavour and overall acceptability with increased in storage period may be due to slight browning and increase of acids or may be due to effect of temperature on storage quality of juice, micro fermentation process at the end of storage. Similar findings have been reported by Khurdiya and Roy (1985), Bhandari (2004) in jamun juice, Shakoor et al., (2013) in strawberry juice.

Table 1: Effect of sodium benzoate concentrations on Total Soluble Solid (°B) of kokum rind juice under ambient storage conditions.

Treatment	TSS (°B)				
	Storage period (Days)				
	0	30	60	90	Mean
T1	12.03	12.67	13.40	13.60	12.92
T2	12.04	12.63	12.87	13.23	12.69
T3	12.04	12.47	12.67	12.77	12.49
T4	12.06	12.40	12.37	12.63	12.36

T5	12.07	12.13	12.23	12.47	12.23
Mean	12.05	12.46	12.71	12.94	
	S.Em ±			CD at 5%	
Treatment (T)	0.06			0.16	
Storage (S)	0.05			0.14	
Interaction (TxS)	0.11			0.32	

Table 2: Effect of sodium benzoate concentrations on titratable acidity (%) of Kokum rind juice under ambient storage conditions

Treatment	Titratable acidity (%)				
	Storage period (Days)				
	0	30	60	90	Mean
T1	3.43	3.74	3.92	4.12	3.80
T2	3.45	3.66	3.79	3.94	3.71
T3	3.44	3.55	3.73	3.80	3.63
T4	3.44	3.52	3.63	3.69	3.57
T5	3.45	3.46	3.57	3.63	3.53
Mean	3.44	3.59	3.73	3.84	
	S.Em ±			CD at 5%	
Treatment (T)	0.04			0.11	
Storage (S)	0.03			0.09	
Interaction (TxS)	0.07			NS	

Table 3: Effect of sodium benzoate concentrations on Reducing sugars (%) of kokum rindjuice under ambient storage conditions

Treatment	Reducing sugars (%)				
	Storage period (Days)				
	0	30	60	90	Mean
T1	8.24	8.62	9.45	9.86	9.05
T2	8.26	8.49	9.02	9.39	8.79
T3	8.26	8.38	8.68	8.89	8.55
T4	8.27	8.30	8.52	8.80	8.47
T5	8.27	8.28	8.39	8.64	8.40
Mean	8.26	8.42	8.81	9.12	
		S.Em ±		CD at 5%	
Treatment(T)		0.09		0.24	
Storage(S)		0.08		0.22	
Interaction (TxS)		0.16		NS	

Table 4: Effect of sodium benzoate concentrations on total sugar (%) content of kokum rindjuice under ambient storage conditions

Treatment	Total sugar (%)				
	Storage period (Days)				
	0	30	60	90	Mean
T1	9.45	10.00	10.29	10.62	10.09
T2	9.46	9.88	10.02	10.22	9.89
T3	9.46	9.68	9.78	9.91	9.71
T4	9.47	9.56	9.69	9.85	9.64
T5	9.47	9.52	9.62	9.73	9.58
Mean	9.46	9.73	9.88	10.07	
		S.Em ±		CD at 5%	
Treatment(T)		0.058		0.17	
Storage(S)		0.052		0.15	
Interaction (TxS)		0.11		NS	

Table 5: Effect of sodium benzoate concentrations on anthocyanins (mg/100g) of kokum rind juice under ambient storage conditions

Treatment	Anthocyanins (mg/100g)				
	Storage period (Days)				
	0	30	60	90	Mean
T1	2.53	2.02	1.63	1.21	1.85
T2	2.54	2.24	1.99	1.55	2.08
T3	2.51	2.39	2.29	2.08	2.32
T4	2.51	2.44	2.44	2.33	2.43
T5	2.53	2.46	2.48	2.44	2.48
Mean	2.52	2.31	2.17	1.92	
		S.Em ±		CD at 5%	
Treatment(T)		0.05		0.15	
Storage(S)		0.05		0.13	
Interaction (TxS)		0.10		0.29	

Table 6: Effect of sodium benzoate concentrations on microbial count (cfu/ml) of Kokum rind juice under ambient storage conditions

Microbial count (No. × 10 ³ cfu/ml)		
Treatment	Storage period (Days)	
	0	90
T1	ND	3.34
T2	ND	2.91
T3	ND	2.68
T4	ND	2.37
T5	ND	1.76
Mean	ND	2.61
S.Em ±	0	0.03
CD at 5%	0	0.07

Table 7: Effect of sodium benzoate concentrations on sensory score for colour of kokum rind juice stored in ambient conditions

Treatment	Sensory score for colour				
	Storage period (Days)				
	0	30	60	90	Mean
T1	7.24	5.70	5.70	5.30	5.99
T2	7.30	6.20	5.80	5.30	6.15
T3	7.60	6.20	6.14	5.40	6.34
T4	7.60	6.10	5.90	5.70	6.33
T5	7.70	6.30	6.10	5.90	6.50
Mean	7.49	6.10	5.93	5.52	
		S.Em ±		CD at 5%	
Treatment(T)		0.06		0.16	
Storage(S)		0.05		0.14	
Interaction (TxS)		0.11		NS	

Table 8: Effect of sodium benzoate concentrations on sensory score for flavour of kokum rind juice under ambient storage conditions

Treatment	Sensory score for flavour				
	Storage period (Days)				
	0	30	60	90	Mean
T1	7.40	6.50	6.17	5.80	6.47
T2	7.40	6.90	6.46	5.79	6.64
T3	7.60	7.24	6.60	6.30	6.94
T4	7.70	7.10	6.60	6.29	6.93
T5	7.80	7.40	6.69	6.59	7.12
Mean	7.58	7.03	6.50	6.15	
		S.Em ±		CD at 5%	
Treatment(T)		0.10		0.29	
Storage(S)		0.09		0.26	
Interaction (TxS)		0.19		NS	

Table 9: Effect of sodium benzoate concentrations on overall acceptability of kokum rind juice under ambient storage conditions

Treatment	Sensory score for overall acceptability				
	Storage period (Days)				
	0	30	60	90	Mean
T1	7.11	5.95	6.14	5.55	6.19
T2	7.35	6.25	6.35	5.65	6.40
T3	7.45	6.30	6.35	5.57	6.42
T4	7.65	6.30	6.60	6.00	6.64
T5	7.72	6.50	6.76	6.25	6.81
Mean	7.46	6.26	6.44	5.81	
		S.Em ±		CD at 5%	
Treatment(T)		0.03		0.10	
Storage(S)		0.03		0.09	
Interaction (TxS)		0.07		NS	

Conclusion

Based on the experiments performed and the results obtained, the following conclusions were reached. The experiment was to compare the different concentration of sodium benzoate on the storage stability of kokum rind juice. In this study, it is evident that 1500ppm sodium benzoate concentration proved to be a better preservative than other at ambient condition and maintain good chemical characteristics as well as sensory quality up to 90 days of storage period.

REFERENCE

1. A.O.A.C. (2020) Official method of analysis. Association of official analytical chemists, Washington D.C., USA.
2. Babu GN. (2015) Development of beverages from ripe pumpkin (*Cucurbita moschata* Duch ex Poir) pulp. M.Sc. Thesis. Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Solan., 116.
3. Bhandari, S.P.(2004) Studies on physico-chemical composition, grading, storage and processing of jamun (*Syzygium cuminii* (Linn.) Skeels). M.Sc. (Ag.) Thesis, Dr.

B.S.K.K.V.,Dapoli.

4. Bharwal and Shreera Sanjay K.(2009) Standardisation of extraction methods and preservation techniques of hill lemon juice.*Journal of Food Safety and Industrial Research*,68 : 608-610
5. Chobe, R.S. (1991). Studies on extraction, clarification, preservation and storage of pomegranate (*Punica granatum* L.) juice. M.Tech (Ag. Engg.) Thesis , M.P.A.U., Rahuri (Maharashtra), India.
6. Das, J.N. (2009). Studies on storage stability of jamun beverages. *Indian Journal Of Horticulture*. 66.:508-510
7. Desai Prabhu (1991). Studies on juice making in ber (*Zizypus mauritiana* Lam.). M.Sc. (Agri.)thesis submitted to MPKV, Rahuri, (M.S.), India.
8. Emamifar, A. (2010). Evaluation of nanocomposite packaging containing Ag and ZnO on shelf life of fresh orange juice. *Innovative Food Science and Emerging Technologies*, 11: 742-748.
9. Gajera RR, Joshi DC. (2015) Development and quality evaluation of bottle gourd, *Lagenaria siceraria* (Mol.) Standl. based blend juice. *Indian Journal of Natural Products and Resources*;6(3):194-199
10. Hussain, I., A. Zeb and M. Ayub.(2011). Evaluation of apple and apricot blend juice preserved withsodium benzoate at refrigeration temperature. *World J. of Agric. Sci.*,7(2): 136-142.
- K.M. (2010). Quality Assessment of Industrially Processed Fruit Juices available in Dhaka City, Bangladesh. *Malaysian Journal of Nutrition*, 16(3):431-438.
11. Kadam D, Lele S. Studies on the physicochemical and nutritional characteristics of ash gourd-carrot juice. *Nutrafoods* 2016;15:39-47.
12. Kaddumukasa, P.P., Imathiu, S.M., Mathara, J.M. and Nakavuma, J.L.(2017). Influence of physicochemical parameters on storage stability: Microbiological quality of fresh unpasteurized fruit juices. *Food Science & Nutrition*, 5(6) : 1098-1105.

13. Kaur G, Aggarwal P. Storage studies on bitter gourd juice preserved with different chemical preservatives. *International Journal of Research in Engineering and Technology* 2014;3(1):223-227
14. Khurdiya, D.S. and Roy, S.K. 1985. Storage studies on jamun (*Syzygium cumini*) juice and nectar. *J. Fd. Sci. Technol*, 22: 217-220.
15. Kiiyukia, C. (2003) Laboratory manual of food microbiology for Ethiopian health and nutrition research institute. UNIDO project (YA/ETH/03/436/11 -52). : 1-197.
16. Krishnamurthy, N., Lewis, Y. S., and Ravindranath, B. (1982). "Chemical constituents of Kokum fruit rind." *J. Food Sci. Technol.*, 19(3), 97–100
1. Majumdar TK, Wadikar D, Bawa AS.(2012) Development and storage stability of aseptically processed ash gourd-mint leaves juice. *International Food Research Journal*;19(3):823-828
2. Majumdar TK, Wadikar D, Bawa AS. Development and storage stability of aseptically processed ash gourd-mint leaves juice. *International Food Research Journal* 2012;19(3):823-828.
3. Majumdar TK, Wadikar DD, Vasudish CR, Premavalli KS, Bawa AS. (2011)Effect of storage on physico-chemical, microbiological and sensory quality of bottle gourd-basil leaves juice. *American Journal of Food Technology*;6(3):226-234.
4. Majumdar TK, Wadikar DD, Vasudish CR, Premavalli KS, Bawa AS. Effect of storage on physico-chemical, microbiological and sensory quality of bottle gourd-basil leaves juice. *American Journal of Food Technology* 2011;6(3):226-234
5. Mehmood Z, Zeb A, Ayub M, Bibi N, Badshah A, Ullah I (2008) Effect of pasteurization and chemical preservatives on the quality and shelf stability of apple juice. *Amer J. Food Tech.* 3(2): 147- 153.
6. Mhapralkar, V.Y. (2007). Studies on unfermented beverages of jamun (*Syzygium cumini*) and Karonda (*Carissa carandus*L.) fruits. M.Sc. (Agri.) thesis submitted to Konkan Krishi Vidyapeeth, Dapoli, and Dist. Ratnagiri (M.S.).
7. Nainegali, B.S., 2021. Studies on Selective Extraction and Purification of Bioactive Compounds from Kokum (*Garcinia Indica*) Fruits using Alcohol Based Aqueous Two-

Phase Systems (Doctoral dissertation, National Institute of Technology Karnataka, Surathkal).

8. Nayak, C.A., Srinivas, P. and Rastogi, N.K. (2010). Characterization of anthocyanin from *Garcinia indica choisy*. *Food Chemistry*. 118: 719-724 Nutrition, Income and Sustainable Development 2008;806:539-546, 22.
9. Oladipo, I.C., Adeleke, D.T. and Adebisi, A.O.(2010). The effect of pH and chemical preservatives on the growth of bacterial isolates from some Nigerian packaged fruit juices. *Pakistan journal of biological sciences: 13*(1), pp.16-21.
10. Padhye, S., Ahmad, A., Oswal, N. and Sarkar, F.H. (2009). Emerging role of Garcinol, the antioxidant chalcone from *Garcinia indica Choisy* and its synthetic analogs. *Journal of hematology & oncology*, 2(1):1-13.
11. Palaniswamy, K.P. and Muthukrishnan, C.R. (1974). Studies on the physico-chemical characters of lemon juices and squashes during storage. *Indian Food Packer*, 28(4): 37-51.
12. Ranganna, S.(1986). Handbook of analysis and quality control for fruits and vegetables.
13. Rasha, H.M., Salha, A., Thanai, A. and Zahar, A., (2015). The biological importance of *Garcinia cambogia*: A review. *Journal of Nutrition and Food Sciences*, 5(4):1-2.
14. Rhim, J. W., (2002). Kinetics of thermal degradation of anthocyanin pigment solutions driven from red flower cabbage. *Food. Sci. Biotech.*, 11: 361-364
15. Shahnawaz, M., Sheikh, S. A. and Minhas, S. (2013) Role of sodium benzoate as a chemical preservative in extending the shelf life of orange juice. *Global Adv Res. J. Food Sci. Technol.* 2(1):7-18.
16. Shakoor, W., Javid Ullah, Alam Zeb and Muhammad Z. (2013). Effect of Refrigeration Temperature, Sugar Concentrations and Different Chemicals Preservatives on the Storage Stability of Strawberry Juice. *International Journal of Engineering and Technology IJET-IJENS.*, 13:02

17. Shrestha, M.K. and Bhatia, B.S. (1982). Apple juice physico-chemical characteristics and storage study. *Indian Food Packer*, 35(3): 53-60.
18. Shrivastava P, Kumar R. 2015. Soil salinity: A serious environmental issue and plant growthpromoting bacteria as one of the tools for its alleviation. *Saudi Journal of Biological Sciences*22:123-131.
19. Subash Chandran, M.D. (2005). Conservation and sustainable use of cultivated and wild tropical fruitdiversity: Introduction to good practices. Good Practices Workshop of UNEP-GEF funded project “Conservation and Sustainable Use of Cultivated and Wild Tropical Fruit Diversity”. Bangkok.Thailand, 4-5 Apr. 2005
20. Tasnim, F. Jr, Hossain, M.A., Nusrath, S., Hossain, M.K., Lopa, D. and Formuzul Haque,
Tata Mc.Graw Hill Publishing Company Limited. New Delhi-110, 7,:9-10.
21. Birnale KS, Pujari KH, Kudale VB. To study the changes in chemical and microbial quality parameters of the blended pineapple (*Ananas comosus* L.) and nutmeg (*Myristica fragrans* L.) rind jelly cubes during storage after product development. *Methods*. 2022;4:90.