

## Original Research Article

### **Enhanced L-lysine Production by UV-irradiated and S-2-aminoethyl-L-cysteine resistant mutants derived from *Bacillus* species using agricultural products as carbon and nitrogen sources**

#### **Abstract**

L-Lysine is essential for human and animal nutrition and may be added to food and feed materials to improve the protein quality. It is utilized in human medicine, in cosmetics and in the pharmaceutical industry, particularly in the formulation of diets with balanced amino acid concentration and in amino acid infusions. Enhanced L-lysine Production by UV irradiated and S-2-aminoethyl-L-cysteine resistant (AEC<sup>R</sup>) mutants derived from *Bacillus* species, using Agricultural Products as carbon and nitrogen sources was studied. The L-lysine producing bacteria had already been isolated from Nigerian soil. They were purified and Identified as *B. subtilis* PR13, *B. subtilis* PR9 and *B. pumilus* SS16, using cultural, biochemical and molecular characteristics. UV irradiated mutants were obtained by exposing the parent strains (wild types) which include *B. subtilis* PR13, *B. subtilis* PR9 and *B. pumilus* SS16, to UV light in a cabinet fitted with UV lamp with a wavelength of 253.7nm. AEC<sup>R</sup> mutants were derived by exposing the parent strains to various concentrations of AEC. The mutants were screened for L-lysine production by cultivating fermentation media. The results obtained showed that, UV mutants PRMU-8 and PRMU-14 derived from *Bacillus subtilis* PR13 accumulated higher lysine yield of 2.47 and 2.55 mg/ml compared to the wild type. Also, UV mutant MUPR-4 derived from *Bacillus subtilis* PR9 recorded a higher lysine yield of 1.77 mg/ml, while enhanced lysine yield of 1.96 and 2.34 mg/ml was observed in UV mutants SSMU-4 and SSMU-9 respectively. AEC<sup>R</sup> mutant PRT-2 derived from *Bacillus*

*subtilis* PR13, accumulated higher lysine yield of 2.35 mg/ml compared to the wild type. Enhanced lysine production of 1.47 and 1.32 mg/ml was observed in mutants ART-5 and ART-8 derived from *Bacillus subtilis* PR9 compared to the wild type. Mutant SRT-4 derived from *Bacillus pumilus* SS16 recorded enhanced lysine accumulation of 2.18 mg/ml compared to the wild type. The results obtained in the study showed that L-lysine production by some UV irradiated and AEC<sup>R</sup> mutants was enhanced.

**Keywords:** *Bacillus* species, L-lysine, UV irradiation, AEC, mutants

## 1. INTRODUCTION

Amino acids are major industrial products derived by fermentation, covering a world market of more than 5 million tons per year. As the building blocks of life, amino acids have long played an important role in both human and animal nutrition and health maintenance (Bercovici and Fuller 1995). Among amino acids are the L-lysine that is one of the leading biotechnological products and the global L-lysine production is estimated to reach 3.0 million tons in 2022 corresponding to 5.6 billion USD of market value according to the current L-lysine market report ([Elder, 2018](#))

with a current production of 2.2 tons per year [Eggeling and Bott M (2015)]. L-Lysine is essential for human and animal nutrition. Lysine cannot be synthesized biologically in the body and its breakdown is irreversible [Shah et al 2002], but may be added to food and feed materials to improve the protein quality [Stilling et al 1971].

Children and growing animals have a high requirement of lysine, since it is needed for bone formation.(Abdul Haleem Shah et al 2002)

Lysine can be produced in different ways including chemical synthesis, extracting from protein hydrolyzate, enzymatic method, fermentation method, protoplast fusion technique and recombinant DNA technology [Anastassiadis S. (2007), Nelofer et al (2008)]. Among these methods, fermentation is the

most economical and practical means of producing lysine, as in this method low temperature, low pressure and low-cost carbon sources are used and a biological form of lysine (L-lysine) is produced [Ekwealor, A. and Obeta AN (2005)].

L-lysine commercial form requires different downstream processing to achieve the degree of purity (implying significant differences in investment costs, losses during down streaming, amount of waste volume, and user friendliness), so that L-lysine-HCl involves higher operating costs due to the larger number of equipment and energy consumption. L-Lysine is being produced on industrial scale using *Corynebacterium glutamicum*, species of *Arthrobacter*, *Bacillus* and *Brevibacterium* as fermenting agent [Sen SK, Chatterjee SP 1989, Kircher M, Pfeerle W 2001 ]. High yielding strains have also been developed from *Bacillus subtilis* and *Escherichia coli* [Leuchtenberger 1996].

L-lysine is useful as chemical agent, food material and feed additive” [Anastassiadis 2007]. In addition, it has pharmaceutical applications both in the formulation of diets with a balanced amino acid composition and in the infusion of amino acids. Lysine supports bone health by insuring adequate absorption of calcium and therefore prevents osteoporosis. It has an important role in production of antibodies for healthy immune system. It is the integral component of musculature .

Due to increasing market demand and price competition, extensive research has been made in order to improve the fermentation process not only from the point of lowering production costs but also of increasing productivity. The lysine industry has shown consistency in developing efficient and better microbial strains either through classical mutagenesis or by applying modern biotechnology tools. These approaches are expected to improve the economics of fermentation process (Costa Ferreira & Duarte, 1992).

A great variety of microorganisms, including auxotrophic as well as regulatory mutants, has been reported to over-produce lysine (Tosaka & Takinami, 1986). The selection of such mutants has led to isolation of high producers, which are used for industrial production of lysine, glutamic acid, threonine and a variety of other amino acids (Han et al., 1991).

We had isolated three *Bacillus* species (which included *Bacillus subtilis* PR13, *B. subtilis* PR9, and *B. pumilus* SS16) from Nigerian soil, which produced various yields of L-lysine [Okpalla and Ekwealor 2019]. In another study, the *Bacillus* species were used for L-lysine production using carbohydrates as

carbon and seed meals as nitrogen sources respectively [Okpalla and Ekwealor 2019].

The present study was aimed at enhancing L-lysine Production by UV irradiation and AEC derived mutants of *Bacillus* species using Agricultural products as carbon and nitrogen Sources.

## **2. MATERIALS AND METHODS**

### **2.1 Microorganisms**

*B. subtilis* PR13, *B. subtilis* PR9 and *B. pumilus* SS16 isolated from different soil [15] in Awka town, were used in the study. They were purified and Identified as *B. subtilis* PR13, *B. subtilis* PR9 and *B. pumilus* SS16 using cultural, biochemical and molecular characteristics. The *Bacillus* species were grown on nutrient agar slants for 24 h at 37 °C. Thereafter, the cultures were then preserved at 4°C and transferring to new slants after 30 days in order to keep them viable for use in L-lysine production.

### **2.2 Inoculum preparation**

Two loopfuls of *B. subtilis* PR13, *B. subtilis* PR9 and *B. pumilus* SS16 were inoculated in an Erlenmeyer flask containing sterile 50 ml of seed medium. The seed medium consisted of peptone, 10.0g; yeast extract, 10.0 g; NaCl, 5.0 g; water, 1litre; pH adjusted to 7.2. The inoculated flasks were incubated for 24 h on a rotary shaker at 120 rpm and 30 °C. Duplicate flasks were used.

### **2.3 Preparation of Fermentation Media**

The submerged production of L-lysine by *B. subtilis* PR13, *B. subtilis* PR9, was conducted in three different fermentation media, namely fermentation medium 1,2 and 3 (FM1, FM2 and FM3). For *Bacillus subtilis* PR 13, L-lysine production was carried out in 100 ml Erlenmeyer flasks, containing FM1. The medium, was composed of  $\text{KH}_2\text{PO}_4$ , 0.5g;  $\text{K}_2\text{HPO}_4$ , 0.5g;  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.001g;  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ , 0.001g;  $\text{FeSO}_4 \cdot 7\text{HO}$ , 0.001g;  $\text{CaCO}_3$ , 50g, the carbon source (glucose) was replaced with millet starch hydrolysate 60g; the nitrogen source (ammonium sulphate) was replaced by soyabean meal 40g; water, 1 litre; pH adjusted to 7.2. For *B. subtilis* PR 9 and *B. pumilus* SS16, the FM2 and FM3 media were used for L-lysine production respectively. FM2 and FM3

were similar to FM1, except that for FM2, the carbon source (glucose) was replaced with sorghum starch hydrolysates 60g, the nitrogen source (ammonium sulphate) was replaced by defatted peanut meal 40g, while for FM3 the carbon source (glucose) was replaced with sorghum hydrolysates 60g, the nitrogen source (ammonium sulphate) was replaced by defatted soyabean meal 20g. The carbon source substrates were prepared in the laboratory using the method of Umerie *et al.* [20].

#### **2.4 UV irradiation and mutant selection**

Two loopful of a 24h culture was used to inoculate a 250ml Erlenmeyer flask containing 50ml of nutrient broth medium. The flask was incubated for 24h on a rotary shaker at 120rpm and 30<sup>0</sup>C. One milliliter of the broth culture was suspended aseptically in 4ml of phosphate buffer(pH 7.2) contained in sterile glass petri dishes of 100mm diameter. The UV light exposure was carried out in a cabinet fitted with UV lamp (253.7nm). The exposure was carried out at a distance of 20 cm and the exposure times were 0, 2, 4, 6, 8,10 and 12 mins. Each UV exposed cell suspension was stored in dark for 10mins to avoid photo-reactivation (getting revertants), then 0.1ml of the irradiated cells was inoculated on Dextrose Peptone agar (DPA) plates for growth and colonies which appeared after 24 h incubation at 30<sup>0</sup>C were subsequently plated out on minimal agar medium. Mutant strains which showed no growth on the minimal medium were selected and stored on Nutrient agar slants at 4<sup>0</sup>C. The mutant strains were screened for lysine production.

#### **2.5 Screening of UV Irradiated Mutant Strains for Lysine Production**

Mutants derived after UV mutagenesis were screened for lysine accumulation. Fermentation was carried out in 100ml Erlenmeyer flasks containing 20 ml of fermentation media (FM1 and FM2) and 25 ml of fermentation 3 (FM3). The preparation of the media were as was previously described. Two milliliters volume of 24h cultures of *Bacillus* species were inoculated into the fermentation media. Uninoculated flasks served as control. The flasks were placed on a rotary shaker (at 160 rpm) and incubated at 30<sup>0</sup>C for 72 h.

Following the termination of fermentation, the fermentation medium were subjected to centrifugation at 5,000 rpm for 15 min to obtain the cell free supernatant which is the crude L- lysine. The cell free supernatant was used for the determination of lysine. The experiments were conducted in triplicate

## **2.6 Derivation of S-2-amino ethyl cysteine(AEC) resistant mutants**

The method of Tosaka *et al.*(1978) was used for the derivation of S-2-aminoethylcysteine mutants. Two loopful of 24h culture of *Bacillus* species was grown in an Erlenmeyer flasks containing 20ml of the seed medium(peptone,1%; yeast extract,1%; NaCl, 0.5%; pH,7.0).The flask was incubated at 30<sup>0</sup>C on a rotary shaker at 120rpm for 24h. A 5ml portion of the broth culture was centrifuged and the cells washed with 0.5ml of 0.1M sodium phosphate buffer (pH 7.0) before resuspending in equal volume of the buffer. A loopful of the cell suspension was inoculated on minimal agar plates containing various concentrations (0.5, 1.0, 1.5, 2.0 mg/ml) of S-2-aminoethylcysteine. S-2-aminoethylcysteine resistant(AEC<sup>R</sup>) colonies that appeared after 3 to 4 days of incubation at 30<sup>0</sup>C were screened for lysine production.

## **2.7 Screening of S-2-amino-ethyl-cysteine resistant mutant for lysine production**

Mutants were screened for lysine accumulation. Fermentation was carried out in 100ml Erlenmeyer flasks containing 20 ml of fermentation media (FM1 and FM2) and 25 ml of fermentation 3 (FM3). The preparation of the media were as was previously described. Two milliliters volume of 24h cultures of *Bacillus* species were inoculated into the fermentation media. Uninoculated flasks served as control. The flasks were placed on a rotary shaker (at 160 rpm) and incubated at 30<sup>0</sup>C for 72 h. Following the termination of fermentation, the fermentation medium were subjected to centrifugation at 5,000 rpm for 15 min to obtain the cell free supernatant which is the crude L- lysine. The cell free supernatant was

used for the determination of lysine. The experiments were conducted in triplicate

**Statistical analysis** Data generated from this work were analyzed using correlation analysis with a software application SPSS version 14

### 3. RESULTS

L-lysine production by UV irradiated mutants derived from *Bacillus subtilis* and *B. pumilus* is shown in Table 1. Four, one and three UV irradiated mutants were derived from the wild types of *B. subtilis* PR13, *B. subtilis* PR9 and *B. pumilus* SS16 respectively (Table 1). Mutants PRMU-8 and PRMU-14 derived from *Bacillus subtilis* PR13 accumulated higher lysine yield of 2.47 and 2.55 mg/ml compared to the wild type (used as control) which recorded 2.13 mg/ml. The mutants strains PRMU-2 and PRMU-15 recorded reduced lysine yield compared to the wild type. Mutant MUPR- 4 derived from *Bacillus subtilis* PR9 recorded a higher lysine yield of 1.77 mg/ml compared to the wild type which recorded 1.28 mg/ml. Enhanced lysine accumulation of 1.96 and 2.34 mg/ml was observed in mutants SSMU-4 and SSMU-9 derived from *Bacillus pumilus* SS16, compared to the wild type which recorded 1.75mg/ml. Mutant SSMU-11 recorded reduced lysine yield of 1.27mg/ml more than the wild type.

Table 2 showed the result of L- lysine production by S-2-Amino-ethyl-L-cysteine resistant (AEC<sup>R</sup>) mutants derived from *Bacillus subtilis* and *B. pumilus*. One, three and one AEC<sup>R</sup> mutants were derived from the wild types of *B. subtilis* PR13, *B. subtilis* PR9 and *B. pumilus* SS16 respectively. AEC<sup>R</sup> mutant PRT-2 derived from *Bacillus subtilis* PR13, accumulated higher lysine yield of 2.35 mg/ml compared to the wild type which recorded 2.13mg/ml. Enhanced lysine production of 1.47 and 1.32 mg/ml was observed in mutants ART-5 and ART-8 derived from *Bacillus subtilis* PR9 compared to the wild type which recorded 1.28mg/ml. Mutant SRT-4 derived from *Bacillus pumilus* SS16 recorded enhanced lysine accumulation of 2.18 mg/ml compared to the wild type which recorded 1.75 mg/ml.

**Table 1: L- lysine production by UV irradiated mutants derived from *Bacillus subtilis* and *B. pumilus***

| Wild type                        | UV irradiated mutant | Treatment time (min) | Lysine (mg/ml) |
|----------------------------------|----------------------|----------------------|----------------|
| <i>Bacillus subtilis</i><br>PR13 | PRMU – 2             | 10                   | 1.83           |
|                                  | PRMU – 8             | 10                   | 2.47           |
|                                  | PRMU – 14            | 12                   | 2.55           |
|                                  | PRMU – 15            | 12                   | 1.74           |
|                                  | Control              |                      | 2.13           |
| <i>Bacillus subtilis</i><br>PR9  | MUPR-4               | 8                    | 1.77           |
|                                  | Control              |                      | 1.28           |
| <i>Bacillus pumilus</i><br>SSI6  | SSMU-4               | 10                   | 1.96           |
|                                  | SSMU-9               | 10                   | 2.34           |
|                                  | SSMU-11              | 8                    | 1.27           |
|                                  | Control              |                      | 1.75           |

**Table 2: L- lysine production by S-2-Amino-ethyl-L-cysteine resistant (AEC<sup>R</sup>) mutants derived from *Bacillus subtilis* and *B. pumilus***

| Wild type                         | AEC <sup>R</sup> mutant | Lysine (mg/ml) |
|-----------------------------------|-------------------------|----------------|
| <i>Bacillus subtilis</i><br>PR 13 | PRT- 2                  | 2.35           |
|                                   | Control(wild type)      | 2.13           |
| <i>Bacillus subtilis</i><br>PR9   | ART-5                   | 1.47           |
|                                   | ART-7                   | 0.91           |
|                                   | ART-8                   | 1.32           |
|                                   | Control                 | 1.28           |
| <i>Bacillus pumilus</i><br>SS16   | SRT-4                   | 2.18           |
|                                   | Control                 | 1.75           |

#### 4. DISCUSSION

Mutant strains of the organisms derived after UV light treatment were found to produce lysine in higher yield than the wild type (parent strain). This is supported by the findings of other researchers. Shah *et al.*(2002) reported an improved lysine production from mutants developed after UV treatment of *Corynebacterium glutamicum*, with the most potent mutants producing 38, 33 and 28.5g/l lysine, in the media containing glucose, molasses and starch hydrolysate. Ekwealor and Obeta(2006) in a study of the effects of UV irradiation on lysine production by *Bacillus megaterium*, observed that mutant MR-10, and MR-25 derived from *Bacillus megaterium* SP 76 and MS-3 and MS-5 derived from *Bacillus megaterium* SP 86 produced higher lysine levels than the parent strains. Sharma(2000) reported a methionine concentration of 4g/l using UV mutants.

Strain improvement by classical mutagenesis techniques is well established and widely used to isolate amino acid over producers (Rowland, 1984; Sharma, 2002).The mutation approach has become the most extensively used tool for industrial organisms (Brautaset *et al.*, 2003, Lee *et al.*, 1996). Mutagenesis is a phenomenon by which changes can be introduced in the metabolic process. The most direct and general method for over production is the genetic removal of the feed back control (Shah *et al.*, 2002). Different mutagens can be used for the desirable changes in the genetics of the strains of interest (Jacobson, 1981). UV light has been recommended as mutagen of the first choice. The ratio of mutation is usually very high and UV light is a relatively safe mutagen for the experimenters (Bridge, 1976).

The result from the study showed increased lysine production from some AEC<sup>R</sup> mutants. The findings of other researchers supported this view. Ekwealor and Obeta (2005) derived AEC resistant mutants from *B. megaterium* SP14 and *B. megaterium* sp76 and were found to enhance lysine accumulation. Odunfa *et al.*(2001) reported the lysine production by AEC resistant *Lactobacillus* and yeast isolates. The result showed that 42.5% of *Lactobacillus* and 83.3% of the yeast isolates tested were capable of lysine production. Siripoke *et al.*(2005) reported the selection of AEC<sup>R</sup> *Bacillus* SWU41, a *Bacillus* resistant to a toxic lysine analogue (AEC), which produced 3.68g/l of lysine (about 6 times of the prototroph) in the medium containing 5% glucose. Nadeem *et al.*(2001) observed lysine production of 0.1-0.5g/l by mutant *Escherichia coli* which showed resistance against AEC a lysine analogue. Though they noted that resistance to AEC does not necessarily mean the overproduction of lysine as seen in their results. Sano and Shiio (1970) developed AEC resistant mutant of *Bacillus subtilis*, *Brevibacterium flavum* and *Escherichia coli*. Among them, *B. flavum* mutant resistant to the growth inhibition of AEC plus threonine was the best, producing 32g L-lysine for 100g glucose. Hilliger and Prauser (1989) reported that AEC resistant mutants of *Oerskonina* accumulated up to 10g/l of lysine. Yakoto and Shiio (1988) described 40g/l of L-lysine production in 10% glucose medium by threonine negative mutant of AEC resistant *Brevibacterium flavum*. Adelberg (1958) noted that some mutants resistant to amino acid analogues are suitable as amino acid producers. The enhanced lysine production by AEC resistant mutants is likely due to the mechanism of control operative in the biosynthesis of lysine in the parent strains being interfered with in the mutants, thus making them over producers. This view is supported by the findings of Samanta and Bhattacharyya (1991), Crociani *et al.* (1991) and Bhattacharyya and Samanta (1992). These researchers working with *Arthrobacter globiformis* and *Bacillus stearothermophilus*, noted the improved lysine yields of AEC resistant mutants of these organisms. Analogue-resistant-mutant isolation is developed for the purpose of overproduction of amino acid

and it has yielded higher production of arginine, lysine, tryptophan threonine etc utilizing different bacterial strains (Aida, 1986). Amino acid analogues can effectively function as true feed back inhibitors without participating in functions in the cell (Morinaga *et al.*, 1982; Chattopadhyays *et al.*, 1995). Mutants resistant to lysine analogues have altered and deregulated enzymes that are not sensitive to feedback inhibition and repression. Liu (1986) however suggested that apart from feed back inhibition and side reaction metabolic interlocks regulate the synthesis of amino acid. He opined that by the release of this regulation the productivity of amino acid increases.

## 5. CONCLUSION

In the study, it was observed that some UV irradiated mutants (from *Bacillus subtilis* PR13) stimulated enhanced lysine production, with mutant PRMU-8 accumulating the highest lysine yield of 2.47 mg/ml. Also, some of the AEC<sup>R</sup> mutants (from *Bacillus subtilis* PR 13) produced an enhanced yield of L-lysine, with mutant PRT- 2 accumulating the highest yield of 2.35mg/ml . The study showed that UV irradiated and AEC<sup>R</sup> mutants can produce enhanced lysine yield. This development indicates that large scale L-lysine production is feasible in Nigeria and it will help to meet present-day needs in its industrial sector.

## References

- [1] Bercovici D, Fuller F. Industrial amino acids in nonruminant animal nutrition. In: Wallace RJ, Chesson A (eds) Biotechnology in animal feeds and animal feeding, VCH, Weinheim, 1995; pp 93–113
- [2] Elder, M. World markets for fermentation ingredients, BCC Research: Market Research Reports. FOD020E. Available at: <https://www.bccresearch.com/market-research/food-and-beverage/world-markets-for-fermentation-ingredients-fod020e.html>, 2018
- [3] Eggeling L, Bott M. A giant market and a powerful metabolism: L-lysine provided by *Corynebacterium glutamicum*. *Appl Microbiol Biotechnol*, 2015; 99(8):3387–3394. doi:10.1007/s00253-015-6508-2
- [4] Shah AH, Hameed A, Khan GM. Fermentative Production of L-Lysine: Bacterial Fermentation-I. *J Med Sci*, 2002; 2: 152-157.

- [5] Stilling BR, Sidwell VD, Hammerle OA. Nutritive Quality of Wheat Flour and Bread Supplement with Either Fish Protein Concentrate or Lysine. *Cereal Chem*, 1971; 292-302.
- [6] Anastassiadis S. L-lysine fermentation. *Recent Patents Biotechnol*. 2007; 1: 11-24.
- [7] Nelofer R, Syed Q, Nadeem M. Statistical Optimization of Process Variables for L-lysine Production by *Corynebacterium glutamicum*. *Turk. J. Biochem*. 2008; 33 (2): 50–57.
- [8] Ekwealor A, Obeta AN. Studies on lysine production by *Bacillus megaterium*. *African J. Biotechnol*. 2005; 4: 633-638.
- [9] Sen SK, Chatterjee SP. Influence of B-vitamins and trace elements on lysine production by *Micrococcus varians* 2fa, *Acta Biotechnol*. 1989; 9:63-67.
- [10] Kircher M, Pfeerle W. The fermentative production of L-lysine as an animal feed additive. *Chemosphere*. 2001; 43:27-31.
- [11] Leuchtenberger W. Amino acids – technical production and uses. In Rehm HJ, Reed G, Phuler A, Stadler P, editors. *Products of primary metabolism. Biotechnology vol. 6.* Wiley- VCH, Weinheim, German. 1996; 492.
- [12] Costa-Ferreira M, Duarte JC. Amino acid accumulation by an analogue– sensitive mutant of *Corynebacterium glutamicum*. *Biotechnol Lett*, 1992; 14: 1025–8.
- [13] Tosaka O, Takinami K. Lysine. In: K. Aida, I. Chibata, K. Nakayama, K. Takinami and H. Yamada (Eds.), *Biotechnology of Amino Acid Production*, Vol. 24, 1986; pp: 152–72. Kodansha Ltd., Tokyo.
- [14] Han JK, Oh JW, Lee H, Chung S, Hyun HH, Lee LH. 1991. Molecular cloning and expression of S-( $\beta$ -aminoethyl)-L-cysteine resistant aspartokinase gene of *Corynebacterium glutamicum*. *Biotechnol, Lett*, 1991; 13: 721–6.
- [15] Okpalla J, Ekwealor IA. Screening for lysine production by bacteria isolated from Nigerian soil. *World wide J Multidiscip Res Develop*, 2019a; 5(4): 10-18.
- [16] Okpalla J, Ekwealor IA. Studies on lysine accumulation in broth culture of *Bacillus* species using carbohydrates as carbon sources and seed meals as nitrogen sources. *Inter J Trend in Scientific Res Development*, 2019b; 3(2): 760- 772.
- [17] Umerie SC, Ekwealor IA, Nwagbo, I.O. Lysine production of *Bacillus laterosporus* from various carbohydrates and seed meals. *Biores Technol* , 2007; 5: 249-252

- [18] Tosaka O, Takinami K, Hirose Y. L-Lysine production by S-2-amino-ethyl-cysteine and  $\alpha$ -amino- $\beta$ -hydroxyvaleric resistant mutants of *Brevibacterium lactofermentum*. *Agri Biol Chem*, 1978; 42: 745-752
- [19] Ekwealor IA, Obeta, JAN. Screening of UV-irradiated and S-2-amino-ethyl-cysteine resistant mutants of *Bacillus megaterium* for improved lysine accumulation. *African Journal of Biotechnology*, 2006; 5(22): 2312-2314
- [20] Sharma S. Strain improvement for the production of methionine. Ph.D. thesis. Indian Institute of Technology, Delhi, India, 2000.
- [21] Rowlands RT. Industrial strain improvement: mutagenesis and random screening procedures. *Enzymes in Microbiology and Technology*, 1984; 6: 3-10.
- [22] Brautaset T, Williams MD, Dillingham DR, Kaufmann C, Bennaars A, Crabbe E, Flickinger EC. Role of the *Bacillus methanolicus* citrate synthase 11 Gene, city, in regulating the secretion of glutamate in L-lysine-secreting mutants. *Applied Environmental Microbiology*, 2003; 69: 3986-3992.
- [23] Lee, G.H., Hur, W., Bremmon, C.E. and Flickinger, M.C. (1996). Lysine production from methanol at 50<sup>0</sup>C using *Bacillus methanolicus*: modeling volume control, lysine concentration and productivity using a three-phase continuous simulation. *Biotechnology and Bioengineering*, 49: 639-653.
- [24] Jacobson, G.K. (1981). Mutation. In: *Microbial fundamental* (Vol.1). Rahman, H.J. and Reed, G. (eds) AVI publishing Co. New York pp281-299.
- [25] Bridge, B.A. (1976). Genetics. In: *second intern. Symposium of industrial microorganism*, Donald, K.D. ed, Academic press, New York pp 7-14.
- [26] Odunfa SA, Adeniran SA, Teniola OD, Nordstorm, J. (2001). Evaluation of lysine and methionine production in some lactobacilli and yeasts from Ogi. *International Journal of Food Microbiology*, 63: 159-63.
- [27] Siripoke S, Potivejkul K, Lerstaveesin, P, Leungsakul, S. Optimal conditions for lysine production by *Bacillus* SWU 41 in a Lab-Scale fermenter. *31<sup>st</sup> Congress on Science and Technology of Thailand at Suranaree University of Technology*. 2005.
- [28] Nadeem S, Ikram A, Rana S.M, Yaqoob NY, Qureshi MJ, Shakoori AR. Enhanced L-lysine production by an *Escherichia coli* mutant WARN 30522 after MNNG treatment. *International Journal of Agricultural Biology*, 2001; 3(4): 447-450.

- [29] Sano K, Shiio I. Microbial production of L-lysine by mutants resistant to S-(2-amino ethyl)-L-cysteine. *General Applied Microbiology*, 1970; **16**: 373-391.
- [30] Hilliger M, Prauser H. (1989). L-Lysine formation in the Nocardioform Taxon. *Oerskovia. Folia Microbiology* 1989; **34**: 427-428.
- [31] Yakota A, Shiio I. Effect of reduced citrate synthetase activity and feed back resistance phosphoenol pyruvate carboxylase on lysine production. *Brevibacterium flavum* mutant. *Agriculture and Biological Chemistry*, 1988; **52**: 455-463.
- [32] Adelberg EA. Selection of bacterial mutants which excrete antagonists of antimetabolites. *J Bacteriol*, 1958; **76**: 326-328.
- [33] Samanta TK, Bhattacharyya S. L-Lysine production by S-2-amino-ethyl-cysteine, Resistant mutants of *Arthrobacter globiformis*. *Folia Microbiol*, 1991; **36**: 59-66
- [34] Crociani F, Selli A, Criseting G, Glosa DG, Matteuzzi D. L-Lysine production at 65°C by auxotrophic-regulatory mutants of *Bacillus stearothermophilus*. *J Industrial Microbiol*, 1991; **8**:127-132.
- [35] Bhattacharyya R, Samanta TK. L-lysine production by double auxotrophic and AEC resistant mutants of *Arthrobacter globiformis*. *Res in Industry*, 1992; **37**: 1-6.
- [36] Aida K. An overview of the microbial production of amino acids. In: K. Aida, I. Chibata, K. Nakayama, K. Takinami and H. Yamada (Eds). *Biotechnology of Amino Acid Production* vol 24. Kodansha Ltd, Tokyo. 1986.
- [37] Morinaga Y, Tani Y, Yamada H. L-methionine production by ethionine resistance mutant of facultative methylotroph, *Pseudomonas* FM 518. *Agri Biol Chem* **46**: 473-80.
- [38] Chattopadhyay MK, Ghosh AK, Sengupta S, Sengupta D. Threonine analogue resistant mutants of *Escherichia coli* K- 12. *Biotech Lett*, 1995; **17**: 567-70.
- [39] Liu YT. Studies on the fermentation production of L-lysine. Optimization of culture condition for L-lysine fermentation with cane molasses. *Rep Taiwan and sugar Res Instit* 1986; **114**: 45-66.