

Original Research Article

Antimicrobial Synergism of Blended Essential Oils from *Piper betle* (L.) and *Melaleuca alternifolia*

ABSTRACT

Aims: The synergistic antibacterial effects of blended essential oils from *Piper betle* and *Melaleuca alternifolia* against nine different microbial strains were investigated.

Methodology: GC-MS was employed for the analysis of essential oil extracted by hydro-distillation. The synergistic antibacterial effects of blended essential oils from *Piper betle* and *Melaleuca alternifolia* against nine different microbial strains were assayed using standard methods including disc diffusion, agar dilution and checkerboard.

Results: Essential oil from *Piper betle* consisted mainly of eugenol (39.21%) followed by other components: eugenol acetate (16.42%), 4-allyl-1,2-diacetoxybenzene (12.24%), terpinen-4-ol (6.58%), α -cadinol (6.13%), γ -terpinene (3.46%), and sabinene (2.14%). Meanwhile, the major components of tea tree essential oil were terpinen-4-ol (49.62%), followed by other components: γ -terpinene (18.08%), α -terpinene (9.16%), *p*-cymene (5.89%), α -terpineol (4.94%), terpinolene (3.47%), and α -pinene (2.02%). The synergistic antibacterial effects of blended essential oils from *Piper betle* and *Melaleuca alternifolia* were recorded in seven of the nine bacteria tested, including *Bacillus subtilis* (FICI = 0.250%), *Enterococcus faecalis* (FICI = 0.375%), *Staphylococcus aureus* (FICI = 0.313%), Methicillin-resistant *Staphylococcus aureus* (FICI = 0.281%), *Acinetobacter baumannii* (FICI = 0.375%), *Escherichia coli* (FICI = 0.375%) and *Salmonella sp* (FICI = 0.375%). However, additive effects were only observed in both Gram-negative bacteria including *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* when treated with blended essential oils resulting in FICI value of 0.625% and 0.750%, respectively.

Conclusion: The combination of essential oils from *Piper betle* and *Melaleuca alternifolia* exhibited more significant antibacterial effects against nine studied microbial strains than single essential oil. Generally, Gram-negative bacteria were partly more resistant to blended essential oils from *Piper betle* and *Melaleuca alternifolia* than Gram-positive strains.

Comment [H1]: Topic is misleading since the antifungal effects of the constituents were not investigated. You may modify to read; Antibacterial synergism

Comment [H2]: Rearrange: To investigate the antibacterial synergistic effect of blended

Comment [H3]: Rearrange: The chemical constituents of the essential oils extracts of the plant were identified using GC-MS technique, while the antibacterial activity of the oils was investigated using disc diffusion assay.

Comment [H4]: How many compounds present in each oils; For example, The GC-MS analysis indicated the presence of 4, 12 chemical compounds for the *Piper betle* and *Melaleuca alternifolia*, respectively

Comment [H5]: Part of the discussion delete

Accordingly, the synergistic antibacterial effects were recorded in seven of the nine microbial strains tested, whereas the additive effects were only observed in both Gram-negative bacteria.

Keywords: Piper betle, Melaleuca alternifolia, GC-MS, synergism, antimicrobial assays.

1. INTRODUCTION

Piper betle L. (*P. betle*), commonly known as betel vine, belongs to genus *Piper* of the Piperaceae family. Betel vine has been traditionally used as herbal medicine mainly as an antiseptic in dental practice. Betel plants, locally known as Pann (India), Ikmo (Philippines) or Trầu không (Vietnam), are popularly cultivated for their leaves in India, Sri Lanka, Thailand and other Asian countries. Betel leaves, the most commonly used plant parts, are widely chewed as not merely a habit practice and a custom, but also an identity of the traditional cultural values which are known as a symbol of love and happiness. In addition, regular chewing of a mixture of betel leaf, areca nut and lime is believed beneficial to health and well-being in several ways such as preventing halitosis, strengthening the gum, protecting the teeth and even improving the digestive health. Phytochemical analysis of betel leaves from different sources of botanical origin revealed the presence of alkaloids, glycosides, saponins, steroids, tannins, flavonoids, volatile oils, proteins and carbohydrates [1]. Commercial betel leaf-based formulations are now available over the counter in pharmacies or drug stores such as mouthwash, toothpaste, shampoo, soap, hand sanitizer or lady care. Betel leaf essential oil (BLO) is a complex chemical mixture consisting of approximately between 20 to 60 natural volatile compounds. These compounds mainly belong to classes of terpenes and phenols in various proportions depending on the botanical origin of the betel leaves. Major components of BLO include eugenol, chavicol, chavibetol, hydroxychavibetol and safrole. Other chemical components of BLO are anethole, estragole, linalool, caryophyllene, *p*-cymene, eucalyptol (1,8-cineole), estragol, sabinene, α -cadinol, γ -terpinene, γ -muurolene, α -muurolene, β -phellandrene, germacrene-D, 4-allyl-1,2-diacetoxybenzene [2-9]. BLO possesses a variety of significant pharmacological properties including antiseptic, antioxidant, antiprotozoal, antifungal, antibacterial and radio-protective properties. Nevertheless, as evidenced from previous studies, the antiseptic property of BLO is attributed chiefly to the major constituents of BLO including chavicol, chavibetol and eugenol. In addition, previous researches showed that BLO exhibits a broad spectrum of antimicrobial activity against pathogenic microorganisms including Gram-positive and Gram-negative bacteria, and fungi such as *Aspergillus*, *Candida*, *Epidermophyton*, *Microsporum*, *Trichophyton*, *Bacillus*, *Escherichia*, *Klebsiella*, *Listeria*, *Pseudomonas*, *Salmonella*, *Staphylococcus* or *Streptococcus* [3-9].

Comment [H6]: Citaion, need references

Comment [H7]: 17

Comment [H8]: 17

Comment [H9]: 17

Comment [H10]: 17

Melaleuca alternifolia (*M. alternifolia*), commonly known as tea tree, is a member of the family Myrtaceae native to Australia. Tea tree essential oil (TTO), a mixture of natural volatile compounds extracted mainly from tea tree leaves, has been widely used as an alternative and complementary therapy for a variety of skin conditions such as acne, wounds or contact dermatitis [10,11]. TTO possesses various significant biological activities including antibacterial, antiviral, antifungal and anti-inflammatory properties. TTO is composed of approximately 100 various chemical compounds. Chemical analysis of TTO reveals the presence of terpinen-4-ol, *p*-cymene, γ -terpinene, α -terpineol, terpinolene, α -pinene, α -terpinene and 1,8-cineole. Other chemical components of BLO are sabinene, limonene, aromandendrene, β -thujene, β -myrcene, β -pinene [11-15]. As indicated from previous literature, the antibacterial and antifungal properties of TTO are attributed mainly to terpinen-4-ol and α -terpineol; however, α -terpineol often contributes an insignificant proportion of TTO which is commonly accounted to form approximately between 2 and 5% of the total composition of essential oil. In addition, several previous scientific studies showed that TTO exhibits a wide spectrum of antimicrobial activity against pathogenic microorganisms including Gram-positive and Gram-negative bacteria, viruses, protozoa and fungi such as *Aspergillus*, *Candida*, *Bacillus*, *Enterococcus*, *Escherichia*, *Klebsiella*, *Haemophilus*, *Mycobacterium*, *Probionibacterium*, *Salmonella*, *Staphylococcus*, *Streptococcus*, Herpes simplex zoster (HSV), *Leishmania* or *Trypanosoma* [10,11,15,16]. Nowadays, TTO is commonly incorporated as an active ingredient of many topical formulations for treatment of skin infectious diseases. TTO-based cosmetics and skin care products are now widely available over the counter in pharmacies or chemists shops.

Comment [H11]: 17

Plants are always regarded as potential natural sources for new drug development. Since drug-resistant bacteria have become a global concern to public health in recent years, the discovery of new antibiotics has always attracted considerable interest from researchers working in several different disciplines. Potential antibacterial agents from plants can be total extracts, fractions, pure secondary metabolites or essential oils. Contemporary data clearly indicated that approximately 300 essential oils have been commercially used in several different fields ranging from folk medicine, aromatherapy to cosmetics, food preservatives and pharmaceutical industries. The antibacterial properties of essential oils have been traditionally and scientifically known for a long time, and since essential oils are a multi-constituent mixture with complex mechanisms of action, the combination of essential oils from different medicinal plants could be one of the potential strategies to enhance the antibacterial effects of blended essential oils against antibiotic-resistant bacteria. In this

Comment [H12]: Give examples

Comment [H13]: Talk directly about the involvement of medicinal plants in antibacterial effects

Comment [H14]: Should be the first paragraph

Comment [H15]: 17

regard, this study was primarily focused on investigating the potential antibacterial synergism of blended essential oils from *P. betle* and *M. alternifolia* against nine different microbial strains using standard microbiological assays including disc diffusion, agar dilution and checkerboard.

2. MATERIALS AND METHODS

2.1 Plant Materials and Essential Oil Extraction

Fresh *P. betle* leaves were collected from a local garden and washed several times with running water to remove any impurities. The plant leaves were immediately air-dried and cut into small pieces. The extraction of essential oil was carried out by hydro-distillation for 3 h using a Clevenger-type apparatus [17]. Meanwhile, fresh *M. alternifolia* leaves and young twigs were harvested from a farm in Mekong Delta and washed thoroughly with running water to eliminate any possible contaminants. The plant materials were then air-dried and cut into small pieces prior to the extraction of essential oil. The plant materials were also subjected to hydro-distillation for 2 h. Once the distillation was complete, the obtained essential oil was dehydrated using anhydrous sodium sulphate and then stored in sealed opaque brown bottles at 4°C until GC-MS analysis.

Comment [H16]: Where you identify the plant, Name of the botanist?

2.2 GC-MS Analysis

Gas chromatography-Mass spectroscopy (GC-MS) was employed for the analysis of obtained essential oil. GC-MS analyses were performed using SCION SQ 456-GC equipped with a Rxi-5ms RESTEX column (30 m x 0.25 mm x 0.25 µm). Helium was used as carrier gas at a constant flow rate of 1 mL/min. The oven temperature was initially programmed at 50°C for 1 min and then increased to 80°C at 30°C/min. After that, it was increased to 230°C at 5°C/min, and finally to 280°C at 25°C/min where it was held for 3 min. The injector temperature was set at 250°C and the rate of Division was 1:30 [18]. Fragmentation was done by electron impact under a field of 70eV. The mass spectra were recorded over the mass range of 50-500 amu with the full-scale mode at a rate of 1s/scan.

Comment [H17]: How you prepare the samples for GC/MS analysis, single, blended, Concentrations,

Comment [H18]: Remove, avoid repetition

2.3 Physicochemical Analysis

The determination of physicochemical parameters of essential oil for specific gravity, acid value, ester value and saponification value was performed as per standard test methods [19]. The obtained essential oils were also subjected to organoleptic evaluation including appearance, color, odor and taste.

Comment [H19]: Indicate the analysis is it for single oil or blended oil

2.4 Microbial Strains

The synergistic antibacterial effectiveness of blended essential oils from *P. betle* and *M. alternifolia* were tested against nine different microbial strains. Gram-positive species were *Bacillus subtilis* (*B. subtilis*), *Enterococcus faecalis* (*E. faecalis*), *Staphylococcus aureus* (*S. aureus*) and Methicillin-resistant *Staphylococcus aureus* (MRSA). Gram-negative strains included *Acinetobacter baumannii* (*A. baumannii*), *Escherichia coli* (*E. coli*), *Klebsiella pneumoniae* (*K. pneumoniae*), *Pseudomonas aeruginosa*, (*P. aeruginosa*) and *Salmonella* sp. The identity of the microorganisms assayed in this research was confirmed by morphological studies and standard biochemical tests [20].

2.5 Microbial Assay

2.5.1 AST

Antimicrobial susceptibility testing (AST) was conducted to investigate the sensitivity of a microorganism to essential oil using disc diffusion technique on agar media. The test was carried out by applying a standardized microbial inoculum of approximately 10^6 CFU/mL to the surface of Mueller-Hinton agar (MHA) plate. Agar plates were then incubated for 24 h at $35-37^{\circ}\text{C}$ prior to determination of results. The AST activity was assessed by measuring the growth inhibition zone (in millimetres, mm). The results were reported qualitatively as susceptible (++) , intermediate (+) or resistant (-) [21-25].

Comment [H20]: Write full name: Antimicrobial susceptibility testing

Comment [H21]: How you prepare the test solutions of the oils,? Concentration, number of samples

Comment [H22]: Positive control must used

2.5.2 MIC Assay

Agar dilution method was employed for the determination of minimum inhibitory concentration (MIC). Two-fold serial dilutions of essential oil were prepared by dilution with dimethyl sulfoxide (DMSO) to produce a series of decreasing concentration from 8% to 0.03125%. MHA plates were inoculated with a standardized inoculum of microbial according to McFarland standard. The plates were then incubated at $35-37^{\circ}\text{C}$ for 24 h. MIC value was defined as the lowest concentration of essential oil that completely inhibited visible growth of the tested microorganisms [26-29].

Comment [H23]: 21

Comment [H24]: How you prepare test solutions, concentration, how many samples

2.5.3 FIC Assay

The antimicrobial synergism of blended essential oils from *P. betle* and *M. alternifolia* against nine selected microbial strains were evaluated by the checkerboard method followed by the calculation of fractional inhibitory concentration index (FICI) using the formulas as described below:

FIC of BLO = MIC of BLO in combination/MIC of BLO alone

FIC of TTO = MIC of TTO in combination/MIC of TTO alone

Comment [H25]: 21

$$FICI = FIC \text{ of BLO} + FIC \text{ of TTO}$$

The interaction was categorized as follows: Synergism as $FICI \leq 0.5$, additive effect as $0.5 < FICI \leq 1$, indifference as $1 < FICI \leq 2$, and $FICI > 2$ was considered to be antagonism [30-32].

2.6 Statistical Analysis

All experiments were conducted in triplicate, and the results were expressed in terms of Mean \pm Standard Error of Mean (SEM). Statistical analysis was performed by SPSS and analysis of variance (ANOVA) with the level of significance $p < 0.05$.

3. RESULTS AND DISCUSSION

3.1 Organoleptic and Physicochemical Properties

Table 1: Organoleptic and physicochemical properties of essential oils from *Piper betle* and *Melaleuca alternifolia*

Organoleptic property/ Physicochemical parameter	Description/Value	
	BLO	TTO
Appearance	Oily liquid	
Solubility	Immiscible in water, but soluble in non-polar organic solvents	
Color	Light yellow	Nearly colorless to pale yellow
Odor	Creosote-like note	Camphor-like note
Taste	Pungent and sharp	Bitter and spicy
Specific gravity	0.97 \pm 0.01	0.89 \pm 0.01
Acid index (mg KOH/g)	7.09 \pm 0.17	1.49 \pm 0.16
Ester index (mg KOH/g)	12.05 \pm 0.56	16.39 \pm 0.16
Saponification index (mg KOH/g)	19.14 \pm 0.42	17.88 \pm 0.01

Note: The data are presented as mean \pm standard deviation and $p < 0.05$.

3.2 Chemical Composition of Essential Oil

Table 2: Chemical composition of essential oils from *Piper betle* and *Melaleuca alternifolia*

BLO		TTO	
Identified compound	Quantity (%)	Identified compound	Quantity (%)
Eugenol	39.21	Terpinen-4-ol	49.62

Comment [H26]: What about blended oil, provide data

Eugenol acetate	16.42	γ -Terpinene	18.08
4-Allyl-1,2-diacetoxybenzene	12.24	α -Terpinene	9.16
Terpinen-4-ol	6.58	<i>p</i> -Cymene	5.89
α -Cadinol	6.13	α -Terpineol	4.94
γ -Terpinene	3.46	Terpinolene	3.47
Sabinene	2.14	α -Pinene	2.02
γ -Muurolene	2.06	Limonene	1.34
Germacrene-D	1.65	Aromandendrene	0.99
β -Phellandrene	1.27	β -Thujene	0.69
Eucalyptol	1.21	β -Myrcene	0.64
α -Muurolene	1.17	β -Pinene	0.52
Caryophyllene	0.92		

The organoleptic and physicochemical properties of essential oils from BLO and TTO are presented in Table 1. Although organoleptic evaluation is highly subjective, it may provide reliable information and is helpful in evaluating the quality of given essential oils. The obtained BLO is light yellow oily liquid, insoluble in water and soluble in non-polar organic solvents. It resembles to the smell of creosote, and with a sharp, pungent taste. Meanwhile, the obtained TTO is nearly colorless or pale yellow oily liquid, non-soluble in water but soluble in non-polar organic solvents. It has a camphoraceous odor, and with a spicy, bitter taste. As can be seen from Table 2, the essential oil from *P. betle* contains 13 compounds representing approximately 95% of the total oil. The chemical composition of BLO which was determined by GC-MS revealed the presence of eugenol (39.21%) followed by other components: eugenol acetate (16.42%), 4-allyl-1,2-diacetoxybenzene (12.24%), terpinen-4-ol (6.58%), α -cadinol (6.13%), γ -terpinene (3.46%), and sabinene (2.14%). Meanwhile other compounds made contributions less than 2% each to the total composition of essential oil. As reported from previous literature, the components of essential oil were influenced by several factors, including plant species, genotypes, climatic conditions, cultivation, ages of the plants, harvesting time, botanical origin and experimental conditions. Thus, the biological activity of essential oil will likely vary depending on various factors, particularly the diversity of microbial strains and the complexity of single molecules present in essential oils. Generally, based on the most common molecules present in BLO obtained from different parts of Asia, constituents of BLO can be classified into eight different chemotypes, including anethole, chavicol, chavibetol, eugenol, eugenol acetate, germacrene-D, isoeugenol, and safrole [5,33]. However, in reality, major constituents of each essential oil are attributed to

Comment [H27]: Modify to highlight the major constituent in each part of the plant, describe the obvious differences in constituents among the various plants and then the yield variations of the predominant common constituent(s) among the various plants

commonly between 2 and 4 chemotypes. In this research, the obtained BLO reveals the absence of five chemotypes including anethole, chavicol, chavibetol, isoeugenol, and safrole. However, eugenol which has been recognized as the most powerful component notably responsible for various biological properties particularly antiseptic potency was accounted to compose approximately 40% of the total constituents of BLO.

On the other hand, the essential oil from *M. alternifolia* consists of 12 compounds accounting for almost 97% of the total oil. The major components of TTO were terpinen-4-ol (49.62%), followed by other components: γ -terpinene (18.08%), α -terpinene (9.16%), *p*-cymene (5.89%), α -terpineol (4.94%), terpinolene (3.47%), and α -pinene (2.02%). Meanwhile, other substances contributed lower than 2% each to the total composition of essential oil. TTO is a mixture of more than 100 natural volatile compounds. The concentration and chemical constituents of TTO may be extremely variable depending on several factors such as chemotypes, places of cultivation, climate conditions or extraction methods. Basically, TTO has been categorized into six chemotypes; each chemotype has been characterized by typical chemical constituents. Of which, the terpinen-4-ol chemotype is specified as type 1, followed by terpinolene as type 2 and four 1,8-cineole chemotypes [11]. Notably, the chemical constituents of TTO are regulated by International Standard ISO 4730:2017 for "Essential oil of *Melaleuca*, Terpinen-4-ol type (Tea Tree oil)", providing specific quantity ranges (in percentage) of 15 major components among more than 100 components identified in pure Australian TTO. Organoleptic characteristics and physical properties are also included in ISO description. As regulated, commercial TTO products to be marketed must be always of terpinen-4-ol chemotype, type 1, typically consisting of at least 35% terpinen-4-ol and eucalyptol (1,8-cineole) must be less than 10% [34,35]. In practice, however, approximately 50% of the commercial products did not meet the ISO specifications. In this study, the obtained TTO reveals the absence of four 1,8-cineole chemotypes. However, terpinen-4-ol which has been confirmed as the principal active component significantly responsible for various biological activities especially antibacterial property was accounted to form approximately 50% of the total composition of TTO. Noticeably, a comparison of constituents between BLO and TTO shows the presence of two major compounds, namely terpinen-4-ol and γ -terpinene, in both essential oils, but at significantly different concentrations (Table 2). Nevertheless, since these two compounds are well known as major molecules substantially responsible for antibacterial activity of both BLO and TTO, the synergistic antibacterial efficacy may be likely variable depending on the ratio of essential oils in combination prepared for the assays against microbial strains.

3.3 AST and MIC Values

Table 3: AST and MIC values of essential oils from *Piper betle* and *Melaleuca alternifolia*

Microorganism	AST and MIC Value			
	BLO		TTO	
	AST	MIC (%)	AST	MIC (%)
<i>Bacillus subtilis</i>	+	0.125	++	0.500
<i>Enterococcus faecalis</i>	++	0.125	++	0.125
<i>Staphylococcus aureus</i>	++	0.125	+	0.500
Methicillin-resistant <i>Staphylococcus aureus</i>	+	2.000	+	0.500
<i>Acinetobacter baumannii</i>	++	0.125	++	0.125
<i>Escherichia coli</i>	+	0.125	+	0.125
<i>Klebsiella pneumoniae</i>	++	0.125	+	0.125
<i>Pseudomonas aeruginosa</i>	+	2.000	+	2.000
<i>Salmonella sp</i>	++	0.125	++	0.125

Note: susceptible (++); intermediate (+).

The results detailed in Table 3 clearly show that BLO and TTO were found effective against all microbial strains tested, but at different levels. Differences in the susceptibility of the tested microorganisms to BLO and TTO were qualitatively identified. Five microbial strains were found susceptible to BLO and four were intermediate, whereas four microorganisms were confirmed susceptible to TTO and five were intermediate. MIC values of both BLO and TTO were varying between 0.125% and 2.000%. BLO was able to powerfully inhibit the growth of *B. subtilis*, *E. faecalis*, *S. aureus*, *A. baumannii*, *E. coli*, *K. pneumoniae* and *Salmonella sp* with the same lowest MIC value of 0.125%, whereas both MRSA and *P. aeruginosa* displayed the least sensitivity to BLO with the same highest MIC value of 2.000%. TTO, on the other hand, efficaciously inhibited the growth of *E. faecalis*, *A. baumannii*, *E. coli*, *K. pneumoniae* and *Salmonella sp* with the same lowest MIC value of 0.125%. TTO also showed potent inhibition against *B. subtilis*, *S. aureus* and MRSA with the same MIC value of 0.500%. Among nine bacteria tested, however, *P. aeruginosa* exhibited the lowest sensitivity to TTO with the highest MIC value of 2.000%.

Numerous studies have been conducted on elucidating the mechanisms of action of essential oils. In general, the antibacterial mechanisms of essential oil are attributed to a variety of factors, including the characteristics of essential oil chemotypes, the quantity of major single chemical constituents present in essential oils and the diversity of microbial strains. Therefore, it might not be possible to postulate a single mode of action of essential oil against various microorganisms. As reported from previous literature, most of the

Comment [H28]: The table do not show concentration dependent effect, Any explanation?

Comment [H29]: There is no positive control used so how true is this statement?

Comment [H30]: Provide the photos of the disc to show the real assay

Comment [H31]: Any statistical evidence to show? Example; p-value?

Comment [H32]: 17

Comment [H33]: 17

researches were mainly involved in the physicochemical properties of pure single molecules and the characteristics of investigated microorganisms. Concerning the nature of chemical constituents, plant essential oils are not true oils, because they are not composed of lipids. Instead, essential oils truly possess a complex mixture of naturally occurring volatile compounds, mainly consisting of terpenoids, phenolics and phenylpropanoids. These compounds, being hydrophobic molecules, are capable of achieving a potent binding affinity to different lipophilic molecular structures such as lipids, proteins, lipoproteins, glycoproteins or glycolipids. It has been previously reported that numerous components present in various essential oils have been identified as key molecules responsible for biological activities and subsequently subjected to investigating the antibacterial mechanism of action [36-41]. As scientifically reviewed elsewhere, in general, the most commonly possible antibacterial mechanism of essential oils is involved in the bacterial cell wall and plasma membrane compositions which are identified as the main target sites of essential oils. Owing to the hydrophobicity nature of key constituents present in essential oils, hydrocarbons are capable of partitioning into the cell wall and cytoplasmic membrane of microorganisms. Different molecules present in essential oils may have different interactions with cell plasma membrane with regards to the types of interactions and strengths of binding affinity. Therefore, the disturbance of cell plasma membrane integrity and permeability may be likely to cause negative impacts in several different ways which can be briefly described as follows: (1) the leakage of essential metabolites; (2) efflux of RNA and DNA from the cytoplasm; (3) loss of vital ions such as Na^+ , K^+ , Ca^{2+} or Mg^{2+} ; (4) depletion of proton pumps in the plasma membrane; (5) depletion of ATP during biosynthesis; (6) disruption of cell respiration; (7) inactivation of various bacterial enzymes such as ATPase, amylase, histidine carboxylase or protease; (8) turbulence of electron transport system; (9) coagulation of cellular components in the cytoplasmic membrane; (10) blockage of protein expression; (11) inhibition of nucleic acid synthesis and function or (12) inhibition of the production and secretion of toxic bacterial metabolites [42-49]. These series of events may be responsible for a number of serious sequelae. Finally, a consequence of any of these impairments may be induction of bacterial cell lysis and cell death.

Moreover, the complexity of antibacterial mechanism of essential oils is even attributed to the difference in susceptibility of Gram-positive and Gram-negative bacteria to a variety of essential oils. In general, Gram-negative bacteria exhibit more natural resistance to various types of essential oils than Gram-positive strains due to the difference in the cell membrane structures and compositions. Gram-negative bacteria possess hydrophilic lipopolysaccharide (LPS)-based outer membrane which play a key role in structural integrity of the bacterial cell

Comment [H34]: 17

Comment [H35]: 17

membrane and also considerably contribute to the negative charge of the cell membrane on the basis of negatively charged carbohydrates and phosphates bonded in the polysaccharides. A thick layer of LPS which is surrounding the cell wall of Gram-negative bacteria establishes a physical barrier limiting the diffusion of macromolecules and hydrophobic compounds, thereby facilitating more resistance to the lipophilicity nature of essential oils. In contrast, Gram-positive cell wall is mainly composed of peptidoglycan, but lack of lipopolysaccharides, which may be likely more vulnerable to essential oils and their components. It should be noted, however, there has been no general principles relevant to properly assessing Gram susceptibility. Practically, some studies have shown no significant difference in susceptibility between Gram-positive and Gram-negative bacteria to essential oils and their components; even in some cases, Gram-negative strains were found greater susceptible than some Gram-positive species to certain essential oils and their constituents. Such conflicting studies may gain considerable attention from researchers in various fields and even trigger more investigation for further understanding the antibacterial mechanism of action of essential oils and their components [50-52]. It is noteworthy to know that some essential oils may be capable of inhibiting the cell-cell signal communication between bacterial cells. This communication, commonly known as quorum sensing, is one of the vital biochemical processes in the biofilm formation. The biofilm is typically composed of a complex mixture of highly polar biomolecules mainly consisting of proteins, nucleic acids and polysaccharides from resident bacteria. The biofilm formation is best known as part of microbial action against antibiotics, owing to the formation of physical barriers to limit the infiltration of antibiotics. Therefore, the inhibition of biofilm formation could be one of the capacities of various potential essential oils in fighting against bacterial infections. Noticeably, the antibacterial mechanisms of eugenol and terpinen-4-ol have been intensively investigated using relevant modern techniques, such as scanning electron microscopy (SEM), transmission electron microscopy (TEM), confocal laser scanning microscopy (CLSM) or measurement of release of 260-nm absorbing materials and proteins. By such means, the physiological and morphological changes have been observed and the biochemical parameters have been recorded. Generally, the antibacterial mechanism of both components against various pathogenic microorganisms might be likely related to the disruption of cell wall and the damage of cytoplasmic membrane. Additionally, eugenol was able to cause the leakage of intracellular components and effectively inactivated various bacterial enzymes such as ATPase, amylase, protease or histidine carboxylase. Terpinen-4-ol, on the other hand, could strongly induce the release of Ca^{2+} , Mg^{2+} and enzyme lactate dehydrogenase (LDH); and powerfully inhibited the synthesis of protein, DNA and ATPase.

Furthermore, both components were found to be capable of effectively inhibiting the biofilm formation resulting in increased number of dead cells [36-41,53-64].

3.4 FICI Values

Table 4: FICI values of blended essential oils from *Piper betle* and *Melaleuca alternifolia*

Comment [H36]: 21

Comment [H37]: How you prepare test solutions ?

Microorganism	Essential oil	MIC (%)		FIC (%)			Interaction
		MIC alone	MIC in combination	FIC BLO	FIC TTO	FICI	
<i>Bacillus subtilis</i>	BLO	0.125	0.0156	1/8 MIC	1/8 MIC	0.250	Synergistic
	TTO	0.125	0.0156				
<i>Enterococcus faecalis</i>	BLO	0.125	0.0313	1/4 MIC	1/8 MIC	0.375	Synergistic
	TTO	0.125	0.0156				
<i>Staphylococcus aureus</i>	BLO	0.125	0.0313	1/4 MIC	1/16 MIC	0.313	Synergistic
	TTO	0.500	0.0313				
Methicillin-resistant <i>Staphylococcus aureus</i>	BLO	2.000	0.0625	1/32 MIC	1/4 MIC	0.281	Synergistic
	TTO	0.500	0.1250				
<i>Acinetobacter baumannii</i>	BLO	0.125	0.0313	1/4 MIC	1/8 MIC	0.375	Synergistic
	TTO	0.125	0.0156				
<i>Escherichia coli</i>	BLO	0.125	0.0313	1/4 MIC	1/8 MIC	0.375	Synergistic
	TTO	0.125	0.0156				
<i>Klebsiella pneumoniae</i>	BLO	0.1250	0.0625	1/2 MIC	1/8 MIC	0.625	Additive
	TTO	0.125	0.0156				
<i>Pseudomonas aeruginosa</i>	BLO	2.000	1.0000	1/2 MIC	1/4 MIC	0.750	Additive
	TTO	2.000	0.5000				
<i>Salmonella sp</i>	BLO	0.125	0.0313	1/4 MIC	1/8 MIC	0.375	Synergistic
	TTO	0.125	0.0156				

The combination of BLO and TTO exhibited significant results as detailed in Table 4. Synergistic antibacterial effects of blended essential oils from *P. betle* and *M. alternifolia* against nine different microbial strains showed the considerable reduction in MIC values of both BLO and TTO at different levels, resulting in FICI values varying between 0.250% and 0.750%. In general, the blended essential oils from *P. betle* and *M. alternifolia* reduced the MIC values of BLO and TTO by 2-32-fold, as gains. There was an 8-fold reduction in MIC value for *B. subtilis*, 2-4 fold for *P. aeruginosa*, 2-8 fold for *K. pneumoniae*, 4-8 fold for *E. faecalis*, *A. baumannii*, *E. coli* and *Salmonella sp*, 4-16 fold for *S. aureus*, and 4-32 fold for MRSA. Meanwhile, the synergistic antibacterial effects of blended essential oils from *P. betle* and *M. alternifolia* were recorded in seven of the nine bacteria tested, including *B. subtilis*

(FICI = 0.250%), *E. faecalis* (FICI = 0.375%), *S. aureus* (FICI = 0.313%), MRSA (FICI = 0.281%), *A. baumannii* (FICI = 0.375%), *E. coli* (FICI = 0.375%) and *Salmonella sp* (FICI = 0.375%). In contrast, additive effects were observed in both Gram-negative bacteria including *K. pneumoniae* and *P. aeruginosa* when treated with blended essential oils resulting in FICI value of 0.625% and 0.750%, respectively.

The combination of essential oils has been explored historically for hundreds of years for both aromatic and therapeutic benefits. Aromatherapy, one of the most popular alternative and complementary therapies, is the practice of blending essential oils topically and aromatically for overall health and well-being. However, there has been not enough scientific research to elucidate their effectiveness in human health. In recent years, numerous efforts have been made to exploit the combination of essential oils in various different fields including cosmetics, food preservatives and pharmaceutical industries. In general, the combination of essential oils has been primarily focused on enhancing the therapeutic efficiency and minimizing the dose of essential oils, thus decreasing the toxicity and adverse reactions. In addition, the interactions between molecules may enhance the solubility and dissolution rate leading to the potential development of new dosage forms and possible improvement of bioavailability. There has been currently an increasing demand for synergistic antibacterial effects of blended essential oils due to not only increasing antibiotic-resistant bacteria at an alarming rate over the last decade, but also negative concerns regarding safety in use of synthetic chemical-based drugs. As mentioned above, the mode of action of individual essential oil has been partially elucidated; however, the mechanism of action of multiple essential oils basically remains unclear, owing to the limited numbers of investigation on the synergistic antibacterial effects of combined essential oils. The diversified and complicated nature of phytoconstituents of essential oils and the complexity of chemical molecules present in blended essential oils would be a big challenge in elucidating the mechanisms of action. Different chemical molecules may possess different characteristics with regards to chemotypes, ring structures, functional groups, physical and chemical properties. As a result, bioactive compounds may have different mechanisms acting on various different target sites. Thus, it is difficult to predict the synergistic antibacterial effects of blended essential oils against various microbial strains. As scientifically reported elsewhere, numerous studies have been conducted on the synergistic antibacterial effects of essential oils blended with other available substances including commonly used antibiotics, and plant-derived components (such as plant extracts, essential oils or single molecules isolated from plants) [65-76]. In principles, the interactions between components from essential oils may lead to antagonistic, indifferent, additive or synergistic

effects. Therefore, microbiological assays would be helpful in not only determining the potency of antibacterial synergism of blended essential oils but also considerably contributing to the elucidation of antibacterial mechanism of action. In this investigation, neither indifferent nor antagonistic effect was detected in any of tested combination of BLO and TTO against nine microbial strains selected for the assay. Instead, the combination of BLO and TTO has demonstrated the powerful synergistic antimicrobial effectiveness against seven of the nine studied microbial strains. Of the seven interactions synergistically detected, four strains were Gram-positive and three others were Gram-negative bacteria. It is noteworthy to know that additive effects were only detected against both Gram-negative bacteria including *K. pneumoniae* and *P. aeruginosa*. This result might partly support the postulation on the difference in susceptibility of Gram-positive and Gram-negative bacteria to antimicrobial agents. Noticeably, among five Gram-negative bacterial strains tested, *P. aeruginosa* was found to be intermediate to both single BLO and TTO with the same highest MIC value of 2.000% (Table 3), apparently confirming the considerable variation of both lipid and polysaccharide components in LPS which may be likely attributed to the notable difference in genera and species of Gram-negative bacteria involved; and even in some cases, to the presence of particular bacterial strain factors. Furthermore, these findings were even consolidated by previous researches as *P. aeruginosa* was also found to be less susceptible to both single BLO and TTO [77-81], thereby partially reflecting the complexity of natural constituents of essential oils and the diversity of biological system of living microorganisms, especially the characteristics of bacterial cell wall and cytoplasmic membrane and the proper function of outer membrane and LPS of Gram-negative bacteria in facilitating resistance against hydrophobicity nature of various molecules present in essential oils.

4. CONCLUSION

Overall, the combination of essential oils from *Piper betle* and *Melaleuca alternifolia* had more significant antibacterial effects against all of the nine studied microbial strains than single essential oil. The synergistic antibacterial effects were recorded in seven of the nine microbial strains tested, whereas the additive effects were only observed in both Gram-negative bacteria including *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*. This promising research scientifically contributes to the herbal medicine database system and promotes the significant potential of synergistic antibacterial efficacy of blended essential oils from BLO and TTO that might be exploited for the development of plant-based therapeutic agents for the topical treatment against skin infectious diseases. Nevertheless, further studies are necessary to investigate comprehensively the possible synergistic effects of

blended essential oils from BLO and TTO against not only other bacteria but also various microorganisms such as viruses, fungi, yeasts, molds or protozoa. Furthermore, the synergistic interactions need to be characterized at a molecular level for a better understanding of the synergistic mechanism, which would be core fundamentals of biochemistry and molecular biology for future research.

Comment [H38]: Modify to highlight the major compounds findings on each plant in accordance with aim of your research

Comment [H39]: Modify to highlight the differences in antibacterial activity between single and blended oils correlate with the aim of the study.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

REFERENCES

- [1] Surjowardojo P, Setyowati E, Ambarwati I. Antibacterial Effects of Green Betel (*Piper betle* Linn.) Leaf Against *Streptococcus agalactiae* and *Escherichia coli*. *AGRIVITA, Journal of Agricultural Science*. 2019;41(3):569–574. DOI: <https://doi.org/10.17503/agrivita.v41i3.2437>.
- [2] Carsono N, Tumilaar SG, Kurnia D, Latipudin D, Satari MH. A Review of Bioactive Compounds and Antioxidant Activity Properties of Piper Species. *Molecules*. 2022;27:6774. DOI: 10.3390/molecules27196774. PMID: 36235309; PMCID: PMC9573611.
- [3] Kapdo R, Ibrahim M. Chemical component analysis of betel leaf essential oil fraction (*piper betle* Linn.) and antibacterial efficacy against different gram-positive bacteria. *International Journal on ObGyn and Health Sciences*. 2022;1(1):1-9. Available: <https://trigin.pelnus.ac.id/index.php/ObGyn/article/view/57>.
- [4] Sakinah D, Rusdi, Misfadhila S. Review of traditional use, phytochemical and pharmacological activity of *Piper betle* L. *Gal Int J Health Sci Res*. 2020;5(3):59-66. Available: https://www.gijhsr.com/GIJHSR_Vol.5_Issue.3_July2020/11.pdf.
- [5] Vandana D, Shalini T. Review study on potential activity of *Piper betle*. *Journal of Pharmacognosy and Phytochemistry*. 2014;3(4):93-98. Available: https://www.phytojournal.com/vol3Issue4/Issue_nov_2014/17.1.pdf.
- [6] Umar RA, Sanusi NA, Zahary MN, Rohin MAK, Ismail S. Chemical Composition and the Potential Biological Activities of Piper Betel – A Review. *Malaysian Journal of Applied Sciences*. 2018;3(1):1-8. Available: <https://journal.unisza.edu.my/myjas/index.php/myjas/article/view/69>.
- [7] Karunanithi S, U Eswaran GM, Guha P, Srivastav PP. A Review on Piper betle L.: Antioxidant, Antimicrobial, Extraction and Application in Food Product Development. *Acta Scientific Nutritional Health*. 2023;7(1):49-61. DOI:10.31080/ASNH.2023.07.1170.

[8] Nayaka NMDMW, Sasadara MMV, Sanjaya DA, Yuda PESK, Dewi NLKAA, Cahyaningsih E, et al. Piper betle (L): Recent Review of Antibacterial and Antifungal Properties, Safety Profiles, and Commercial Applications. *Molecules*. 2021;26(8):2321. DOI.org/10.3390/molecules26082321.

[9] Delia DCO, Teresita C. The activity of the leaf essential oil of Philippine *Piper betel* against dermatophytes and *Candida albicans*. *Philippine Journal of Systematic Biology*. 2019;13(2). DOI 10.26757/pjsb2019b13002.

[10] Sunita L, Malay KD, Sudarshana B. An Overview on Tea Tree (*Melaleuca Alternifolia*) Oil. *International Journal of Pharmaceutical and Phytopharmacological Research*. 2013;3(3):250-253. Available: <https://ejppr.com/storage/models/article/ghKc7OvScRawdtASi05OTQlJvSPgU5BE DsPAdEGZShNEqqzkLxwlrz4Fxy5j/an-overview-on-tea-tree-melaleuca-alternifolia-oil.pdf>.

[11] Carson CF, Hammer KA, Riley TV. *Melaleuca alternifolia* (Tea Tree) Oil. A Review of Antimicrobial and Other Medicinal Properties. *Clinical Microbiology Reviews*. 2006;19(1):50-62. DOI: 10.1128/CMR.19.1.50-62.2006. PMID: 16418522; PMCID: PMC1360273.

[12] de Groot AC, Schmidt E. Tea tree oil: contact allergy and chemical composition. *Contact Dermatitis*. 2016;75(3):129-43. DOI: 10.1111/cod.12591. Epub 2016 May 13. PMID: 27173437.

[13] Li X, Shen D, Zang Q, Qiu Y, Yang X. Chemical Components and Antimicrobial Activities of Tea Tree Hydrosol and Their Correlation With Tea Tree Oil. *Natural Product Communications*. 2021;16(9):1-7. DOI:10.1177/1934578X211038390.

[14] Zhang X, Guo Y, Guo L, Jiang H, Ji Q. *In Vitro* Evaluation of Antioxidant and Antimicrobial Activities of *Melaleuca alternifolia* Essential Oil. *BioMed Research International*. 2018; Article ID 2396109. DOI.org/10.1155/2018/2396109.

1. [15] Sevik R, Akarca G, Kilinc M, Asciglu Ç. Chemical Composition of Tea Tree (*Melaleuca alternifolia*) (Maiden & Betche) Cheel Essential Oil and Its Antifungal Effect on Foodborne Molds Isolated from Meat Products. *Journal of Essential Oil Bearing Plants*. 2021;24(3):561-570.
2. DOI:10.1080/0972060X.2021.1942232.

[16] Ibrahim K. Critical Evaluation of *Melaleuca alternifolia*. A Review of the Phytochemical Profile, Pharmacological Attributes and Medicinal Properties in the Botanical, Human and Global Perspectives. *Open Journal of Medicinal Chemistry*. 2021;11(1). DOI: 10.4236/ojmc.2021.111001.

[17] Moshe B. Small and efficient distillation apparatus for extraction of essential oils from plant matter. United States Patent Application Publication. 2008;Pub. No.: US 2008/0128260A1. Available: <https://patents.google.com/patent/US20080128260A1/en>.

[18] Scion Instruments 456-GC Reference Manual. Available: <https://www.manualslib.com/manual/1626438/Scion-Instruments-456-Gc.html>.

[19] AOAC. Official Methods of Analysis of the Association of Official Analytical Chemists. 15th Ed. Washington DC. 1990;1:1223.

[20] Baron S, editor. Medical Microbiology. 4th edition. Galveston (TX): University of Texas Medical Branch at Galveston; 1996.
Available: <https://www.ncbi.nlm.nih.gov/books/NBK7627/>.

[21] Laboratory Guide: Methodologies for Antimicrobial Susceptibility Testing (Manual). Committee on Trade and Investment (CTI), Sub-Committee on Standards and Conformance (SCSC). 2020.
Available: <https://www.apec.org/Publications/2020/05/Laboratory-Guide-Methodologies-for-Antimicrobial-Susceptibility-Testing>.

[22] Jorgensen JH, Ferraro MJ. Antimicrobial Susceptibility Testing: A Review of General Principles and Contemporary Practices. Clinical Infectious Diseases. 2009;49(11):1749-1755.
DOI: 10.1086/647952. PMID: 19857164.

[23] Mark L. Antimicrobial Susceptibility Testing. American Association for Clinical Chemistry (AACC). 2017.
Available: <https://www.aacc.org/science-and-research/clinical-chemistry-trainee-council/trainee-council-in-english/pearls-of-laboratory-medicine/2017/antimicrobial-susceptibility-testing>.

[24] Antimicrobial Susceptibility Testing (AST): Introduction, Methods, Procedure and Result Interpretation. Universe84a.
Available: <https://universe84a.com/antimicrobial-susceptibility-testing-ast/>.

[25] Gajic I, Kabic J, Kekic D, Jovicevic M, Milenkovic M, Mitic CD, et al. Antimicrobial Susceptibility Testing: A Comprehensive Review of Currently Used Methods. Antibiotics. 2022;11(4):427.
DOI.org/10.3390/antibiotics11040427.

[26] Lila R. Minimal inhibitory concentration (MIC) test and determination of antimicrobial resistant bacteria. In Laboratory manual of standardized methods for antimicrobial sensitivity tests for bacteria isolated from aquatic animals and environment (pp.31-55). Tigbauan, Iloilo, Philippines: Aquaculture Department, Southeast Asian Fisheries Development Center. 2004.
Available: <https://repository.seafdec.org.ph/bitstream/handle/10862/1637/Chapter3-Minimal-Inhibitory-Concentration-Test.pdf?sequence=1&isAllowed=y>.

[27] Kowalska-Krochmal B, Dudek-Wicher R. The Minimum Inhibitory Concentration of Antibiotics: Methods, Interpretation, Clinical Relevance. Pathogens. 2021;10(2):165.
DOI: 10.3390/pathogens10020165. PMID: 33557078; PMCID: PMC7913839.

[28] Determination of minimum inhibitory concentrations (MICs) of antibacterial agents by agar dilution. European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID). 2009.
Available: <https://onlinelibrary.wiley.com/doi/10.1046/j.1469-0691.2000.00142.x>.

[29] Balouiri M, Sadiki M, Ibsouda SK. Methods for *in vitro* evaluating antimicrobial activity: A review. J Pharm Anal. 2016;6(2):71-79.
DOI: 10.1016/j.jpha.2015.11.005. Epub 2015 Dec 2. PMID: 29403965; PMCID: PMC5762448.

[30] van Vuuren S, Viljoen A. Plant-based antimicrobial studies--methods and approaches to study the interaction between natural products. *Planta Med.* 2011;77(11):1168-82. DOI: 10.1055/s-0030-1250736. Epub 2011 Jan 31. Erratum in: *Planta Med.* 2012 Feb;78(3):302. PMID: 21283954.

[31] Faleiro ML, Miguel MG. Chapter 6 - Use of Essential Oils and Their Components against Multidrug-Resistant Bacteria. In: Mahendra KR, Kateryna VK, editors. *Fighting Multidrug Resistance with Herbal Extracts, Essential Oils and their Components.* Academic Press; 2013. DOI:10.1016/B978-0-12-398539-2.00006-9.

[32] Kalpna DR, Mital JK, Sumitra VC. Chapter 11 - Medicinal Plants as Alternative Sources of Therapeutics against Multidrug-Resistant Pathogenic Microorganisms Based on Their Antimicrobial Potential and Synergistic Properties. In: Mahendra KR, Kateryna VK, editors. *Fighting Multidrug Resistance with Herbal Extracts, Essential Oils and their Components.* Academic Press; 2013. DOI:10.1016/B978-0-12-398539-2.00011-2.

[33] Guha P, Nandi S. Essential Oil of Betel Leaf (*Piper betle* L.): A Novel Addition to the World Food Sector. 2019. DOI:10.1007/978-3-030-16546-8_5.

[34] International Standard. Essential oil of *Melaleuca*, terpinen-4-ol type (Tea Tree oil). iTeH STANDARD PREVIEW (standards.iteh.ai) ISO 4730:2017. Third edition 2017-02. Available: <https://cdn.standards.iteh.ai/samples/69082/302bafb843a64a95b8c74676e282d921/ISO-4730-2017.pdf>.

[35] ISO4730: 2017 Standard. Australia Tea tree oil *Melaleuca alternifolia*. *A Natural Treatment-Nature's Best*. Available: <https://teatree.org.au/standards.php#:~:text=ISO4730%3A%202017%20Standard,a%20number%20of%20physical%20parameters>.

[36] Ashrafudoulla M, Mizan MFR, Ha AJ, Park SH, Ha SD. Antibacterial and antibiofilm mechanism of eugenol against antibiotic resistance *Vibrio parahaemolyticus*. *Food Microbiol.* 2020;91:103500. DOI: 10.1016/j.fm.2020.103500. Epub 2020 Apr 11. PMID: 32539983.

[37] Su R, Bai X, Liu X, Song L, Liu X, Zhan X, et al. Antibacterial Mechanism of Eugenol Against *Shigella sonnei* and Its Antibacterial Application in Lettuce Juice. *Foodborne Pathogens and Disease.* 2022:779-786. DOI.org/10.1089/fpd.2022.0046.

[38] Mak KK, Kamal MB, Ayuba SB, Sakirolla R, Kang YB, Mohandas K, et al. A comprehensive review on eugenol's antimicrobial properties and industry applications: A transformation from ethnomedicine to industry. *Phcog Rev.* 2019;13:1-9. Available: https://www.phcogrev.com/sites/default/files/PhcogRev_2019_13_25_1.pdf.

[39] Bai X, Li X, Liu X, Xing Z, Su R, Wang Y, et al. Antibacterial Effect of Eugenol on *Shigella flexneri* and Its Mechanism. *Foods.* 2022;11:2565. DOI.org/10.3390/foods11172565 PMID: 36076751; PMCID: PMC9455010.

- [40] Shapira S, Pleban S, Kazanov D, Tirosh P, Arber N. Terpinen-4-ol: A Novel and Promising Therapeutic Agent for Human Gastrointestinal Cancers. *PLoS One*. 2016;11(6):e0156540. DOI: 10.1371/journal.pone.0156540. PMID: 27275783; PMCID: PMC4898785.
- [41] Cordeiro L, Figueiredo P, Souza H, Sousa A, Andrade-Júnior F, Medeiros D. Terpinen-4-ol as an Antibacterial and Antibiofilm Agent against *Staphylococcus aureus*. *Int. J. Mol. Sci*. 2020;21:4531. DOI.org/10.3390/ijms21124531 PMID: 32630600; PMCID: PMC7350221.
- [42] Li WR, Li HL, Shi QS, Sun TL, Xie XB, Song B, et al. The dynamics and mechanism of the antimicrobial activity of tea tree oil against bacteria and fungi. *Applied Microbiology and Biotechnology*. 2016;100(20):8865-8875. DOI: 10.1007/s00253-016-7692-4. PMID: 27388769.
- [43] Xiang F, Bai J, Tan X, Chen T, Yang W, He F. Antimicrobial activities and mechanism of the essential oil from *Artemisia argyi* Levl. et Van. var. *argyi* cv. Qiai. *Industrial Crops and Products*. 2018;125(1):582-587. DOI.org/10.1016/j.indcrop.2018.09.048.
- [44] Chouhan S, Sharma K, Guleria S. Antimicrobial Activity of Some Essential Oils-Present Status and Future Perspectives. *Medicines (Basel)*. 2017;4(3):58. DOI: 10.3390/medicines4030058. PMID: 28930272; PMCID: PMC5622393.
- [45] Zhang J, Ye KP, Zhang X, Pan DD, Sun YY, Cao JX. Antibacterial Activity and Mechanism of Action of Black Pepper Essential Oil on Meat-Borne *Escherichia coli*. *Front. Microbiol*. 2017(7):2094. DOI: 10.3389/fmicb.2016.02094 PMID: 28101081; PMCID: PMC5209337.
- [46] Yang SK, Tan NP, Chong CW, Abushelaibi A, Lim SH, Lai KS. The Missing Piece: Recent Approaches Investigating the Antimicrobial Mode of Action of Essential Oils. *Evolutionary Bioinformatics*. 2021;17. DOI:10.1177/1176934320938391 PMID: 34017165; PMCID: PMC8114247.
- [47] Li ZH, Cai M, Liu YS, Sun PL, Luo SL. Antibacterial Activity and Mechanisms of Essential Oil from *Citrus medica* L. var. *sarcodactylis*. *Molecules*. 2019 Apr 22;24(8):1577. DOI: 10.3390/molecules24081577. PMID: 31013583; PMCID: PMC6515347.
- [48] Tang C, Chen J, Zhang L, Zhang R, Zhang S, Ye S, et al. Exploring the antibacterial mechanism of essential oils by membrane permeability, apoptosis and biofilm formation combination with proteomics analysis against methicillin-resistant *Staphylococcus aureus*. *International Journal of Medical Microbiology*. 2020;310(5):151435. DOI.org/10.1016/j.ijmm.2020.151435.
- [49] Lopez-Romero JC, González-Ríos H, Borges A, Simões M. Antibacterial Effects and Mode of Action of Selected Essential Oils Components against *Escherichia coli* and *Staphylococcus aureus*. *Evidence-Based Complementary and Alternative Medicine*. 2015, Article ID 795435. DOI.org/10.1155/2015/795435.
- [50] Puškárová A, Bučková M, Kraková L, Pangallo D, Kozics K. The antibacterial and antifungal activity of six essential oils and their cyto/genotoxicity to human HEL 12469 cells. *Sci Rep*. 2017;7(1):8211.

DOI: 10.1038/s41598-017-08673-9. PMID: 28811611; PMCID: PMC5557807.

[51] Zairi A, Nour S, Zarrouk A, et al. Chemical composition, Fatty acids profile and Biological properties of *Thymus capitatus* (L.) Hoffmanns, essential Oil. *Sci Rep.* 2019;9: 20134.

DOI.org/10.1038/s41598-019-56580-y.

[52] Skala E, Rijo P, Garcia C, Sitarek P, Kalembe D, Toma M, et al. The Essential Oils of *Rhaponticum carthamoides* Hairy Roots and Roots of Soil-Grown Plants: Chemical Composition and Antimicrobial, Anti-Inflammatory, and Antioxidant Activities. *Oxidative Medicine and Cellular Longevity.* 2016; Article ID 8505384.

DOI.org/10.1155/2016/8505384. Epub 2016 Dec 18. PMID: 28074117; PMCID: PMC5203915.

[53] Mak KK, Kamal MB, Ayuba SB, Sakirolla R, Kang YB, Mohandas K, et al. A comprehensive review on eugenol's antimicrobial properties and industry applications: A transformation from ethnomedicine to industry. *Phcog Rev* 2019;13(25).

Available: https://www.phcogrev.com/sites/default/files/PhcogRev_2019_13_25_1.pdf.

[54] Angane M, Swift S, Huang K, Butts CA, Quek SY. Essential Oils and Their Major Components: An Updated Review on Antimicrobial Activities, Mechanism of Action and Their Potential Application in the Food Industry. *Foods.* 2022;11: 464.

DOI.org/ 10.3390/foods11030464. PMID: 35159614; PMCID: PMC8833992.

[55] Das B, Mandal D, Dash SK, et al. Eugenol Provokes ROS-Mediated Membrane Damage-Associated Antibacterial Activity against Clinically Isolated Multidrug-Resistant *Staphylococcus aureus* Strains. *Infectious Diseases: Research and Treatment.* 2016;9.

DOI:10.4137/IDRT.S31741. PMID: 26917967; PMCID: PMC4756864.

[56] Devi KP, Nisha SA, Sakthivel R, Pandian SK. Eugenol (an essential oil of clove) acts as an antibacterial agent against *Salmonella typhi* by disrupting the cellular membrane. *Journal of Ethnopharmacology.* 2010;130(1):107-115.

DOI: 10.1016/j.jep.2010.04.025. PMID: 20435121.

[57] Mohammadi Nejad S, Özgüneş H, Başaran N. Pharmacological and Toxicological Properties of Eugenol. *Turk J Pharm Sci.* 2017;14(2):201-206.

DOI: 10.4274/tjps.62207. Epub 2017 Aug 15. PMID: 32454614; PMCID: PMC7227856.

[58] Devi KP, Sakthivel R, Nisha SA, Suganthi N, Pandian SK. Eugenol alters the integrity of cell membrane and acts against the nosocomial pathogen *Proteus mirabilis*. *Arch Pharm Res.* 2013;36(3):282-92.

DOI: 10.1007/s12272-013-0028-3. Epub 2013 Feb 27. PMID: 23444040.

[59] Pramod K, Ansari SH, Ali J. Eugenol: a natural compound with versatile pharmacological actions. *Nat Prod Commun.* 2010;(12):1999-2006. PMID: 21299140.

Available:https://www.researchgate.net/publication/49815369_Eugenol_A_Natural_Compound_with_Versatile_Pharmacological_Actions.

[60] Elbestawy MKM, El-Sherbiny GM, Moghannem SA. Antibacterial, Antibiofilm and Anti-Inflammatory Activities of Eugenol Clove Essential Oil against Resistant *Helicobacter pylori*. *Molecules.* 2023;28(6):2448.

DOI.org/10.3390/molecules28062448. PMID: 36985419; PMCID: PMC10058968.

- [61] Bruna FMTA, Lidiane NB, Fernanda CBA, Mariana A, Vera LMR, José MS. The antibacterial effects of *Melaleuca alternifolia*, *Pelargonium graveolens* and *Cymbopogon martinii* essential oils and major compounds on liquid and vapor phase, *Journal of Essential Oil Research*.2016;28(3):227-233.
DOI: 10.1080/10412905.2015.1099571.
- [62] Carson CF, Mee BJ, Riley TV. Mechanism of action of *Melaleuca alternifolia* (tea tree) oil on *Staphylococcus aureus* determined by time-kill, lysis, leakage, and salt tolerance assays and electron microscopy. *Antimicrob Agents Chemother*. 2002;46(6):1914-20.
DOI: 10.1128/AAC.46.6.1914-1920.2002. PMID: 12019108; PMCID: PMC127210.
- [63] Mondello F, Fontana S, Scaturro M, Girolamo A, Colone M, Stringaro A. Terpinen-4-ol, the Main Bioactive Component of Tea Tree Oil, as an Innovative Antimicrobial Agent against *Legionella pneumophila*. *Pathogens*. 2022;11(6):682.
DOI.org/10.3390/pathogens11060682. PMID: 35745536; PMCID: PMC9229490.
- [64] Cordeiro L, Figueiredo P, Souza H, Sousa A, Andrade-Júnior F, Medeiros D, et al. Terpinen-4-ol as an Antibacterial and Antibiofilm Agent against *Staphylococcus aureus*. *Int J Mol Sci*. 2020;21(12):4531.
DOI: 10.3390/ijms21124531. PMID: 32630600; PMCID: PMC7350221.
- [65] Yap PS, Yiap BC, Ping HC, Lim SH. Essential Oils, A New Horizon in Combating Bacterial Antibiotic Resistance. *The Open Microbiology Journal*. 2014;8:6-14.
DOI: 10.2174/1874285801408010006. PMID: 24627729; PMCID: PMC3950955.
- [66] Fadila M, Tajelmolk A. Evaluation of antibacterial activity and synergistic effect between antibiotic and the essential oils of some medicinal plant. *Asian Pacific Journal of Tropical Biomedicine*. 2016;6(1):32-37.
DOI.org/10.1016/j.apjtb.2015.09.024.
- [67] Sharma K, Guleria S, Razdan V, Babu V. Synergistic antioxidant and antimicrobial activities of essential oils of some selected medicinal plants in combination and with synthetic compounds. *Industrial Crops and Products*. 2020;154:112569.
DOI.org/10.1016/j.indcrop.2020.112569.
- [68] Huang X, Lao Y, Pan Y, Chen Y, Zhao H, Gong L, et al. Synergistic Antimicrobial Effectiveness of Plant Essential Oil and Its Application in Seafood Preservation: A Review. *Molecules*. 2021;26(2):307.
DOI.org/10.3390/molecules26020307.
- [69] Alam M, Bano N, Ahmad T, Sharangi AB, Upadhyay TK, Alraey Y, et al. Synergistic Role of Plant Extracts and Essential Oils against Multidrug Resistance and Gram-Negative Bacterial Strains Producing Extended-Spectrum-Lactamases. *Antibiotics*. 2022;11:855.
DOI.org/10.3390/antibiotics11070855.
- [70] Wessal O, Balouiri M, El Houssaine H, Sandrine M, Hassane G. Synergistic antimicrobial activity of two binary combinations of marjoram, lavender, and wild thyme essential oils, *International Journal of Food Properties* 2017;20(12):3149-3158.
DOI: 10.1080/10942912.2017.1280504.
- [71] Fahimi S, Hajimehdipoor H, Shabanpoor H, Bagheri F, Shekarchi M. Synergic antibacterial activity of some essential oils from Lamiaceae. *Research Journal of Pharmacognosy*. 2015;2(3):23-29.

Available:https://www.researchgate.net/publication/313049591_Synergic_antibacterial_activity_of_some_essential_oils_from_Lamiaceae.

[72] Raquel R, María V, Amparo C. Study of the potential synergistic antibacterial activity of essential oil components using the thiazolyl blue tetrazolium bromide (MTT) assay. *LWT - Food Science and Technology*. 2019;101:183–190. DOI.org/10.1016/j.lwt.2018.10.093.

[73] Low WL, Kenward K, Britland ST, Amin MC, Martin C. Essential oils and metal ions as alternative antimicrobial agents: a focus on tea tree oil and silver. *Int Wound J*. 2017;14(2):369-384. DOI: 10.1111/iwj.12611. Epub 2016 May 5. PMID: 27146784; PMCID: PMC7949732.

[74] Yustina SH, Yohanes MSD, Rakhel NP, Lia ES. Antagonistic Antibacterial Effect of Betel and Red Betel Combination against Gram-positive and Gram-negative Bacteria. *International Journal of Current Microbiology and Applied Sciences*.2018;7(5). DOI.org/10.20546/ijcmas.2018.705.035al.

[75] Utchariyakiat I, Surassmo S, Jaturanpinyo M, et al. Efficacy of cinnamon bark oil and cinnamaldehyde on anti-multidrug resistant *Pseudomonas aeruginosa* and the synergistic effects in combination with other antimicrobial agents. *BMC Complement Altern Med* 16. 2016;158. DOI.org/10.1186/s12906-016-1134-9. PMID: 27245046; PMCID: PMC4888607.

[76] Altun M, Yapici BM. Determination of chemical compositions and antibacterial effects of selected essential oils against human pathogenic strains. *Annals of the Brazilian Academy of Sciences*. 2022;94(1). DOI 10.1590/0001-3765202220210074. PMID: 35293514.

[77] Suppakul P, Sanla-Ead N, Phoopuritham P. Antimicrobial and antioxidant activities of betel oil. *Kasetsart Journal - Natural Science*. 2006;40.91-100. Available:https://www.researchgate.net/publication/233727986_Antimicrobial_and_antioxidant_activities_of_betel_oil.

[78] Longbottom CJ, Carson CF, Hammer KA, Mee BJ, Riley TV. Tolerance of *Pseudomonas aeruginosa* to *Melaleuca alternifolia* (tea tree) oil is associated with the outer membrane and energy-dependent cellular processes. *Journal of Antimicrobial Chemotherapy*. 2004;54(2):386–392. DOI.org/10.1093/jac/dkh359. Epub 2004 Jul 14. PMID: 15254026.

[79] Papadopoulos CJ, Carson CF, Hammer KA, Riley TV. Susceptibility of pseudomonads to *Melaleuca alternifolia* (tea tree) oil and components. *J Antimicrob Chemother*. 2006;58(2):449-51. DOI: 10.1093/jac/dkl200. Epub 2006 May 30. PMID: 16735435.

[80] Mann CM, Cox SD, Markham JL. The outer membrane of *Pseudomonas aeruginosa* NCTC 6749 contributes to its tolerance to the essential oil of *Melaleuca alternifolia* (tea tree oil). *Lett Appl Microbiol*. 2000;30(4):294-7. DOI: 10.1046/j.1472-765x.2000.00712.x. PMID: 10792649.

[81] Carson CF, Papadopoulos CJ, Hammer KA, Riley TV. Tolerance of *Pseudomonas aeruginosa* to Tea Tree Oil. A report for the Rural Industries Research and Development Corporation. Publication No. 05/126 Project No. UWA-79A. 2005.

Available: <https://agrifutures.com.au/wp-content/uploads/publications/05-126.pdf>.

UNDER PEER REVIEW