

## Original Research Article

### **Influence of Sulphur and Zinc on Growth and Yield of Pearl millet (*Pennisetum glaucum* L.)**

#### **ABSTRACT**

A field experiment was conducted during the Zaid season of 2022 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) India. To study the response of Sulphur and zinc on the growth and yield of Pearl millet. The treatments consist of Sulphur @ 20, 30, and 40 kg/ha and zinc @ 5, 10, and 15 kg/ha. There were 9 treatments each replicated thrice. The soil of the experimental plot was sandy loamy in texture, nearly neutral in soil reaction (pH 7.8), low in organic carbon (0.35%), available N (163.42 kg/ha), available P (21.96 kg/ha) and available K (256.48 kg/ha). Results revealed that the higher plant height (198.06 cm), plant dry weight (75.83 g/plant), crop growth rate (47.70 g/m<sup>2</sup>/day), ear head length (25.2 cm), grains/ear head (2006.0), test weight (8.57 gm), higher grain yield (40.95 q/ha) and higher stover yield (40.95 q/ha) were significantly influenced with the application of Sulphur 40 kg/ha + Zinc 15 kg/ha. Higher gross returns (INR 92145.00/ha), higher net returns (INR 62365.00/ha) and a higher B:Cratio (2.09) were also recorded in treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha).

**Keywords:** Pearl millet, Sulphur, zinc, growth parameters and yield attributes.

#### **INTRODUCTION**

Pearl millet is commonly known in India as Bajri or Bajra. Pearl millet (*Pennisetum glaucum* L.) is also known as 'bullrush millet', originated in tropical western Africa, where the greatest number of both wild ancestors and cultivated forms occur. It belongs to the family Gramineae (Poaceae). Pearl millet is a staple diet of the rural population and also an important fodder crop in arid and semi-arid regions of India. The realized productivity of pearl millet is below its potential. The main reasons for poor crop yield are low moisture availability to crops during the growing season and lack of proper nutrient management (Choudhary *et al.*

Formatted: Indent: Left: 0.75"

Comment [AE1]: Incomplete sentence

Comment [AE2]: Its above near neutral range... slightly alkaline would be better term to use here

Comment [AE3]: In continue sentence it seems you want to say that available N P K are also low like organic carbon. But values shows only OC and N are in low range whereas P and K are in medium range. So rewrite the sentence with clear meanings.

Comment [AE4]: 3 – 4 reference need to including in introduction relevant to the contents

Formatted: Left

Formatted: Right: 0.11", Space Before: 8.95 pt

2016). The foodgrain demand of in India will increase to about 291 m tons by 2025 and to 377 m tons by 2050 (Amarasinghetal., 2010). Since, there is limited scope to increase the net cultivated area (142 million ha), the improved per unit area productivity could trigger an overall increase in food grain production (Reddy et al., 2022). Pearl millet is a heavy nutrient feeder and leads to a large withdrawal of plant nutrients from the soil. This depletion will result in a decline in the yield of the crop.

Comment [AE5]: In reference its year is 2014

Comment [AE6]: Missing in citation

Among various nutrients, sulphur and zinc play a crucial role in pearl millet production (Vinay Singhetal., 2015). Sulphur performs important functions in pearl millet. It is the best known for its role in the synthesis of proteins, oils and vitamins in pearl millet. Sulphur is associated with the production of crops for superior nutritional and market-quality produce (Chaudhary et al., 2014). Sulphur plays an important role in the formation of S-containing amino-acids, which act as building blocks in the synthesis of proteins. It has to play in increasing chlorophyll formation and aiding photosynthesis in pearl millet (Pandey et al., 2018).

Comment [AE7]: Correct format both here and in citations

Zinc plays a key role as a structural constituent or regulatory cofactor of a wide range of different enzymes and proteins in many important biochemical pathways. These are mainly concerned with carbohydrate metabolism, both in the conversion of sugars to starch, protein metabolism, auxin (growth regulator) metabolism, pollen formation the maintenance of the integrity of biological membranes, the resistance to infection by certain pathogens. Zinc deficiency is a common phenomenon in cereals, particularly in coarse treatment, soil semi-arid regions. Balanced fertilization is the key to achieve higher productivity and nutrient use efficiency (Vinay Singhetal., 2015).

Comment [AE8]: Same as above

Formatted: Font: Not Bold

Keeping these points in view, the present investigation regarding titled “Effect of S sulphur and Z zinc on G growth and Y yield of pearl millet (*Pennisetum glaucum* L.)” was conducted during Zaid-2022, at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences (SHUATS), Prayagraj (U.P.) India crop research farm, SHUATS, Prayagraj (U.P.).

Comment [AE9]: Elaborate the methodology e.g.

1. plot sizes
2. row to row
3. plant to plant distances
4. agronomic practices
5. NPK application...
6. Explain statistical analysis briefly.....

## MATERIALS AND METHODS

A field experiment was conducted during the Zaid season of 2021-22 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University

of Agriculture Technology and Sciences (SHUATS), Prayagraj (U.P.) India. The soil of the experimental plot was sandy loamy in texture, nearly neutral slightly alkaline in soil reaction (pH 7.8), and low inorganic carbon (0.35%). The treatments consist of Sulphur 20 kg/ha + zinc 5 kg/ha, Sulphur 20 kg/ha + zinc 10 kg/ha, Sulphur 20 kg/ha + zinc 15 kg/ha, Sulphur 30 kg/ha + zinc 5 kg/ha, Sulphur 30 kg/ha + zinc 10 kg/ha, Sulphur 30 kg/ha + zinc 15 kg/ha, Sulphur 40 kg/ha + zinc 5 kg/ha, Sulphur 40 kg/ha + zinc 10 kg/ha, Sulphur 40 kg/ha + zinc 15 kg/ha. The experiment was laid out in Randomized Block Design, with 9 treatments replicated thrice. The observations were recorded for plant height, plant dry weight, Crop Growth Rate ( $\text{g/m}^2/\text{day}$ ), Relative Growth Rate ( $\text{g/g/day}$ ), Ear head length (cm), Grains/ear head (g), Test weight (g), Grain yield ( $\text{kg/ha}$ ) and Stover yield ( $\text{kg/ha}$ ). The data were subjected to statistical analysis by analysis of variance method (Gomez and Gomez, 1976).

## RESULT AND DISCUSSION

### GROWTH PARAMETERS

**Plant height (cm)**- At 80 DAS, the significantly higher plant height (198.06 cm) [Table-1] was observed in treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). However, treatment-8 (Sulphur 40 kg/ha + zinc 10 kg/ha) was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). The significantly higher plant height was observed with the application of Sulphur 40 kg/ha. This increase may be due to the involvement of Sulphur in the biosynthesis of Indole 3 acetic acid. Also, Sulphur plays an important role in the growth and development of crops. It owes an important role in the formation of S-containing amino acids like cystine (27% S), Cysteine (26% S), and methionine (21% S), which act as building blocks in the synthesis of proteins. Similar results were reported by Chauhan *et al.*, (2017). Further increase of plant height might be due to the application of zinc 15 kg/ha, due to the role of zinc as a "catalyst" in most physiological, metabolic, and tryptophan synthesis processes. Certain protein elements are required to produce growth hormones (auxins) including IAA. There were similar findings described by Reddy *et al.* (2022).

**Plant dry weight (g/plant)** - At 80 DAS, the significantly higher plant dry weight (75.83 gm/plant) [Table-1] was observed in treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). However, treatment-8 (Sulphur 40 kg/ha + zinc 10 kg/ha) was found to

**Comment [AE10]:** Reviewers comment... not addressed... Discussion Paragraph? This manuscript does not discuss the results obtained, it does not explain the reasons for the results and the relationships, therefore, I suggest the authors add a couple of discussion paragraphs with other papers.

**Comment [AE11]:** Missing in citation

best statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). ~~The~~ A significantly higher plant dry weight (75.83 gm) was observed with the application of Sulphur 40 kg/ha. It has a role ~~in play~~ in increasing chlorophyll formation and aiding photosynthesis. Sulphur plays a role in the activation of enzymes ~~and~~ nucleic acids and forms parts of biotin and thiamine. Further increase in the dry weight might be due to the application of zinc at 15 kg/ha. Zinc is essential for promoting certain metabolic reactions. It is necessary for the production of chlorophyll and carbohydrates. Zinc is directly or indirectly required by several enzymes, auxin and protein synthesis. The nutrients applied in one crop are not fully utilized which leads to their residual effect on succeeding crops. Similar results conformity with [Kkaushal and Vvinay \(2021\)](#).

Comment [AE12]: Correct the format in citation

**Crop growth rate (g/m<sup>2</sup>/day)** - At 60-80 DAS, the significantly higher crop growth rate (47.70 g/m<sup>2</sup>/day) [Table-1] was observed in treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). However, treatment-8 (Sulphur 40 kg/ha + zinc 10 kg/ha) was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). ~~the~~ The significantly higher crop growth rate was observed with the ~~Sulphur sulphur~~ application 40 kg/ha, may be due to the ~~Increase increase~~ in plant height with an increase in ~~Sulphur sulphur~~ level might be due to the beneficial effect of Sulphur on the various metabolic activities and also because of its important role in cell division, photosynthetic process and formation of chlorophyll in the leaf. Therefore, it shows a positive reaction ~~on to~~ crop growth rate. Similar results ~~were reported by~~ ~~in accordance with~~ [Dadhichand Gupta \(2003\)](#).

## YIELD ATTRIBUTES

**Ear head length (cm)** - The significant and higher ear head length (25.2 cm) [Table-2] was observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 15 kg/ha), which was significantly superior over ~~the~~ rest of the treatments. However, treatment-8 (Sulphur 40 kg/ha + Zinc 10 kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). ~~The~~ A significant and higher ear head length was observed with the application of zinc 15 kg/ha. Zn ~~are is~~ involved in cell division, ~~and~~ enzyme activation and with their increased supply, their availability, acquisition, mobilization and influx into the plant tissue increased and thus improved growth attributes and yield components. Similar results ~~were~~ ~~observed by in accordance with~~ [Vvinay and Mmamta \(2018\)](#).

Comment [AE13]: Correct the format in citation

**No. of Grains/ear head** - The significant and higher Grains/ear head (2006.0) [Table-2] was observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 15 kg/ha), which was significantly superior over the rest of the treatments. However, treatment-8 (Sulphur 40kg/ha + Zinc 10 kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40kg/ha+Zinc15kg/ha). This significant and higher Grains/ear head was observed might be due to Sulphur application. Because Sulphur is a part of the amino acid (Cystine) which helps in chlorophyll formation, photosynthetic process, activation of enzymes and seed formation. Similar results were observed by **Degraet al. (2008)**.

**Test Weight** - The significantly higher Test weight (8.57 gm) [Table-2] was observed in treatment-9 with (Sulphur 40kg/ha+Zinc15kg/ha) treatments. However, treatment-8 (Sulphur 40 kg/ha + Zinc 10 kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). The significant and higher test weight was recorded with the application of Zinc (15kg/ha), which might be due to the application of zinc might have increased the photosynthetic efficiency due to improved enzymatic activity and thus might have increased the thousand grains weight. Similar results are in conformity with the findings of **Arshad et al. (2016)**.

**Grain yield** - The significant ~~ly~~ ~~Maximum~~ ~~maximum~~ Grain yield (40.95 q/ha) [Table-2] was observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 15 kg/ha), which was significantly superior over the rest of the treatments. However, treatment-8 (Sulphur 40kg/ha+Zinc10kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). The significant and higher Grain yield was observed with increasing the Sulphur application level. And also, it performs important functions in pearl millet. It is the best known for its role in the synthesis of proteins, oils and vitamins in pearl millet. Sulphur is associated with the production of crops for superior nutritional and market quality produce (**Chaudhary et al., 2014**). Further, with the application of Zinc (15 kg/ha), might be due to the greater photosynthesis efficiency or more nutrients availability due to the increasing decomposition rate of organic matter or improved individual plant performance might be the possible reasons for higher grain yield in zinc applied plots compared to other plots. These results are in conformity with the findings of **Arshad et al. (2016)**, **Norwood (1992)** and **Jan et al. (2013)**.

**Stover yield** - The significant ~~ly~~ ~~Maximum~~ ~~maximum~~ stover yield (65.33 q/ha)

Formatted: Indent: Left: 0.94", First line: 0.35"

[Table-2] was observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 15 kg/ha), which was significantly superior over the rest of the treatments. However, treatment-8 (Sulphur 40 kg/ha + Zinc 10 kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 15 kg/ha). The A significant and higher stover yield was observed by the application of Sulphur 40 kg/ha. Sulphur plays a vital role in proving vegetative structure for nutrient absorption, increasing sink strength through the development of reproductive structures by the production of assimilates to fill economically important sinks (Choudhary *et al.*, 2016). Further increase of stover yield may be due to the application of zinc 15 kg/ha. Zinc is critical to the growth and development of tryptophan, a necessary amino acid for plant growth and development. Similar results were in conformity with Reddy *et al.* (2022).

Comment [AE14]: missing

The influence of sulfur and zinc on the growth and yield of pearl millet has been a topic of research interest in recent years. Sulfur and zinc are essential micronutrients for plant growth and development, and their deficiency can result in reduced crop yields (Olivares, 2016; Araya-Alman *et al.* 2020; Olivares, 2022; Olivares *et al.* 2022a; 2022b). Several studies have investigated the effects of sulfur and zinc on pearl millet growth and yield, and these studies have provided valuable insights into the importance of these micronutrients for crop productivity.

The influence of agro-environmental factors on tropical crops in Latin America is well known. These factors include soil fertility (Hernandez *et al.* 2018; Hernandez and Olivares, 2019; Lobo *et al.* 2023), water availability (Parra *et al.* 2017; Cortez *et al.* 2017), temperature, and humidity (Olivares, 2018; Hernandez *et al.* 2020). In this study, the soil was deficient in both sulfur and zinc, which could have affected the growth and yield of pearl millet. The regular watering of the plants could have compensated for the water availability factor. However, the effect of temperature and humidity on the growth and yield of pearl millet was not studied in this experiment.

## CONCLUSION

It was concluded that with the application of Sulphur 40 kg/ha along with the zinc 15 kg/ha (Treatment-9), has performed positively and improves growth and yield parameters. Maximum grain yield, gross returns, net returns and benefit

benefit-cost ratio were also recorded with the application of Sulphur 40 kg/ha along with the zinc 15 kg/ha (Treatment-9). These findings are based on one season therefore; further trials may be required for further confirmation.

#### REFERENCES

Amarasinghe UA, Shah T, Singh OP (2010). Changing consumption patterns: Implications on food and water demand in India, International Water Management Institute, Colombo, Sri Lanka. *IWMIResearchReport.*;119:43.

Anonymous(2011). Area and production of pearl millet in India. Agricultural Statistics, Directorate of Economics & Statistics, Govt. of India.

[Araya-Alman, M., Olivares, B., Acevedo-Opazo, C. et al. \(2020\). Relationship Between Soil Properties and Banana Productivity in the Two Main Cultivation Areas in Venezuela. J Soil Sci Plant Nutr. 20 \(3\): 2512-2524. <https://doi.org/10.1007/s42729-020-00317-8>](#)

Arshad, Muhammad & Adnan, Muhammad & Ahmed, Sher & Khan, Abdul & Ali, Irshad & Ali, Muhammad & Ali, Azaz & Khan, Azam & Kamal, Muhammad & Gul, Farhana & Khan, Muhammad. (2016). Integrated Effect of Phosphorus and Zinc on Wheat Crop. *American-Eurasian J. Agric. & Environ. Sci.*, **16**:455-459.

Chaudhary, N.N., Khaif, H.R., Raj, A.D., Yadav, V. and Yadav, P. (2014). Effect of nutrients (K and S) on growth, yield and economics of pearl millet [*Pennisetum glaucum (L.)*]. *International Journal of Forestry and Crop Improvements*, **5**(1):9-12.

Chauhan T.M., Ram Lakhan and Vinay Singh (2017). Effect of potassium and sulphur on yield of and nutrient uptake by pearl millet (*Pennisetum glaucum*) in alluvial soil. *Annals of Plant and Soil Research* 19(4): 434-437.

[Cortez A, Olivares B, Parra R, Lobo D, Rodríguez M.F, Rev JC. \(2017\). Evaluation of agricultural vulnerability to drought weather in different locations of Venezuela. Rev. Fac. Agron. \(LUZ\); 34 \(1\): 103-129. <https://n9.cl/d827w>](#)

Dadhich, L. K. and Gupta, A. K. (2003). Productivity and economics of pearl millet fodder as influenced by sulphur, zinc and planting pattern. *Forage Research*. **28**(4):207-209.

Gomez, K.A., Gomez, A. A., (1976) Three or more factor experiment. (In:) *Statistical Procedure for Agricultural Research 2nd ed.*, 1976, pp.139-141.

[Hernández, R., Olivares, B. \(2019\). Ecoterritorial sectorization for the sustainable agricultural](#)

**Comment [AE15]:** Extension revision of references format is required. All references should be in journals acceptable format. In text I have found many references which are not listed in this section. Address missing references problem.

**Formatted:** Portuguese (Brazil)

**Formatted:** Portuguese (Brazil)

[production of potato \(\*Solanum tuberosum\* L.\) in Carabobo, Venezuela. \*Agricultural Science and Technology\*. 20\(2\): 339-354. \[https://doi.org/10.21930/rcta.vol20\\\_num2\\\_art:1462\]\(https://doi.org/10.21930/rcta.vol20\_num2\_art:1462\)](#)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

[Hernández, R; Olivares, B., Coelho, R., Molina, J.C., Pereira, Y. \(2018\). Analysis of climate types: Main strategies for sustainable decisions in agricultural areas of Carabobo, Venezuela. \*Scientia Agropecuaria\*; 9\(3\): 359 – 369. <https://doi.org/10.17268/sci.agropecu.2018.03.07>](#)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

[Hernandez, R.; Olivares, B.; Arias, A; Molina, J.C., Pereira, Y. \(2020\). Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. \*Idesia\*, 38\(2\):95-102. <http://dx.doi.org/10.4067/S0718-34292020000200095>](#)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Kaushal and [vinayVinay](#) (2021). Effect of sulphur and zinc management on yield, quality and nutrient uptake in pearl millet (*Pennisetum glaucum*) – lentil (*Lens culinaris*) crop sequence. *Annals of Plant and Soil Research* 23(2):135-139.

[Lobo, D; Olivares, B; Rey, J.C; Vega, A; Rueda-Calderón, A. \(2023\). Relationships between the Visual Evaluation of Soil Structure \(VESS\) and soil properties in agriculture: A meta-analysis. \*Scientia agropecuaria\*, 14-1, 67- 78. <https://doi.org/10.17268/sci.agropecu.2023.007>](#)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

[Mukesh C., K.S. Rana, D.S. Rana and R.S. Bana \(2016\). Tillage and crop residue effects in rainfed pearl millet \(\*Pennisetum glaucum\*\) in conjunction with sulphur fertilization under pearl millet-Indian mustard. \*Indian Journal of Agronomy\* 61 \(1\):1519.](#)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: Indent: Left: 0"

[Olivares, B. \(2016\). Description of soil management in agricultural production systems of sector hammock in Anzoátegui, Venezuela. \*La Granja: Revista de Ciencias de la Vida\*. 23\(1\): 14-24. <https://doi.org/10.17163/lgr.n23.2016.02>](#)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

[Olivares, B. \(2022\). Determination of the potential influence of soil in the differentiation of productivity and in the classification of susceptible areas to banana wilt in Venezuela. UCOPress: Spain, pp. 89-111, <https://helvia.uco.es/handle/10396/22355>](#)

[Olivares B. \(2018\). Tropical conditions of seasonal rain in the dry-land agriculture of Carabobo, Venezuela. \*La Granja: Journal of Life Sciences\*; 27\(1\):86-102. <http://doi.org/10.17163/lgr.n27.2018.07>](#)

[Olivares, B.O.; Rey, J.C.; Perichi, G.; Lobo, D. \(2022a\). Relationship of Microbial Activity with Soil Properties in Banana Plantations in Venezuela. \*Sustainability\* 14, 13531. <https://doi.org/10.3390/su142013531>](#)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Olivares B, Vega A, Calderón MAR, Rey JC, Lobo D, Gómez JA, Landa BB. (2022b). Identification of Soil Properties Associated with the Incidence of Banana Wilt Using Supervised Methods. Plants, 11(15):2070. <https://doi.org/10.3390/plants11152070>

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Parra R, Olivares B, Cortez, A. (2017). Characterization of precipitation patterns in Anzoátegui state, Venezuela. Ería: 3 (3): 353-365. <https://doi.org/10.17811/er.3.2017.353-365>

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Pandey, M., Abidi, A. B. and Singh, R. P. (2010). Effect of sulphur and phosphorus on yield and quality of pearl millet (*Pennisetum glaucum* L.). *Annals of Plant and Soil Research*, **20**(3):272-275.

~~Chauhan T. M., Ram Lakhan and Vinay Singh (2017). Effect of potassium and sulphur on yield and nutrient uptake by pearl millet (*Pennisetum glaucum*) in alluvial soil. *Annals of Plant and Soil Research* **19**(4):434-437.~~

Vangala S, Rajesh and Chandu (2022). Effect of phosphorus and zinc on growth and yield of pearl millet (*Pennisetum glaucum* L.). *The Pharma Innovation Journal*; **11**(4):542-545.

Vinay and Mamta (2018). Direct effect of sulphur and zinc on productivity, quality and nutrient uptake by pearl millet (*Pennisetum glaucum*) and their residual effect on succeeding wheat (*Triticum aestivum*) in pearl millet – wheat crop sequence. *Annals of Plant and Soil Research* **20**(3):233–238.

Vinay, Harvendra, Seema, Javed and Jagpal (2015). Balanced use of nutrients for sustaining higher production of pearl millet in alluvial soil. *Annals of Plant and Soil Research* **17**(4):346-349.

UNDER PEER REVIEW

**Table-1. Effect of sulphur and zinc on the growth of pearl millet.**

**Comment [AE16]:** It is more appropriate to add letters which show the significant difference among treatments.

S. No.	Treatment combinations	Plant height (cm)	Dry weight (g/plant)	Crop growth rate (g/m <sup>2</sup> /day)
1.	Sulphur 20kg/ha + Zinc 5 kg/ha	180.38	62.90	36.73
2.	Sulphur 20kg/ha + Zinc 10 kg/ha	183.53	64.22	37.31
3.	Sulphur 20kg/ha + Zinc 15 kg/ha	184.62	65.87	36.65
4.	Sulphur 30kg/ha + Zinc 5 kg/ha	186.38	67.64	39.20
5.	Sulphur 30kg/ha + Zinc 10 kg/ha	188.94	68.22	40.22
6.	Sulphur 30kg/ha + Zinc 15 kg/ha	189.96	68.95	38.93
7.	Sulphur 40kg/ha + Zinc 5 kg/ha	192.27	71.20	41.65
8.	Sulphur 40kg/ha + Zinc 10 kg/ha	195.05	73.87	42.90
9.	Sulphur 40kg/ha + Zinc 15 kg/ha	198.06	75.83	47.70
	Ftest	S	S	S
	SEm.(±)	1.12	0.84	1.71
	CD (p=0.05)	3.37	2.53	5.12

Comment [AE17]: It is more appropriate to add letters which show the significant difference among treatments.

**Table-2. Effect of sulphur and zinc on yield of pearl millet.**

S. No.	Treatment combinations	Ear head length (cm)	No. of Grains/earhead	Test weight (g)	Grain yield (q/ha)	Stover yield (q/ha)	Harvest index (%)
1.	Sulphur 20kg/ha + Zinc 5 kg/ha	19.8	1837.1	6.75	32.29	47.55	40.44
2.	Sulphur 20kg/ha + Zinc 10 kg/ha	20.8	1876.0	6.81	33.94	48.20	41.32
3.	Sulphur 20kg/ha + Zinc 15 kg/ha	21.5	1902.4	7.19	34.61	49.11	41.34
4.	Sulphur 30kg/ha + Zinc 5 kg/ha	21.3	1911.1	7.45	35.24	50.99	40.87
5.	Sulphur 30kg/ha + Zinc 10 kg/ha	22.4	1929.6	7.81	36.16	54.03	40.10
6.	Sulphur 30kg/ha + Zinc 15 kg/ha	22.7	1938.4	7.99	37.03	58.36	38.83
7.	Sulphur 40kg/ha + Zinc 5 kg/ha	23.5	1965.3	8.10	37.82	59.33	38.93
8.	Sulphur 40kg/ha + Zinc 10 kg/ha	24.5	1981.1	8.31	39.03	63.32	38.14
9.	Sulphur 40kg/ha + Zinc 15 kg/ha	25.2	2006.0	8.57	40.95	65.33	38.53
	Ftest	S	S	S	S	S	S
	Sem(±)	0.27	6.07	0.13	0.40	0.65	0.41
	CD(P=0.05)	0.81	18.19	0.40	1.21	1.94	1.23