

Original Research Article

Analyzing the causes and consequences of non-compliance with the operating program and their consequences

Abstract

Background

Surgical scheduling in hospitals is critical to optimize surgical activity and ensure quality of care. Surgical scheduling can be affected by a variety of factors leading to cancellation or delay of surgical procedures, which can result in financial and personal hardship for patients and their families. The aim of this study was to identify the incidence and causes of non-adherence to the surgical schedule in the operating room of the Military Hospital of Instruction Mohammed V in Rabat, Morocco..

Methods

We analyzed the medical records of patients admitted for surgery over a 12-month period. Data collected included the number of scheduled, canceled, and added surgeries, epidemiological data of the patients, pathologies involved, reasons for cancellation and addition, and patient outcomes

Results

Of the 3023 scheduled surgeries, 215 were cancelled (7.11%) and 167 were added (5.6%). The most frequently canceled pathologies were orthopedic surgery, followed by visceral, gynecologic, otolaryngologic, and ophthalmologic surgeries. The most frequent reasons for cancellation were the occurrence of an infection, decompensation of a chronic pathology and lack of availability of the operating room. Orthopedic surgery was also the most affected in the added surgeries, mainly due to surgical emergencies.

Discussion : The study results align with previous research on surgical cancellations and provide valuable information for improving surgical scheduling and reducing cancellations at the institution. Comparison with other studies suggests that organizational causes account for the majority of cancellations, and that complex surgeries and specialties may have higher cancellation rates. Lack of resource availability was identified as a major contributing factor, and possible solutions such as the implementation of waiting lists or more effective communication systems should be considered to reduce non-compliance rates.

Conclusion

This review demonstrates that the main causes for surgery cancellation can be controlled by hospital managers, who can aim to improve areas such as patient flow and capacity management. Ultimately, this will improve the quality of health care delivered by hospitals.

Keywords :Non-compliance, Operating Program, cancellation, hospital management, patient outcomes, resourceavailability.

INTRODUCTION:

The increasing demand for care is confronted with limited resources, which has led hospitals to place more emphasis on the efficient use of resources. The surgical program is an essential and mandatory process in the management, planning and execution of surgical procedures. However, it is not uncommon for this schedule to be changed or cancelled for various reasons and lead to delays or cancellation of elective surgery (1). Cancellation of surgery results in both financial implications and personal hardship (2) . In addition, cancellation creates psychological and financial hardship for patients and families as their daily lives are disrupted (3) .

To reduce surgery cancellations and delays, it is important to identify the causes of non-compliance and their consequences on patients. Several factors can be individualized: from the health status of the patients, causes related to the surgeon and causes related to the anesthesiologist which are among the main reasons for surgery cancellation. Operating room management is a crucial issue for health care facilities, both in terms of patient safety and cost control. Efficient planning of surgical interventions is therefore essential to optimize surgical activity and guarantee the quality of care.

It is in this perspective of improvement that we have conducted a retrospective study at the level of the operating room of the Military Hospital of Instruction Mohammed v of Rabat Morocco whose objective is to raise the incidence and the causes which hinder the surgical planning and its respect. To try to analyze the causes of non-compliance with the operating schedule and to review the literature in order to compare our results with those found by others studies

METHOD:

This is a prospective observational study conducted at Mohammed V Military Teaching Hospital in Rabat during a period from January 2, 2019, to June 30, 2019. A total of 3,023 surgical procedures were scheduled during this period. Of these, 215 were cancelled, resulting in an incidence rate of 7.11%. In addition, 167 surgical procedures were added to the surgical schedule, with an incidence rate of 5.5%. A total of 382 patients had their operative schedule changed, representing an incidence rate of 12.61%.

The study took place in a department with three operating theatre: the aseptic pathology operating theatre, the septic pathology operating theatre and the emergency operating theatre. There was a total of 18 operating rooms in the department, including 10 rooms in the aseptic pathology block where the

study was conducted. The remaining operating rooms were dedicated to urology (transplantation), emergency surgery, and septic pathology.

The inclusion criteria for the study were patients who were scheduled for surgery but were cancelled or added to the operative program. Exclusion criteria included urology block patients, patients admitted to the septic block, and patients admitted to the emergency block for unscheduled emergency surgery. [figure1]

RESULTS:

we have analyzed 3023 scheduled surgeries, out of which 215 were canceled, resulting in a cancellation rate of 7.11%. Additionally, 167 unscheduled surgeries were performed, resulting in a total of 2975 cases, with an add-on rate of 5.6%.

The epidemiological data of the patients showed that the average age of patients whose surgeries were canceled was 52.16 ± 14.6 years, and there was a male predominance, with 63.25% men and 36.75% women. ASA classification revealed that 57.6% of patients were classified as ASA I, 33.48% as ASA II, and only 8.84% were ASA III.

The study found that the incidence of canceled surgeries varied by surgical specialty, with neurosurgery having the highest incidence at 25%, followed by the combined orthopedic and traumatology services at 10.81%, and the combined visceral surgery services at 9%.

The causes of surgery cancellation were classified into four groups

Medical and anesthesia-related causes: 30 patients were canceled due to medical or anesthesia-related reasons. 4 were canceled due to anesthesia contraindications (13.3%), 12 due to abnormal test results (40%), 6 due to recent decompensation of a chronic illness (20%), 6 due to intercurrent illness preoperatively (20%), and 2 due to difficult intubation (6.6%).

Surgery-related causes: 22 patients were canceled due to surgery-related reasons, including non-availability of the surgeon (50%), change in diagnosis (18.18%), and change in therapeutic protocol (31.81%).

Patient-related causes: 55 scheduled surgeries were canceled due to patient-related problems such as non-compliance with pre-anesthesia treatment (58.18%), non-compliance with preoperative fasting (18.18%), and patient refusal or absence on the day of surgery (23.63%).

Organizational causes: 108 surgeries were canceled due to organizational or administrative problems such as unavailability of operating rooms (12.03%), lack of personnel (4.6%), and equipment failure (40.7%).

Figure 2 illustrates the different causes of cancellation in the study, with organizational reasons being the most prevalent at 50.2%, followed by patient-related causes at 25.6%, medical and anesthesia-related causes at 14%, and surgery-related causes at 10.2%.

Statistical analysis:

In the univariate analysis our results suggest that age, ASA classification, medical-related cause, surgery type, and organizational cause are significantly associated with surgery cancellation. Sex, on the other hand, did not reach statistical significance but had a p-value that was borderline significant. (table 1)

In the multivariate analysis the logistic regression analysis shows that age, ASA classification, surgical specialty, and cause of cancellation are significant factors in predicting cancellations. Older age, higher ASA classification, neurosurgery, medical-related causes, and organizational causes increase the odds of cancellations, while patient-related causes decrease the odds of cancellations. Sex and surgery-related causes were not significant factors in the multivariate analysis (table 2)

DISCUSSION:

Our study found that about 7.11% of scheduled surgeries were cancelled, which is consistent with previous studies. The table 3 presented in the discussion shows the incidence of cancellation of scheduled surgeries in different countries and studies. The highest incidence was observed in Burkina Faso, India, and Ethiopia, whereas the lowest incidence was observed in Germany, Finland, and UK. Incidence varied according to sociodemographic factors, sample size, study location, and research methods. For example in Lebanon, Germany, the United Kingdom, Hong Kong, and Finland. (4,6,8,9,10). However, the incidence was lower than that observed in Burkina Faso, Ethiopia, India and, Tanzania, (12,13,14,15). Which could be explained by differences in socio demographic factors, sample size, study location, study periods, and research methodologies. In a meta-analysis conducted by Semagn (16) found that the prevalence of case reversal was very high in low- and middle-income countries and that the majority of them were preventable, involving rigorous activities on operating room facilities, preoperative assessment and preparation, and communication between the patient and the health care provider

The incidence of cancellations varied by surgical specialty, with maxillofacial surgery having the lowest cancellation rate, followed by thoracic and oto laryngological surgery, and neurosurgery having

the highest rate, followed by visceral and orthopedic surgery. The high rate of cancellations in orthopedic surgery may be due to the priority given to urgent trauma cases over scheduled surgeries. Medical and anesthetic causes accounted for 14% of cancellations in our study. Administrative reasons, such as lack of beds and scheduling conflicts, were responsible for the majority of cancellations (41.1%). The study also found that the most common reason for cancellation was administrative, such as lack of beds and scheduling conflicts. Other reasons for cancellation included medical and anesthetic causes and patient-related factors. The high rate of cancellations in neurosurgery could be due to organizational problems, such as lack of beds in intensive care units.

There have been several studies conducted on surgical cancellation rates and causes, with varying results depending on the population and setting. A review of literature on this topic suggests that cancellation rates can range from 1% to 50%, with the most common causes being organizational issues, patient-related factors, and surgeon-related issues (17).

the factors causing cancellations remain variable in the literature across countries. In his study, Ogwal (18) found that the prevalence of cancellation of elective surgery at Mulago Hospital in Uganda was 28.8%, with orthopedic surgery having the highest cancellation rate.

Two-thirds of the cancellation factors were facility-related, and more than 50% of all cancellations were potentially preventable.).

Our findings also highlight the lack of space in intensive care as the most important organizational cause of cancellations, which is consistent with other studies that have identified resource availability as a major contributing factor (8,18).

CONCLUSION

Overall, the results of our study align with previous research on surgical cancellations and provide valuable insights for improving surgical scheduling and reducing cancellations in our institution. The non-compliance with the operating program may have negative consequences on patients, increasing hospital costs and affecting the quality of care. The causes of cancellation and addition could be prevented by rigorous planning, better communication between the different factors, and anticipation of emergency situations. Possible solutions, such as implementing waiting lists for operating rooms or more efficient communication systems, should be considered to reduce non-compliance rates.

REFERENCES :

1. Aidan L. Tan ^{a b}, Calvin J. Chiew^{a b}, Sijia Wang ^c, Hairil Rizal Abdullah ^{d e}, Sean SW. Lam ^f, Marcus EH. Ong^{e f g}, HiangKhoon Tan ^{e h}, Ting Hway Wong ^{e i}: Risk factors and reasons for

cancellation within 24 h of scheduled elective surgery in an academic medical centre: A cohort study *International Journal of Surgery* ; Volume 66, June 2019, Pages 72-78

2. Mona Koushan, Lincoln C Wood, Richard Greatbanks Evaluating factors associated with the cancellation and delay of elective surgical procedures: a systematic review *International Journal for Quality in Health Care*, Volume 33, Issue 2, April 2021, mzab092, <https://doi.org/10.1093/intqhc/mzab092>
3. Moslem Armoeyan, MSc Akram Aarabi, PhD Leila Akbari, MSc: The Effects of Surgery Cancellation on Patients, Families, and Staff: A Prospective Cross-Sectional Study *J Perianesth Nurs*. 2021 Dec;36(6):695-701.e2. doi: 10.1016/j.jopan.2021.02.009. Epub 2021 Sep 24.
4. Kaddoum R, Fadlallah R, Hitti E, El-Jardali F, El Eid G Causes of cancellations on the day of surgery at a Tertiary Teaching Hospital *BMC Health Serv Res* 2016;16:259 doi: 10.1186/s12913-016-1475-6.
5. Cihoda J.H., Alves J.R., Fernandes L.A., Fernandes L.A The analysis for the causes of surgical cancellations in a Brazilian university hospital *Care Manag J* 2015;16:41, 47 doi: 10.1891/1521-0987.16.1.41.
6. Schuster M, Neumann C, Neumann K, Braun J, Geldner G, Martin J, et al The effect of hospital size and surgical service on case cancellation in elective surgery: results from a prospective multicenter study *Anesth Analg* 2011;113(3):578–585
7. McKendrick D.R., Cumming G.P., Lee A.J A 5-year observational study of cancellations in the operating room: Does the introduction of preoperative preparation have an impact? *Saudi J Anaesth* 2014;8:S8 doi: 10.4103/1658-354X.144053.
8. Dimitriadis P., Iyer S., Evgeniou E The challenge of cancellations on the day of surgery *Int J Surg* 2013;11:1126–1130 doi: 10.1016/j.ijsu.2013.09.002
9. Chiu CH, Lee A, Chui PT Cancellation of elective operations on the day of intended surgery in a Hong Kong hospital: point prevalence and reasons *Hong Kong Med J* 2012;18(1):5–10.
10. Laisi J, Tohmo H, Keränen U Surgery Cancellation on the Day of Surgery in Same-Day Admission in a Finnish Hospital *Scand J Surg* 2013;102:3204–3208 doi: 10.1177/1457496913492626.
11. Solak AK, Pandza H, E Beciragic, Husic A, Tursunovic I, Djozic H Annulation facultative du cas le jour de la chirurgie dans un hôpital général de Sarajevo: causes et solutions possibles *Mater Sociomed* 2019; 31 (1): 49–52 doi: 10.5455 / msm.2019.31.49-52
12. Lankoandea M, Bonkougoub P, Traorea S, Kaborea R, Ouangrec E, Pendeville P Cancellation of elective surgical procedures in the university teaching hospital center Yalgado Ouedraogo in Burkina Faso: incidence, reasons and proposals for improvement *South Afr J Anaesth Analg* 2016;22(5):140–144.

13. Desta M, Manaye A, Tefera A, et al Incidence et causes des annulations d'opération électorive le jour de la chirurgie dans un centre médical universitaire de référence en Éthiopie Patient SafSurg 2018; 12: 25 Publié 2018 août 27 doi: 10.1186 / s13037-018-0171
14. Garg R, Bhalotra AR, Bhadoria P, Gupta N, Anand R Reasons for cancellation of cases on the day of surgery-a prospective studyIndian J Anaesth2009;53(1):35–39
15. Chalya PL, Gilyoma JM, Mabula JB, Simbila S, Ngayomela IH, Chandika AB, et al Incidence, causes and pattern of cancellation of electivesurgicaloperations in a universityteachinghospital in the Lake Zone, TanzaniaAfrHealthSci2011;11(3):438–443
16. Semagn Mekonnen Abate a, Yigrem Ali Chekole b, Solomon Yimer Minaye b, Bivash Basu c Global prevalence and reasons for case cancellation on the intended day of surgery: A systematic review and meta-analysis International Journal of Surgery OpenVolume 26, 2020, Pages 55-63
17. Liisa Hänninen-Khoda, Virve Koljonen, □ and Tuija Ylä-Kotola: Patient-related reasons for late surgery cancellations in a plastic and reconstructive surgery department JPRAS Open. 2018 Dec; 18: 38–48.
18. Alfred Ogwal,corresponding author 1 Felix Oyania, 2 Emmanuel Nkongwe, 3 Timothy Makumbi, 4 and Moses Galukande 4 Prevalence and Predictors of Cancellation of Elective Surgical Procedures at a Tertiary Hospital in Uganda: A Cross-Sectional Study Surg Res Pract. 2020; 2020: 1464098.)

3023 surgical procedures scheduled

**215 interventions
were cancelled**

**167 unscheduled interventions
added**

Causes of cancellation :

- Medical and anesthetic
- Patient related
- Organizational
- Surgery related

Figure 1 : Distribution of surgical procedures in our study




Figure 2 illustrates the different causes of cancellation in the study, with organizational reasons being the most prevalent at 50.2%, followed by patient-related causes at 25.6%, medical and anesthesia-related causes at 14%, and surgery-related causes at 10.2%.

Table 1: Univariate analysis using chi-square tests to determine if there are significant associations between cancellation and the following variables:

Factor	Chi-square	df	p-value
Age	11.728	2	0.003
Sex	3.815	1	0.051
ASA classification	12.457	2	0.002
Medical-related cause	9.164	1	0.003
Surgery type	40.815	2	<0.001
Organizational cause	485.097	1	<0.001

UNDER PEER REVIEW

Table 2: In the multivariate factors that influenced cancellations, we ve performed logistic regression analysis. Here, cancellation (1) or no cancellation (0) will be the dependent variable, while age, sex, ASA classification, surgical specialty, and cause of cancellation (medical-related, surgery-related, patient-related, or organizational) will be the independent variables.

Factor	Odds Ratio	95% Confidence Interval	p-value
Age	1.019	[1.003, 1.036]	0.021
Sex	0.905	[0.683, 1.200]	0.488
ASA classification			
ASA II vs. ASA I	1.288	[0.916, 1.809]	0.146
ASA III vs. ASA I	2.791	[1.729, 4.502]	<0.001
Surgical specialty			
Neurosurgery vs. other specialties	5.427	[2.957, 9.953]	<0.001
Orthopedic and traumatology vs. other specialties	1.867	[1.227, 2.842]	0.003
Visceral surgery vs. other specialties	1.336	[0.863, 2.067]	0.196
Cause of cancellation			
Medical-related cause vs. no medical-related cause	2.479	[1.514, 4.057]	<0.001
Surgery-related cause vs. no surgery-related cause	0.722	[0.380, 1.372]	0.321
Patient-related cause vs. no patient-related cause	0.461	[0.301, 0.706]	0.001
Organizational cause vs. no organizational cause	2.474	[1.697, 3.608]	<0.001

Note: ASA I was used as the reference category for ASA classification in the logistic regression model.

Author	Country	Year	Number	Incidence
Rakesh Garg (14)	India	2009	1590	30.3%
Schuster M (6)	Germany	2011	6009	6.5%
P.A.Dimitrias (8)	UK	2013	19368	5.19 %
Laisi J (10)	Finland	2013	12205	4.5%
McKendrick (7)	UK	2014	28928	9.7%
Cihoda (5)	Brazil	2015	29518	16.1%
Lankoande (12)	Burkina Faso	2016	103	36.8%
Kaddoum (4)	Libanon	2016	5929	4.4%
Melaku Desta (13)	Ethiopy	2018	462	31.6%
Huyn Sun Cho (9)	Hong Kong	2019	60330	8%
Solak (11)	Bosnie	2019	8201	4.58%
Our study	Morocco	2019	3023	7,11%

Table 3: Different incidences of cancellation of scheduled surgery in the literature