

Original Research Article

“Effect of organic manures on growth and yield of rice varieties (*Oryza sativa* L.) and yield validation using SPSS model”

Abstract

The experiment was conducted during the *Kharif* season 2022, at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (U.P.) to find out the “**Effect of Organic manures on growth and yield of rice varieties (*Oryza sativa* L.) and yield validation using SPSS model.** The experiment was laid out in Randomized Block Design comprising of 9 treatments which include 3 varieties NDR-359, BPT-5204 and MTU-7029 and 3 organic manures Poultry manure 5(t/ha), Vermicompost 10(t/ha) and Poultry manure 2.5(t/ha) + Vermicompost 5(t/ha). Plant growth characters, yield attributes and yield were significantly influenced by different treatments except plant height, plant dry weight and no. of panicles per hill all the parameters were found to be highest with the treatment T9 (MTU-7029 + Poultry manure 2.5(t/ha) + Vermicompost 5(t/ha)). The economic analysis also showed that the treatment T9 maximized the net returns, gross returns and benefit-cost ratio (BCR). Treatment 9 has shown 44.66% increase over predicted yield whereas there was 29.50% increase in treatment 8 over predicted yield through SPSS model. From the basis of one year experimentation, it is concluded that Poultry manure 2.5(t/ha) + Vermicompost 5(t/ha) was found more productive for the variety MTU-7029.

Keywords: Manure, yield, poultry, vermicompost, paddy, yield validation and SPSS.

Introduction

Rice is one of the most important cereal crops in the world, feeding half of the world's population and accounting for 35–60% of all calories consumed by humans (**Tayefe et al. 2014**). Rice is derived from the cereal grass *Oryza Sativa*, a genus in the family Gramineae.

Currently, Asia accounts for more than 90% of global rice production and consumption. Rice provides 60–70% of the energy needs of more than two billion people in Asia (**Sridhar et al. 2019**).

India has the largest area (43.50 m ha) among the nations that grow rice and is the second-largest producer (163.51 MT), after China (203.14 MT), with an average productivity of 3.76 t/ha. Around 44 million ha of land in India are used for rice farming, and 109.70 million tonnes were produced there between 2016 and 2017. With a 5.98 million ha area, Uttar Pradesh produces 14.64 million tonnes of rice with a productivity of 2.447 t/ha (**GOI, 2016**).

For the country's food security, India must produce an additional 1.7 million tonnes of rice per year (**Dass and Chandra, 2013**).

Instead of artificial fertilizer, rice may be grown using the nutrient-rich poultry manure, but it must first be well mixed into the soil. Before using it in your field, you must give it time to mature. The least suggested time to mature chicken dung before putting it to a field is three to four months, and sooner than six months is more prudent. Poultry manure is an excellent fertilizer material because it contains essential plant nutrients, especially nitrogen (N), phosphorus (P), and potassium (K). It contains a lot of nitrogen. In order to replace the usage of urea on the majority of commercial farms, chicken manure can improve the N content of soil. These nutrients plus others come largely from the bird faeces. Manures decompose (mineralize) in the soil releasing nutrients for crop uptake. Poultry manure has more nutrients than other animal manures. It contains significant amounts of micronutrients such Cu, Zn, Fe, Mn, and others as well as nitrogen (3-5 %), phosphorus (1.5-3.5 %), potassium (1.5-.03 %) (**Mohamed, 2010**).

Vermicompost was employed as an organic source for the nutrients nitrogen, phosphorus, and potassium, as needed for the treatment. It has 1.0-1.5% K, 1.8-2.1% P, and 1.5-2.2% N. The term "vermicompost" refers to the compost created by the use of earthworms, and "vermiculture" or "Verm technology" refers to a contemporary method of harnessing the ecosystem for efficient utilisation of the organic waste with the aid of earthworms, which results in the emergence of useful organic manure. With this method, we can compost biodegradable waste while also utilising the compost's byproducts to boost crop production and cut down on the use of artificial fertilisers (**Sujit, 2012**).

Weather has an impact on crop development at various phenological stages, which explains why yields vary from year to year and location to location. The response of crops to weather has been measured using a variety of statistical approaches, including multiple regressions, principal component analysis (**Jain et al., 1984**), Markov chain analysis (**Ramasubramanian et al., 1999**), and agro-meteorological models (**Walker, 1989**). In India, agricultural yields were predicted using multiple regression models (**Appa Rao, 1983**). To assess yield patterns and forecast yields in various circumstances, time series analysis is utilised.

Materials and methods

The experiment was conducted during the *Kharif* season 2022, at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (U.P.) which is located at 25.4071753 N latitude, 81.8490776 E longitude and 98 m altitude above the mean sea level (MSL). This area is situated on the right side of the Yamuna River of Prayagraj - Rewa road about 12 km from the city. The soil of the experiment plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.4), low in organic carbon (0.51%), available N (78.9 kg/ha), available P (32.88 kg/ha), available K (385.10 kg/ha). Nutrient sources were Poultry manure and Vermicompost to fulfil the requirement of Nitrogen, Phosphorus, Potassium, respectively. The experiment was laid out in Randomized Block Design (RBD) with nine treatments replicated thrice. The treatments were 1- NDR-359 + Poultry manure 5(t/ha), 2- NDR-359 + Vermicompost 10(t/ha), 3- NDR-359 + Poultry manure 2.5(t/ha) + Vermicompost 5(t/ha), 4- BPT-5204 + Poultry manure 5(t/ha), 5- BPT-5204 + Vermicompost 10(t/ha), 6- BPT-5204 + Poultry manure 2.5(t/ha) + Vermicompost 5(t/ha), 7- MTU-7029 + Poultry manure 5(t/ha), 8- MTU-7029+ Vermicompost 10(t/ha), 9- MTU-7029+ Poultry manure 2.5(t/ha) + Vermicompost 5(t/ha). The growth parameters of the plants were recorded at frequent intervals from transplanting up until 100 DAT and finally, the yield parameters were recorded after harvest. Analysis of Variance (ANOVA) was used statistically to examine these variables using the Randomized Block design. The Pearson's correlation between the measured yield and the individual weather parameters as well as the combination of weather parameters was calculated using SPSS (Statistical Product and Service Solutions). The correlation coefficient has been obtained from the sum of weather parameters and the sum product of various weather parameters. The dependant variable (yield) and the independent variables (time, sum, and sum products for various meteorological conditions) were regressed many times. The regression formula was used to create the regression equation.

Result and Discussion

Plant height (cm)

The highest plant height (108.3 cm) was recorded with application of treatment 3 [NDR 359 + Poultry manure 2.5t/ha + Vermicompost 5t/ha] as compare to other treatments and statistically at par with NDR- 359 + Poultry manure 5t/ha. Whereas minimum plant height was recorded in treatment 8 [MTU-7029 + Vermicompost 10t/ha].

The increased concentration of N in poultry manure, which is also more easily available to crops, may account for the taller plants in treated plots. The findings of **Edward and Daniel (1992)**, **Suvarna Latha and Sankara Rao (2001)**, and **Santosh Kumar Jha et al. (2004)** are all in agreement with these findings. Vermicompost may perform better than FYM because of the nutrients' increased concentration and quicker release.

Number of tillers per hill

The highest number of tillers (13.1) was recorded with application of treatment 9 [MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha] as compare to other treatments and statistically at par with MTU-7029 + Poultry manure 5t/ha. Whereas minimum number of tillers was recorded in treatment 1 [NDR- 359 + Poultry manure 5t/ha].

Among the organic manure treatments, the better performance of poultry manure-treated plots might be due to the involvement of certain growth-promoting substances, which might have also accelerated the number of tillers. This is in accordance with the findings of **Prabhakaran (2000)**, and **Helen Belephant-Miller (2007)**. Better performance of vermicompost might be due to its presence of macronutrients and micronutrients (**Shinde 1992**).

In general, tiller number showed an increasing trend up to 60 DAT, and thereafter decreased up to harvest. A decrease in the number of tillers after 60 DAT could be due to the senescence of lower tillers because of the shading effect, which was also supported by **Singh's (2001)** findings. The variation in tiller production might be attributed to the availability of nutrients.

Dry weight (g/hill)

The highest Dry weight (53.32 g/hill) was recorded with application of treatment 3 [NDR 359 + Poultry manure 2.5t/ha + Vermicompost 5t/ha], there are no statistically at par values in the

treatment combination. Whereas minimum Dry weight was recorded in treatment 5 [BPT-5204 + Vermicompost 10t/ha].

Increased dry matter accumulation in plots treated with poultry manure and vermicompost may be attributed to the ongoing, slow release of nutrients that have allowed the leaf area duration to increase, giving plants the chance to increase their photosynthetic rate, which may have in turn resulted in higher dry matter accumulation. **Amanullah (2006), Suvarna Latha and Sankara Rao (2001) and Altaf Hussaine (2012)** all came to similar conclusions.

Crop growth rate (g/m²/day) and Relative growth rate (g/g/day)

The significantly maximum crop growth rate (18.63 g/m²/day) was recorded with application of treatment 8 [MTU-7029 + Vermicompost 10t/ha] as compare to other treatments and statistically at par with NDR 359 + Vermicompost 10t/ha and [BPT-5204 + Poultry manure 5t/ha]. Whereas minimum Crop growth rate was recorded in treatment 5 [BPT-5204 + Vermicompost 10t/ha].

The significantly maximum relative growth rate (0.0123 g/g/day) was recorded with application of treatment 8 [MTU-7029 + Vermicompost 10t/ha] as compare to other treatments and statistically at par with NDR 359 + Vermicompost 10t/ha and [BPT-5204 + Poultry manure 5t/ha].

The higher leaf chlorophyll content and better cellular nutrition appear to have accelerated photosynthetic activity. Therefore, it follows that the enhanced metabolic activities and photosynthetic rate that result in an improvement in the accumulation of dry matter at the successive growth stages may be the cause of the improved growth and development of the crop plants in the current investigation. This in turn leads to an increase in the crop growth rate and relative growth rate in all stages of plants. The current study takes into account the conclusions of **Das (2016)**.

Yield parameters

The highest number of effective tillers per hill was recorded in Treatment T9 (MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha) (11.33) as compare to other treatments and statistically at par with (MTU-7029 + Vermicompost 10t/ha). Whereas minimum number of effective tillers per hill are seen in Treatment T1 (NDR- 359 + Poultry manure 5t/ha).

The highest number of panicles per hill was recorded in Treatment T6 (Bpt-5204 + Poultry manure 2.5t/ha + Vermicompost 5t/ha) (9.21), there are no statistically at par values in the

treatment combination. Whereas minimum number of panicles per hill are seen in Treatment T7 (MTU-7029 + Poultry manure 5t/ha).

The highest number of grains per panicle was recorded in Treatment T9 (MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha) (112.37), there are no statistically at par values in the treatment combination. Whereas minimum number of grains per panicle are seen in Treatment T1 (NDR- 359 + Poultry manure 5t/ha).

The highest test weight was recorded in Treatment T9 (MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha) (22.35 g), there are no statistically at par values in the treatment combination. Whereas minimum test weight are seen in Treatment T5 (BPT-5204 + Vermicompost 10t/ha).

The nutrients in the poultry manure are readily available to the crop due to well-developed decomposition, which may have led to increased yield attributes such as productive tillers, number of filled grains per panicle, 1000 grain weight, and significantly lower sterility percent, which in turn led to higher yield. Researchers like **Datta (1994), Singh (2001) and Hossan (2010)** all found similar findings.

Yield (t/ha)

The highest grain yield was recorded in Treatment T9 (MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha) (5.21 t/ha) as compare to other treatments and statistically at par with (NDR- 359 + Poultry manure 2.5t/ha + Vermicompost 5t/ha). Whereas minimum grain yield is seen in Treatment T7 (MTU-7029 + Poultry manure 5t/ha).

The highest straw yield was recorded in Treatment T9 (MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha) (6.43 t/ha) as compare to other treatments and statistically at par with (NDR- 359 + Poultry manure 5t/ha), (NDR- 359 + Poultry manure 2.5t/ha + Vermicompost 5t/ha), (BPT-5204 + Poultry manure 5t/ha) and (BPT-5204 + Poultry manure 2.5t/ha + Vermicompost 10t/ha). Whereas minimum straw yield is seen in Treatment T5 (BPT-5204 + Vermicompost 10t/ha).

The highest harvest index was recorded in Treatment T9 (MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha) (44.7%) as compare to other treatments and statistically at par with (NDR- 359 + Poultry manure 2.5t/ha + Vermicompost 5t/ha). Whereas minimum harvest index is seen in Treatment T5 (BPT-5204 + Vermicompost 10t/ha).

The supremacy of enriched poultry manure compost lies in the fact that it can supply the nutrients in soluble form for a quite longer period by not allowing the entire soluble form into solution to come into contact with soil and other inorganic constituents, thereby minimising fixation and precipitation from the enriched manures. The plant roots can very well compete with loss mechanisms and absorb more nutrients, leading to a better yield.

The maximum grain and straw yield were due to marked improvement in dry matter accumulation, yield attributes and greater nutrient content and their uptake by rice crop. These findings are in direct conformity with that of **Barik et al. (2008)**.

Yield validation using SPSS model

The multi-regression analysis using SPSS has been employed for the estimation of rice yield. The regression for SPSS model is

$$Y = 13.661 + (0.263 \times Z21 \text{ of prediction year}) + (0.066 \times \text{time})$$

Here, Z21 is the sum product of minimum temperature.

The yield obtained in treatment 9 with MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha (5.21t/ha) showed 44.66% increase over the predicted yield through SPSS model (2.88 t/ha).




Fig 1. Percentage of yield increase over SPSS model

Table 1. Effect of organic manures on growth attributes of rice varieties

Sl. no	Treatments	100 DAT			80-100 DAT	
		Plant height (cm)	Number of tillers/hill	Dry weight (g/hill)	Crop growth rate (g/m ² /day)	Relative growth rate (g/g/day)
1.	NDR- 359 + Poultry manure 5t/ha	108.2	9.0	52.95	17.43	0.0110
2.	NDR- 359 + Vermicompost 10t/ha	106.3	9.3	51.88	18.23	0.0120
3.	NDR- 359 + Poultry manure 2.5t/ha + Vermicompost 5t/ha	108.3	9.9	53.32	17.76	0.0110
4.	BPT-5204 + Poultry manure5t/ha	106.7	9.5	52.13	18.13	0.0120
5.	BPT-5204 + Vermicompost 10t/ha	105.8	9.7	50.09	16.97	0.0110
6.	BPT-5204 + Poultry manure 2.5t/ha + Vermicompost 5t/ha	107.1	10.4	52.34	17.97	0.0113
7.	MTU-7029 + Poultry manure 5t/ha	106.5	11.8	52.64	17.90	0.0110
8.	MTU-7029 + Vermicompost 10t/ha	105.4	12.3	51.47	18.63	0.0123
9.	MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha	106.9	13.1	52.79	17.88	0.0110
	F test	S	S	S	S	S
	SEm(+)	0.20	0.36	0.09	0.21	0.0002
	CD(P=0.05)	0.60	1.09	0.28	0.64	0.0005

Table 2. Effect of organic manures on yield attributes of rice varieties

Sl. no	Treatments	No. of effective tiller/hill	No. of panicles/hill	No. of grains/panicle	Test weight (g)	Grain yield (t/ha)	Stover yield (t/ha)	Harvest index (%)
1.	NDR- 359 + Poultry manure 5t/ha	8.33	6.40	100.47	21.29	4.58	6.20	42.49
2.	NDR- 359 + Vermicompost 10t/ha	8.60	6.27	103.00	20.69	4.51	6.14	42.34
3.	NDR- 359 + Poultry manure 2.5t/ha + Vermicompost 5t/ha	9.00	6.60	101.67	21.20	4.88	6.30	43.67
4.	BPT-5204 + Poultry manure5t/ha	9.13	8.70	101.93	14.72	4.45	6.30	41.39
5.	BPT-5204 + Vermicompost 10t/ha	9.13	8.73	103.47	14.60	4.49	6.17	42.10
6.	BPT-5204 + Poultry manure 2.5t/ha + Vermicompost 5t/ha	10.00	9.21	101.13	15.14	4.76	6.40	42.67
7.	MTU-7029 + Poultry manure 5t/ha	9.93	5.40	106.73	21.02	4.08	5.82	41.24
8.	MTU-7029 + Vermicompost 10t/ha	10.73	5.60	106.23	21.17	4.28	5.84	42.29
9.	MTU-7029 + Poultry manure 2.5t/ha + Vermicompost 5t/ha	11.33	6.14	112.37	22.35	5.21	6.43	44.74
	F test	S	S	S	S	S	S	S
	SEm(+)	0.31	0.14	1.07	0.18	0.09	0.09	0.50
	CD(P=0.05)	0.93	0.41	3.21	0.55	0.28	0.26	1.49

Conclusion

Based on the findings, it can be said that the inclusion of poultry manure at a rate of 2.5 t/ha alongside vermicompost at a rate of 5 t/ha improved the output and economic return among the three types of rice in MTU-7029. Additional studies are required to corroborate the outcomes because they are based on only one season.

References

- Altaf Hussain, S., Sheraz Mahdi, R.A., Bhat Faisul-ur-Rasool and Raihana Habib Kanth. Integrated nutrient management of rice (*Oryza sativa* L.) under temperate conditions of Kashmir. *Agricultural Science Digest*. 2012; 32(1): 18-22.
- Amanullah, M.M., Alagesan, A.,Vaiyapuri, K., Pazhanivelan, Sand Sathyamoorthi, K. Intercropping and organic manures on the growth and yield of cassava(*Manihotesculenta* crantz.). *Research Journal of Agriculture and Biological Sciences*. 2006:2: 183-189.
- Appa Rao GJ. Estimation of wheat yields over Punjab using district and state models. *Mausam*. 1983;34(3):275-80.
- Barik, A. K.; Raj, A. and Saha. R. K. Yield performance, economics and soil fertility through organic sources (vermicompost) of nitrogen as substitute to chemical fertilizers in wet season rice. *Crop Research* (Hisar). 2008; 36: 1/3, 4-7.
- Das S, Das A, Idapuganti R, Layek J, and Chowdary S. Growth and physiology of groundnut as influenced by micronutrients and liming in acid soil of North East India. *Indian Journal of Hill farming*. 2016;29:40-47.
- Dass, A. and Chandra. Irrigation, spacing and cultivar effects on net photosynthetic rate, dry matter partitioning and productivity of rice under system of rice intensification in Mollisol of northern India. *Experimental Agriculture*. 2013; 49(4), 504-523.

- Datta, M and Banik, S. Effect of poultry manure and phosphate dissolving bacteria on rice (*Oryza sativa*) in arid soil. *Indian Journal of Agricultural Sciences*. 1994;64(11): 791-793.
- Edward, D.Rand Daniel, T.C. Environmantal impacts of on-farm poutry waste disposal - A review. *Bioresource Technology*. 1992: 41(1): 9-33.
- Government of India(GOI). Agricultural statistics at a glance: Ministry of Agriculture& Farmers Welfare, Government of India. 2016.
- Helen Belefant-Miller. Poultry litter induces tillering in rice. *Journal of Sustainable Agriculture*. 2007;3(1):151-159.
- Hossan, A.T.M.S., Rahman, F., Saha, P.Kand Solaiman, A.R.M. Effects of different aged poultry litter on the yield and nutrient balance in boro rice cultivation. *Bangladesh Journal of AgriculturalResearch*. 2010;35(3): 497-505.
- Jain RC, Sridharan H, Agarwal R. Principal component technique for forecasting of sorghum yield. *Indian Jornal of Agricultural Sciences*. 1984;54(6):467-70.
- M. Mohamed Amanullah, S. Sekar and P. Muthukrishnan, 2010. Prospects and Potential of Poultry Manure. *Asian Journal of Plant Sciences*, 9(4): 172-182.
- Prabhakaran, C. Studies on organic farming in tomato. M.Sc., (Ag) Thesis, Tamil Nadu Agricultural University, Coimbatore. 2000.
- Ramasubramanian V, Jain RC. Use of growth indices in Markov Chain model for crop yield forecasting. *Biometrical Journal* 1999;41(1):99-109.
- Santosh Kumar Jha, Tripathi, R.S and Sanjeev Malaiya. Influence of integrated nutrient management practices on growth and yield of scented rice (*Oryza sativa* L.). *Annals of Agricultural Research New Series*. 2004;25(1): 159-161.
- Shinde, P.H., Naik, R.L., Nazirkar, R.B., Kadam, S.K and Khaire, V.N. Evaluation of vermicompost. Proceedings of the National Seminar on Organic Farming, 18 –19 April, 1992.
- Singh, G., Wade, L. J., Singh, R. K. Nayak, R. Singh, B. B. and Singh, O. N. Nutrient management for rainfed lowland rice and its effect on succeeding lentil crop. *Oryza*, 2001, 38(3) 123-126.

- Singh, G.R., Parihar and Chaure, N.K. Effect of poultry manure and its combination with nitrogen in rice-wheat cropping system. *Indian Agriculturist*. 2001;45(3&4):235-240.
- Sridhar, K., Srinivas, A., Kumar, A.K., Ramprakesh, T., & Rao, R.P. Physiological growth parameters of rabi rice (*Oryza sativa* L.) under alternate wetting and drying irrigation with varied nitrogen levels. *International Journal Current Microbiology and Applied Sciences*. 2019;8, 01-15.
- Sujit Adhikary. Vermicompost, the story of organic gold: A review. *Agricultural Sciences*. 2012;03-07: 905-917.
- Suvarna Latha, A.J and Sankara Rao, V. Integrated use of fertilizers and poultry manure on nutrient availability and yield of rice. *Journal of the Indian Society of Coastal Agricultural Research*. 2001;19 (2):153-157.
- Tayefe, M., Akif, G., Amiri, E., & Azin, N.Z. Effect of nitrogen on rice yield, yield components and quality parameters. *African Journal of Geology, Ecology, and Landscapes*. 2014; **13**(1), 91-105.
- Walker G.K.. Model for operational forecasting of western Canada wheat yield. *Agricultural and Forest Meteorology*. 1989;44(3-4):339-351.