

Evaluation of Haemostatic effects of *Petroselinum crispum* (Parsley) in Wistar rats

Abstract

Petroselinum crispum (parsley or garden parsley), besides its use as an edible vegetable, is used in the treatment of various ailments with documented evidence validating some of these medicinal uses. The present study evaluates the impact of oral administration of *P. crispum* on blood coagulation parameters. Fresh leaves of *P. crispum* were locally sourced, authenticated and extracted using ethanol to obtain an ethanolic extract of *P. crispum* (PCE). Twenty-five (25) male Wistar rats were used for the study and were randomly assigned to five (5) animals of five (5) rats each. Group I served as the control group and was given distilled water, whereas groups II, III, and IV were treated as experimental groups and administered PCE at 200, 400, and 800mg/kg via oral gavage, respectively. Group V also received 800mg/kg of PCE with the administration terminated on the 28th day to allow for a 14-day recovery period. Platelet count (PLT) and other platelet indices were determined using an automated Haematology analyzer. Bleeding time (BT), clotting time (CT), prothrombin time (PT), activated partial thromboplastin time (aPTT) and plasma fibrinogen (FIB) were assayed following standard laboratory protocols. The results of the study show that the mean PLT were significantly reduced in the experimental animals compared to the control ($p < 0.05$) while BT, CT, PT, aPTT and FIB significantly increased compared to the control ($p < 0.05$). Also, BT, CT, PT, and aPTT recovered towards the control values after the 14-day withdrawal period. The present study shows ethanolic leaf extract of *P. crispum* prolonged blood coagulation. The current evidence suggests that *P. crispum* could be a candidate for further exploration as a natural plant-based antithrombotic agent.

Keywords: *Petroselinum crispum*, haemostasis, bleeding time, clotting time, prothrombin time, activated partial prothrombin time.

INTRODUCTION

Petroselinum crispum, commonly referred to as parsley or garden parsley is a fragrant and flavorful herb that belongs to the Apiaceae family [1, 2]. It is native to the Central and Eastern Mediterranean region but naturalized and used in Europe, Middle Eastern and American cuisine. It has been employed as food, perfume and cosmetics and in the treatment of various ailments [3, 4]. It is grown worldwide for its edible leaves, which can be used fresh or dried as a seasoning. It is also used as a garnish for salads, snacks, vegetable soups, stews and casseroles. The ancient Greeks and Romans used it in funeral decoration and as deodorants [1, 5]. *P. crispum* is used in traditional medicine in different parts of the world in the management of various illnesses, including gastrointestinal disorders, amenorrhea, dysmenorrhea, diabetes mellitus, skin diseases, fever, ulcers, rheumatism, and hypertension. It has been found useful as an anti-inflammatory, anti-urolithiasis, carminative, gastrotonic, antiseptic and diuretic [1, 4, 6, 7].

Several bioactive compounds and phytochemical constituents have been extracted from various parts of *P. crispum*, including its seeds, roots, leaves, and petioles. These compounds are predominantly composed of carbohydrates, flavonoids such as apigenin, quercetin, isorhamnetin, luteolin, chrysoeriol and eugenol. Also, coumarins and essential oils such as myristicin and apiol, and other compounds [1, 8, 9]. Furthermore, the plant is a good source of iron, phosphorous, calcium, zinc and vitamins A & C [10-12]. Screened pharmacological actions of *P. crispum* include antimicrobial [13, 14], anti-inflammatory [15, 16], antioxidant [9, 17], anti-diabetic [18, 19], immunomodulatory [20, 21], genitourinary [22, 23], neuroprotective [24], analgesic [25, 26], spasmolytic [27, 28], anti-cancer [29, 30], anti-hypertensive [4], and anti-platelet [31] activities. Despite the array of documented pharmaceutical benefits, *P. crispum* and its essential oils have to been demonstrated to be hepatotoxic and nephrotoxic at dosages equal to or higher than 1000mg/kg [32].

Haemostasis refers to the process of stopping or preventing bleeding in response to damaged blood vessels. This involves a series of physiological events that result in the production of a localized clot at the site of injury in a very short amount of time, usually seconds to minutes [33, 34]. When a blood vessel is damaged, the body naturally produces a reflex vasoconstriction to slow the loss of blood and allow more time for the formation of a platelet plug and the initiation of blood coagulation. The process of haemostasis involves the formation of a platelet plug, followed by the activation of coagulation factors to form a fibrin mesh, fibrinolysis, and eventually the repair of the blood vessel [33, 35]. The haemostatic cascade is made up of the intrinsic and extrinsic pathways, which originate differently but converge at the point of fibrin activation to stabilize the platelet plug. The intrinsic pathway is activated by trauma within the vascular system, while the extrinsic pathway is activated by trauma outside the vascular system [36, 37]. This process largely relies on the platelet count and function, as well as other coagulation parameters to achieve a steady balance between the fluid and solid states of blood and prevent pathological thrombosis. Thrombosis is the formation of a blood clot within a blood vessel, which can partially or completely obstruct blood flow leading to cardiovascular complications and death [38, 39]. Due to the negative side effects, high cost, and limited availability of synthetic medications, the utilization of plant-derived natural substances derived from plants has increasingly gained popularity and has become the primary approach in human pharmacotherapy [40-42]. This has led to the screening of possible natural anti-thrombotic agents derived from natural products. Although there has been increasing interest in exploring the pharmacological properties of *P. crispum*, the impact it has on blood coagulation and its potential as an anti-thrombotic agent remain unclear or have yielded conflicting findings. This present study, therefore, aims to evaluate the haemostatic effect of *P. Crispum* using Wistar rat models.

MATERIALS AND METHODS

Source and Preparation of Plant Extract

Fresh leaves of *P. crispum* were procured from Fruit Garden Market, Port Harcourt, Nigeria and authenticated at the Plant Science and Biotechnology Department of the University of Port Harcourt. The leaves were carefully washed to eliminate any sand or dirt and allowed to air dry at room temperature for four (4) weeks. Following air-drying, the leaves were subjected to Soxhlet extraction using 99.5% ethanol [43]. The resulting extract of *P. crispum* (PCE) was stored at a temperature of 5°C until administration.

Experimental Animals

For the study, a total of 25 male Wistar rats weighing between 150 to 200g were obtained from the animal house of the Faculty of Basic Medical Sciences, University of Port Harcourt. These rats were given two weeks to acclimatize during which they had unlimited access to standard rat pellets and water. The rats were kept in standard cages with appropriate ventilation, temperature, and humidity, and were subjected to a regular 12-hour light/dark cycle. Throughout the study, the animals were treated in accordance with the Animal Use and Care Guidelines of the National Veterinary Research Council of Nigeria.

Experimental Design

A total of 25 male Wistar rats were divided into five groups of five animals each, through a random allocation process. Group I served as the control group and was given distilled water, whereas groups II, III, and IV were treated as experimental groups and administered with *P. crispum* ethanolic extract at 200, 400, and 800mg/kg via oral gavage, respectively. Group V also received 800mg/kg of PCE. However, the administration of PCE was stopped for Group V after 28 days and they were given access to standard rat pellets and water for 14 days to recover.

Laboratory Analysis

Determination of Bleeding Time

Bleeding time (BT) was determined using a modified Duke's method [44, 45]. A surgical blade was utilized to create a small incision at the end of each rat's tail. This incision was sufficient to cause bleeding, and a stopwatch was started as soon as the first drop of blood was visible. Every 15 seconds, the bleeding area was gently cleaned using a dry cotton wool swab, and the stopwatch was stopped once the bleeding ceased completely. The time taken for the bleeding to stop was recorded for each rat as the bleeding time.

Determination of Clotting Time

Clotting time (CT) was assayed using a modified Ivy's method [45, 46]. The tail tip of each rat was cut using a dissection blade to initiate bleeding. A single drop of blood was collected from the incision on a clean and dry microscope slide, with the stopwatch started simultaneously as the blood came in contact with the slide. At 15-second intervals, the tip of a clean and dry pin was cautiously passed through the blood droplet on the slide, and the stopwatch was stopped immediately upon observation of a fibrin thread, this was recorded as the clotting time.

Determination of Platelet Count and Clotting Parameters

First, the animals were humanely anaesthetized by cervical dislocation and blood samples were taken directly from animals by cardiac puncture. These samples were collected in two types of **sample** bottles: EDTA **sample** bottles for platelet analysis and sodium citrate **sample** bottles for coagulation tests. The platelet count (PLT), plateletcrit (PCT), mean platelet volume (MPV), platelet distribution width (PDW) and platelet large ratio (PL-R) were measured using an automatic Haematology analyzer (Mindray, China) while the prothrombin time (PT), activated partial thromboplastin time (aPTT) and fibrinogen (FIB) were analyzed using the protocol provided by standard assay kit (Helena Laboratories, USA).

The animals received care and handling in compliance with the Animal Use and Care Guidelines set forth by the National Veterinary Research Council of Nigeria and the guide for the care and use of laboratory animals [47].

Statistical Analysis

Data were analyzed using IBM Statistical Product and Service Solutions (SPSS version 25). For each research group, the mean and standard error of the mean for each parameter was calculated. The mean values of the experimental groups (II, III, IV, and V) were compared to the control group (Group I) using analysis of variance (ANOVA), followed by a least significant difference (LSD) posthoc analysis. A p-value less than 0.05 was considered statistically significant.

RESULTS

Table 1: The effect of *P. crispum* on platelet count and platelet indices of male Wistar rats

Parameters	Control (n=5)	200mg/kg PCE (n=5)	400mg/kg PCE (n=5)	800mg/kg PCE (n=5)	800mg/kg PCE + Recovery (n=5)
PLT (x10 ⁹ /L)	456.00±46.05	492.80±79.81	230.40±52.29*	361.40±70.69	487.60±93.07
PCT (%)	0.33±0.4	0.37±0.07	0.14±0.06	0.26±0.05	0.38±0.07
MPV (fL)	7.24±0.11	7.46±0.11	7.24±0.16	7.42±0.22	7.68±0.13
PDW (%)	10.00±0.13	10.32±0.16	9.98±0.22	10.3±0.36	10.60±0.12
PL-R	12.30±0. ±97	12.32±0.84	10.30±1.32	12.16±1.73	13.84±0.61

Result is given as mean±standard error of mean; *significantly different compared to control (p<0.05)

Table 1 displays the impact of oral administration of ethanolic leaf extract of *Petroselinum crispum* on male Wistar rats' platelet count and other platelet indices for 28 days, followed by a 14-day stoppage of administration for animals in Group V. The findings indicate a significant decrease in platelet count among animals in Group III (400mg/kg PCE) compared to the control (P<0.05). Additionally, during the 14-day recovery period, the platelet counts increased, with no significant difference observed between Group V (800mg/kg PCE + recovery) animals and the control (p>0.05).

Table 2: The effect of *P. crispum* on some coagulation parameters of male Wistar rats

Parameters	Control (n=5)	200mg/kg PCE (n=5)	400mg/kg PCE (n=5)	800mg/kg PCE (n=5)	800mg/kg PCE + Recovery (n=5)
------------	------------------	-----------------------	-----------------------	-----------------------	----------------------------------

BT (secs)	73.20±6.47	93.80±9.52	95.80±5.52	109.80±6.36*	72.160±10.83
CT (secs)	45.00±3.2	62.00±7.89	68.00±8.74	79.40±14.97*	40.40±5.33
PT (secs)	18.80±2.08	24.40±1.80	26.60±2.37	31.80±7.33*	17.00±1.14
aPTT (secs)	16.20±0.60	18.40±3.12	21.20±1.20	22.60±1.96*	19.40±1.81
FIB (mg/dL)	174.40±8.63	246.40±7.13*	240.80±9.28*	252.00±7.67*	257.2±8.16

Result is given as mean±standard error of mean; *significantly different compared to control (p<0.05)

Table 2 shows the effect of oral administration of ethanolic leaf extract from *Petroselinum crispum* on the bleeding time, clotting time, prothrombin time, activated partial thromboplastin time and plasma fibrinogen on male Wistar rats after 28-day oral administration followed by 14 days recovery period for animals in Group V. The results show that the BT, CT, PT and aPTT was significantly raised for rats in Group IV (800mg/kg PCE) compared to the control (P<0.05). Similarly, the mean FIB level increased for all the experimental groups (II, III, IV & V). Following a 14-day recovery, mean BT, CT, PT and aPPT reduced towards the control values while FIB continued to increase when compared with the control (P<0.05).

DISCUSSION

Due to the high cost of antithrombotic treatment, there is now a focus on exploring various natural and plant-based alternatives as potential antithrombotic agents. With over 400 promising natural substances in different phases of clinical evaluation as possible antithrombotic agents [2, 48], the present study evaluated the effect of ethanolic extract of *P. crispum* on blood coagulation and its potential as an anti-thrombotic agent using Wistar rat models.

Effect on Platelet Count and Other Platelet Indices

There was a significantly reduced platelet count for animals supplemented with 400mg/kg (Group III) of PCE compared to the control (p<0.05) (Table 1). Discontinuation of the administration of PCE appeared to improve the platelet count raising the mean values to 487.60±93.07 (x10⁹/L) for group IV compared to the control value of 456.00±46.05 (x10⁹/L) after 14 days of recovery. No significant difference was observed in mean PCT, MPV, PDW and PL-R between the experimental groups and control. Platelets are the smallest blood cells with primary function in the process of blood coagulation, the body's natural response to stop bleeding. When there is an injury to a blood vessel, platelets become activated and clump together to form a plug, which stops the bleeding. They also release chemicals that attract other cells to the site of injury and promote healing [49, 50]. Platelets can contribute to the development of diseases either directly, as a primary disorder of their function or quantity, or indirectly, through their critical role in blood clotting, which can lead to conditions like coronary artery disease, stroke, peripheral vascular disease, and diabetes [51, 52]. The significantly reduced platelet count observed among the animals that received 400mg/kg PCE could be attributed to the reported anti-platelet aggregation activities of *P. crispum* [2, 31], hence PCE may have interfered with clot formation by reducing platelet counts. Furthermore, platelet counts have been reported to increase with inflammation [52, 53] while *P. crispum* has been shown to possess anti-inflammatory activity [15, 16], hence it is possible that *P. crispum* attenuates

inflammation by reducing platelet count. The improved platelet count among the Group V animal appears to be due to the cessation of the effect of PCE to interfere with platelet production or destruction. Shcherazade *et al.*, observed no significant difference in platelet count among phenylhydrazine-induced anaemic mice treated with *P. crispum* [54] while Al-Daraji *et al.*, reported that dietary supplementation of local geese with *P. crispum* as a food additive resulted in increased platelet count [55].

Effects of Coagulation Parameters

The result of the present study shows significantly raised BT, CT, PT and aPTT for the highest dose of PCE (800mg/kg) compared to the control ($p < 0.05$) (Table 2). Coagulation parameters evaluate the clotting ability of the blood while measuring various components of the coagulation process. These prolonged coagulation parameters may be due to the potent anti-platelet aggregation activities of *P. crispum* [2, 31]. The activation and aggregation of blood platelets serve a significant physiological purpose in the process of haemostasis. There is substantial evidence suggesting that numerous cardiovascular disorders are connected to abnormal and excessive activation of platelets [31, 53]. Platelet activation and aggregation may have been disrupted by PCE, which could have led to the prolongation of BT, CT, PT, and aPTT. The study also observed that following a 14-day recovery period, BT, CT, PT and aPTT values returned towards normal compared to the control. This suggests that the reduced coagulation parameters could be due to the termination of the effect of PCE to inhibit platelet aggregation. Plasma fibrinogen was significantly increased among the experimental groups compared to the control with the increase continuing after the caseation of the PCE supplementation (Table 2). Plasma fibrinogen is a vital component of the coagulation cascade and serves as a major factor in the determination of blood flow and viscosity [56]. It basically serves to create fibrin, which aids in the formation of a hemostatic plug by binding platelets and plasma proteins which ensures the caseation of bleeding [57]. Although increased plasma fibrinogen concentration is associated with enhanced blood clotting (procoagulant), plasma fibrinogen at certain concentrations has been associated with anti-coagulant activities [58]. While the elevated plasma fibrinogen following PCE administration may not be clearly understood, it is possible that the elevated FIB, in this case, acted as an anticoagulant leading to the observed prolonged BT, CT, PT and aPTT. Chao *et al.* observed that lower plasma concentrations enhanced platelet aggregation while higher concentrations inhibited platelet aggregation [59]. Similarly, it has been observed that following low tissue factor stimulus, elevated fibrinogen served as an anticoagulant by prolonging clotting time and decreasing clot stability. This goes to show that increased plasma fibrinogen may not be thrombogenic as previously conceived [58]. It is also possible that PCE may have in some way affected the production of fibrinogen in the liver.

CONCLUSION

The result of the present study shows that ethanolic extract of *Petroselinum crispum* reduced platelet count, increased, bleeding clotting, prothrombin and activated partial thromboplastin time as well as plasma fibrinogen concentration. The current evidence suggests that *P. crispum* could be a candidate for further exploration as a natural plant-based antithrombotic agent.

Ethical Approval

The research protocol and design were approved by the University of Port Harcourt Research Ethics Committee (UPH/CEREMAD/REC/MM87/030).

REFERENCES

1. Agyare C, Appiah T, Boakye YD, Apenteng JA. *Petroselinum crispum*: a review. *Medicinal spices and vegetables from Africa*. 2017;527-47.
2. Chaves DS, Frattani FS, Assafim M, de Almeida AP, Zingali RB, Costa SS. Phenolic chemical composition of *Petroselinum crispum* extract and its effect on haemostasis. *Natural Product Communications*. 2011;6(7):1934578X1100600709.
3. Lopez M, Sanchez-Mendoza I, Ochoa-Alejo N. Comparative study of volatile components and fatty acids of plants and in vitro cultures of parsley (*Petroselinum crispum* (Mill) nym ex hill). *Journal of agricultural and food chemistry*. 1999;47(8):3292-6.
4. Ajebli M, Eddouks M. Antihypertensive activity of *Petroselinum crispum* through inhibition of vascular calcium channels in rats. *Journal of ethnopharmacology*. 2019;242:112039.
5. El-Ahmady S, Ibrahim N, Farag N, Gabr S. *Apiaceae Plants Growing in the East: Centuries of Healing Traditions and Culture*. *Ethnopharmacology of Wild Plants*: CRC Press; 2021:246-300.
6. Eddouks M, Ajebli M, Hebi M. Ethnopharmacological survey of medicinal plants used in Daraa-Tafilalet region (Province of Errachidia), Morocco. *Journal of ethnopharmacology*. 2017;198:516-30.
7. Farzaei MH, Abbasabadi Z, Ardekani MRS, Rahimi R, Farzaei F. Parsley: a review of ethnopharmacology, phytochemistry and biological activities. *Journal of traditional Chinese medicine*. 2013;33(6):815-26.
8. Pápay ZE, Kósa A, Boldizsár I, Ruzskai Á, Balogh E, Klebovich I, et al. Pharmaceutical and formulation aspects of *Petroselinum crispum* extract. *Acta Pharmaceutica Hungarica*. 2012;82(1):3-14.
9. Zhang H, Chen F, Wang X, Yao H-Y. Evaluation of antioxidant activity of parsley (*Petroselinum crispum*) essential oil and identification of its antioxidant constituents. *Food research international*. 2006;39(8):833-9.
10. Popović M, Kaurinović B, Jakovljević V, Mimica-Dukic N, Bursać M. Effect of parsley (*Petroselinum crispum* (Mill.) Nym. ex AW Hill, Apiaceae) extracts on some biochemical parameters of oxidative stress in mice treated with CCl₄. *Phytotherapy research*. 2007;21(8):717-23.
11. Caunii A, Cuciureanu R, Zakar AM, Tonea E, Giuchici C. Chemical composition of common leafy vegetables. *Studia Universitatis "Vasile Goldis" Arad. Seria Stiintele Vietii (Life Sciences Series)*. 2010;20(2):45.
12. Hossein M. Determination of mineral nutrients in some leafy vegetables. *Acta horticulturae*. 2010;877:1229-32.
13. Linde G, Gazim Z, Cardoso B, Jorge L, Tešević V, Glamočlija J, et al. Antifungal and antibacterial activities of *Petroselinum crispum* essential oil. *Genetics and Molecular Research*. 2016;15(3).
14. Petrolini FVB, Lucarini R, Souza MGMd, Pires RH, Cunha WR, Martins CHG. Evaluation of the antibacterial potential of *Petroselinum crispum* and *Rosmarinus officinalis* against

- bacteria that cause urinary tract infections. *Brazilian Journal of Microbiology*. 2013;44:829-34.
15. Slighoua M, Mahdi I, Di Cristo F, Amaghnoije A, Grafov A, Boucetta N, et al. Assessment of in vivo estrogenic and anti-inflammatory activities of the hydro-ethanolic extract and polyphenolic fraction of parsley (*Petroselinum sativum* Hoffm.). *Journal of Ethnopharmacology*. 2021;265:113290.
 16. Al-Howiriny TA, Al-Sohaibani MO, El-Tahir KH, Rafatullah S. Preliminary evaluation of the anti-inflammatory and anti-hepatotoxic activities of parsley (*Petroselinum crispum*) in rats. *Journal of natural remedies*. 2003;54-62.
 17. Fejes S, Kery A, Blazovics A, Lugasi A, Lemberkovics E, Petri G, et al. Investigation of the in vitro antioxidant effect of *Petroselinum crispum* (Mill.) Nym. ex AW Hill. *Acta Pharmaceutica Hungarica*. 1998;68(3):150-6.
 18. Yanardağ R, Bolkent Ş, Tabakoğlu-Oğuz A, Özsoy-Saçan Ö. Effects of *Petroselinum crispum* extract on pancreatic B cells and blood glucose of streptozotocin-induced diabetic rats. *Biological and Pharmaceutical Bulletin*. 2003;26(8):1206-10.
 19. Soliman HA, Eltablawy NA, Hamed MS. The ameliorative effect of *Petroselinum crispum* (parsley) on some diabetes complications. *Journal of Medicinal Plants Studies*. 2015;3(4):92-100.
 20. Yousofi A, Daneshmandi S, Soleimani N, Bagheri K, Karimi MH. Immunomodulatory effect of Parsley (*Petroselinum crispum*) essential oil on immune cells: mitogen-activated splenocytes and peritoneal macrophages. *Immunopharmacology and Immunotoxicology*. 2012;34(2):303-8.
 21. Karimi MH, Ebadi P, Amirghofran Z. Parsley and immunomodulation. *Expert Review of Clinical Immunology*. 2012;8(4):295-7.
 22. Yoshikawa M, Uemura T, Shimoda H, Kishi A, Kawahara Y, Matsuda H. Medicinal foodstuffs. XVIII. Phytoestrogens from the aerial part of *Petroselinum crispum* Mill. (Parsley) and structures of 6"-acetylapiin and a new monoterpene glycoside, petroside. *Chemical and pharmaceutical bulletin*. 2000;48(7):1039-44.
 23. Abdel-Wahhab MA, Abbas S, Salah-Abbas JB, Hassan AM, Oueslati R. Parsley oil protects against Zearalenone-induced alteration in reproductive function in male mice. *Toxicology Letters*. 2006(164):S266.
 24. Allam AA, Maodaa SN, Abo-Eleneen R, Ajarem J. Protective effect of parsley juice (*Petroselinum crispum*, Apiaceae) against cadmium deleterious changes in the developed albino mice newborns (*Mus musculus*) brain. *Oxidative medicine and cellular longevity*. 2016;2016.
 25. Eidi A, Eidi M, Badieli L. Antinociceptive effects of ethanolic extract of parsley (*Petroselinum crispum* L.) leaves in mice. *Medical Science Journal of Islamic Azad University-Tehran Medical Branch*. 2009;19(3):181-6.
 26. Al-khazraji SM. Studying the Analgesic, Anti-inflammatory and Antipyretic Properties of The Aqueous Extract of *Petroselinum crispum* in Experimental Animal Models. *IOSR Journal Of Pharmacy*. 2015;5(9):17-23.
 27. Branković S, Kitić D, Radenković M, Ivetić V, Veljković S, Nešić M. Relaxant activity of aqueous and ethanol extracts of parsley (*Petroselinum crispum* (Mill.) Nym. ex AW Hill, Apiaceae) on isolated ileum of rat. *Medicinski preglod*. 2010;63(7-8):475-8.

28. Damabi NM, Moazedi AA, Seyyednejad SM. The role of α - and β -adrenergic receptors in the spasmolytic effects on rat ileum of *Petroselinum crispum* Latifolium (parsley). *Asian Pacific Journal of Tropical Medicine*. 2010;3(11):866-70.
29. Aissani N, Albouchi F, Sebai H. Anticancer effect in human glioblastoma and antioxidant activity of *Petroselinum crispum* L. methanol extract. *Nutrition and Cancer*. 2021;73(11-12):2605-13.
30. Griffiths IB, Douglas RGA. Phytophotodermatitis in pigs exposed to parsley (*Petroselinum crispum*). *Veterinary Record*. 2000;146(3):73-4.
31. Gadi D, Bnouham M, Aziz M, Ziyyat A, Legssyer A, Bruel A, et al. Flavonoids purified from parsley inhibit human blood platelet aggregation and adhesion to collagen under flow. *Journal of Complementary and Integrative Medicine*. 2012;9(1).
32. Awe EO, Banjoko SO. Biochemical and haematological assessment of toxic effects of the leaf ethanol extract of *Petroselinum crispum* (Mill) Nyman ex AW Hill (Parsley) in rats. *BMC Complementary and Alternative Medicine*. 2013;13(1):1-6.
33. Zaidi A, Green L. Physiology of haemostasis. *Anaesthesia & Intensive Care Medicine*. 2019;20(3):152-8.
34. Sira J, Eyre L. Physiology of haemostasis. *Anaesthesia & Intensive Care Medicine*. 2016;17(2):79-82.
35. Chinko BC, Amah-Tariah FS, Ekenna IC. Evaluation of the effects of calabash chalk on the haematological profile of Wistar rats. *Notulae Scientia Biologicae*. 2022;14(3):11281-.
36. Chaudhry R, Babiker HM. Physiology, coagulation pathways. StatPearls: StatPearls Publishing; 2018.
37. Fareed J, Iqbal O. Physiology of Hemostasis. *Anticoagulation and Hemostasis in Neurosurgery*: Springer; 2016:3-16.
38. Zaman AG, Helft G, Worthley SG, Badimon JJ. The role of plaque rupture and thrombosis in coronary artery disease. *Atherosclerosis*. 2000;149(2):251-66.
39. Lippi G, Franchini M, Targher G. Arterial thrombus formation in cardiovascular disease. *Nature Reviews Cardiology*. 2011;8(9):502-12.
40. Atanasov AG, Waltenberger B, Pferschy-Wenzig E-M, Linder T, Wawrosch C, Uhrin P, et al. Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnology advances*. 2015;33(8):1582-614.
41. Ugoeze KC, Oluigbo KE, Chinko BC. Phytochemical and Nutritional Benefits of the GC-FID Quantified Phytochemicals of the Aqueous Extract of *Azadirachta indica* leaves. *Journal of Pharmacy and Pharmacology Research*. 2020;4(4):149-63.
42. Hart VO, Chinko BC. Effect of *Terminalia catappa* Seed Extract on Liver Enzymes and Lipid Profile in Male Wistar Rat. *Asian Journal of Research in Botany*. 2022;8(4):25-32.
43. Evans WC. *Trease and Evans' Pharmacognosy*. 16th ed. New York: Elsevier Health Sciences; 2009.
44. Russeau AP, Vall H, Manna B. Bleeding time. StatPearls [Internet]: StatPearls Publishing; 2021.
45. Bain BJ, Bates I, Laffan MA. *Dacie and lewis practical haematology e-book*: Elsevier Health Sciences; 2016.
46. Mokwenye I, Ikewuchi CC, Okoye NF. Evaluation of the haemostatic potentials of aqueous extract of *Hyptis suaveolens* leaves in wistar rats. *Bio-Research*. 2023;21(1):1818-27.
47. Albus U. Guide for the care and use of laboratory animals (8th edn). SAGE Publications Sage UK: London, England; 2012.

48. Hilbrich L, Truelsen T, Yusuf S. Stroke and cardiovascular diseases: the need for a global approach for prevention and drug development. *International Journal of Stroke*. 2007;2(2):104-8.
49. Jurk K, Kehrel BE. Platelets: physiology and biochemistry. *Seminars in thrombosis and hemostasis*, 2005. Copyright© 2005 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New ...: 381-92.
50. Gremmel T, Frelinger AL, Michelson AD. Platelet Physiology. *Semin Thromb Hemost*. 2016;42(03):191-204.
51. Andrews RK, Berndt MC. Platelet physiology and thrombosis. *Thrombosis Research*. 2004;114(5):447-53.
52. Andrews RK, Berndt MC. Platelet physiology and thrombosis. *Thrombosis research*. 2004;114(5-6):447-53.
53. Gawaz M, Langer H, May AE. Platelets in inflammation and atherogenesis. *The Journal of clinical investigation*. 2005;115(12):3378-84.
54. Shcherazade O-SF, Pétronille A-Z, Zana C, John AGKK, Bertin YK. Study of the Anti-Anemic Activity of the Aqueous Extract of the Leaves of *Petroselinum Crispum* (Apiaceae) In the White Mouse (*Mus Musculus*) Of SWISS Strain. *J Blood Disord Ther*. 2022;2:101.
55. Al-Daraji HJ, Al-Mashadani HA, Al-Hassani AS, Mirza HA, Al-Hayani WK. The Influence of parsley (*Petroselinum crispum*) as feed additive on hematological traits of local Iraqi geese. *Advances in Nutrition Research*. 2012;1(1):1-5.
56. Kamath S, Lip GYH. Fibrinogen: biochemistry, epidemiology and determinants. *QJM: An International Journal of Medicine*. 2003;96(10):711-29.
57. Stang LJ, Mitchell LG. Fibrinogen. *Haemostasis: Methods and Protocols*. 2013:181-92.
58. Rea CJ, Sorensen B, Ingerslev J, Laursen P. Fibrinogen: a procoagulant and an anticoagulant. *Blood*. 2011;118(21):384.
59. Chao FC, Tullis JL, Kenney DM, Conneely GS, Doyle JR. Concentration effects of platelets, fibrinogen and thrombin on platelet aggregation and fibrin clotting. *Thrombosis and Haemostasis*. 1974;32(04):216-31.