

## Original Research Article

# **Synthesis of (E)-N'-(pyridine-2-ylmethylene) picolino hydrazide with promising antimicrobial and anti-inflammatory activities along with their *in-silico* studies**

### **ABSTRACT**

In recent years, there has been a significant amount of interest in hydrazide-hydrazone compounds as a result of their extensive use in the pharmaceutical industry. In this study, the new compound *(E)-N'-(pyridin-2-ylmethylene) picolino hydrazide*, with the formula  $C_{20}H_{19}N_3O_2$ , was synthesised and confirmed using different spectroscopic techniques. The newly synthesised compound was tested for anti-inflammatory and antibacterial properties *in vitro*. The results of an *in vitro* anti-inflammatory activity test using the denaturation of bovine serum albumin as the measurement kit revealed a highly effective inhibitory effect. In addition, the results of the antimicrobial test showed that the compound demonstrated a maximum zone of inhibition (18.2 mm) against the *Staphylococcus aureus* strain at a higher concentration (32  $\mu\text{g/ml}$ ) of the tested sample (32  $\text{g/ml}$ ), which is greater than the standard drug (ciprofloxacin: 17.5 mm). The antifungal activity of *(E)-N'-(pyridin-2-ylmethylene) picolinohydrazide* indicated a maximum inhibition zone of 9.2 mm at a higher concentration (32  $\mu\text{g/ml}$ ) of the tested sample, which is comparable to that of the standard drug (ketoconazole: 9.0 mm). The anti-inflammatory activity performed by the carrageenan-induced paw edema method also gave a promising result. Molecular docking analysis revealed that the active compounds had a higher docking score and interacted effectively with the target protein. *In silico* molecular docking, results revealed the COX-5 and LL-37 (residues 17–29) proteins mainly depends on H-bonds between the pyridine groups of the synthesized ligand and the catalytic residues of the binding site,  $\pi$ -cation/anion interactions, (iii)  $\pi$ -alkyl interactions, and formation of carbon-hydrogen bond,  $\pi$ -  $\pi$  bond that stabilizes the interaction with the active site. Among the analogs of the hydrazine hydride class, the newly

synthesized compound: (*E*)-*N'*-(pyridin-2-ylmethylene) picolino hydrazide proved to be a vital bioactive molecule.

**Keywords:** Hydrazide-hydrazone, Antimicrobial, Anti-inflammatory, Docking

## 1. Introduction

Inflammation can be induced by a variety of factors, including bacterial and viral infections, allergies, irritants, toxic substances, and injury to any tissue in the body (Zabrodskii). Inflammation can be generated from the outside (exogenously) or the inside (endogenously) (Gallucci and Matzinger; Lon, Liu, and Jusko). This may also involve trauma, surgery, autoimmunity, adult respiratory distress syndrome, prolonged exposure to industrial toxins, and reoxygenation damage (Weigand *et al.*, Guoquiand Li *et al.*, Chengjian Lou *et al.*).

The inflammatory response can be sped up by molecules that are readily available in the body, such as histamine, prostaglandins, leukotrienes, oxygen- and nitrogen-based free radicals, serotonin, bradykinin, and interleukins. An excessive inflammatory response can increase the risk of having a stroke or developing heart disease, as well as lupus, cancer, tissue injury, physiological decompensation, organ damage, and even mortality (Ward and Lentsch). Researchers in the field of medicinal chemistry face a problem when it comes to the hunt for more potent anti-inflammatory drugs, because of the toxic chemical involved with the inflammation process. Aspirin, indomethacin, flufenamic acid, ibuprofen, and a host of other anti-inflammatory medications are acidic in nature (Whitehouse; John R Vane and Renia M Botting; John R Vane and Regina M Botting). One of the most important classes of medications for treating inflammation is known as non-steroidal anti-inflammatory drugs (NSAIDs). These drugs work by reducing inflammation in the body by inhibiting an enzyme called cyclooxygenase (COX 5), which is involved in the production of prostaglandins (Kovala-Demertzi; Brideau *et al.*).

In the process of organic synthesis, hydrazines and the derivatives of these compounds make up an important class of substances that have been utilized in a variety of diverse applications (Rollas, Gulerman and Erdeniz; Gürsoyet *al.*, Nabel B. Ayrimet *al.*). Hydrazines have traditionally been employed as reagents for the derivatization and analysis of carbonyl compounds. In recent years, the N-N bond has emerged as a key structural motif in numerous bioactive drugs. A growing number of N-N bond-containing heterocycles and

peptidomimetics have found their way into commercial use as therapeutic medicines. Particularly, N-N bond-containing heterocycles and peptidomimetics are increasingly utilized as medicinal and agricultural agents in the industry (Vicini *et al.*; Mamolo *et al.*). In addition, hydrazide-hydrazones have gained prominence due to their various biological applications, which include antibacterial (Alaa, El-Subbagh and Kunieda), anti-HIV (Metobo *et al.*), anticancer (Kamat *et al.*), anti-inflammatory, antioxidant, and anticoagulant (de Candia *et al.*; Chaban *et al.*) activities.

Infections caused by microorganisms have become one of the primary causes of death in recent decades (RoukiatouTraoré, Emmanuel Sampo and Savadogo; Laurretta). Researchers are actively looking for antimicrobial candidates that are more potent and effective against bacteria that are both drug-sensitive and drug-resistant to overcome current challenges. Bruno Leal *et al.* (Leal *et al.*), Jacques *et al.* (Bompart *et al.*), Rakha *et al.* (Rakha *et al.*), and Altundas *et al.* (Altundas, Ayvaz and Logoglu) found numerous antibacterial candidates including pyridine. The pyridine nucleus has been regarded as one of the most potent bioactive scaffolds in many biological systems based on the reported literature.

Various anti-inflammatory and antibacterial drugs containing pyridine have also been identified in the literature. For instance, Abdelmonsef *et al.*, (Abdelmonsef *et al.*), Santosh *et al.*, (Undare *et al.*), and Santhisudha *et al.*, (Sarva *et al.*) reported on anti-inflammatory compounds derived from pyridine. Huan *et al.*, (Du *et al.*), Yusufzai *et al.*, (Khan Yusufzai *et al.*) Desai *et al.*, (Desai *et al.*) Mathew (Bijo *et al.*) and Zahra, *et al.* (Najafi *et al.*) have discovered antibacterial agents. In light of these findings, we also sought to synthesize bioactive scaffolds consisting of hydrazides connected to picolinic-containing pyridine. The newly synthesized compound such as: (*E*)-*N'*-(pyridin-2-ylmethylene) picolino hydrazide have been characterized and confirmed by different spectroscopic techniques. In addition, *in vitro* antibacterial and *in vivo* anti-inflammatory properties, as well as *in silico* molecular docking investigations, were methodically covered here.

## 2. Methods


### 2.1. Chemicals, solvents, and instrumentations

All of the chemicals and solvents used in this study were acquired from Hi-Media laboratory, India, and Spectrochem Pvt. Ltd., India, and was utilized untreated. The reactions were conducted according to standard protocol. In an open capillary tube, the uncorrected melting points of newly synthesized compounds were determined. Thin-layer

chromatography (TLC, silica gel 60 F254) was used to evaluate the reaction's progress and the purity of the result. The solvent mixture of ethyl acetate and hexane was used as the mobile phase, and it was monitored at ~254 nm under UV light. The Bruker-400 spectrometer was used to record  $^1\text{H}$  NMR spectra, using  $\text{DMSO-d}_6$  as a solvent and TMS as an internal standard (the chemical shift in  $\delta$  ppm and coupling constants (J) were represented in parts per million (ppm) and Hz, respectively).

## 2.2. Synthesis of (*E*)-*N'*-(pyridine-2-ylmethylene) picolino hydrazide.

The picolino hydrazide (0.137g, 1.0 mM) and picolinaldehyde (0.107g, 1.0 mM) are dissolved in methanol and refluxed for 2 hrs. This reaction mixture was poured into ice-cold distilled water and kept for 30 mins at room temperature. The precipitate formed was filtered and air dried and re-crystallized using THF.



*N'*-[(*E*)-(pyridin-2-yl)methylidene]pyridine-2-carbohydrazide

(74% Yield)

### Scheme 1. Synthetic Pathway of the (*E*)-*N'*-(pyridine-2-ylmethylene) picolinohydrazide

## 2.3. *In vitro* antimicrobial activity

Human pathogens such as *Staphylococcus aureus* (ATCC 25923), *Escherichia Coli* (ATCC 25922), and *Candida albicans* (ATCC 22972) were used for the microbial studies. The clinical isolates were obtained from the National Chemical Laboratories in Pune, India. The antimicrobial activity of all samples was evaluated on Muller Hinton agar (MHA) for bacteria and sterile Sabouraud dextrose broth (SDB) for the fungi using the disc diffusion method (Bharadwaj et al.). The discs were prepared by punching sterile Whatman filter paper with a sterile punching machine. To determine antimicrobial activity, each standard antibiotic disc was dipped in DMSO containing synthesized organic compound at the following

concentrations: 4, 8, 16, and 32 µg/ml of test compounds was made by adding 10 µl of different concentration and allowed to dry (RT for 30 min). The MHB and SDB tubes with and without test samples were used as the control. A standard antibacterial drug (ciprofloxacin) and an antifungal drug (ketoconazole) were used for comparison. The bacterial and fungal strains were added to the nutrient broth and incubated on a rotary shaker at 200 rpm and 37°C for 24 hours. Post incubation, the optical density at 600nm was adjusted to get an inoculum size of approximately 10<sup>7</sup> CFU/ml and used for the assay. On MHA plates, the turbidity of broth cultures was compared to that of standard 0.5 McFarland solutions, along with a standard antibiotic disc and a prepared disc containing different amounts of test samples. After 24 hours at 37°C, digital calipers were used to measure the inhibitory zones (Mitutoyo Rochester, New York). Three experiments were conducted, and the average zone diameter was determined.

#### 2.4 Details animal grouping and *in vivo* anti-inflammatory activity

Male Swiss albino mice of weight 100 to 120 g were considered for the *in vivo* anti-inflammatory studies. With sufficient water, standard *ab libitum* diet was employed for about 7 days prior to the experiments. The animals were allowed for fasting overnight at the start of the experiments. Animal Ethical committee guidelines from Chromed Biosciences Pvt. Ltd. Hirehalli Industrial Area, Tumkur with the clearance number CBPL-IAEC-014/07/2022 were followed and the good care was taken for the animals.

For *in vivo* anti-inflammatory activity, carrageenan induced paw edema method. The animals were divided into six groups with six animals each in a group. Group I was administered with only DM water (10 ml/kg p.o/day). Group II was administered with standard drug indomethacin (10 mg/kg). Group III to Group VI animals were administered with synthesised (*E*)-*N'*-(pyridine – 2 - ylmethylene) picolinohydrazide(10, 25, 50 and 100 mg/kg, respectively). Using Plethysmometer, initial paw volume was measured and then the drugs were induced accordingly. The drug was administered and after 30 min, 1 % of carrageenan (25 µl) was injected in saline into the sub-plantar region of the right hind paws. The volume of paw was then recorded at 1, 2, 4, 6, and 24 h. The final edema was calculated using initial and final paw volumes at each time points.

#### 2.5 Statistical Analysis

SPSS version 20.0 were utilized for the statistical analysis. The studies were run in triplicate and the findings were represented as mean standard deviation (MSD) by one-way

ANOVA. Turkey's Multiple comparisons were utilized to compare standard and synthesized compounds. Using Pearson's correlation analysis,  $p < 0.01$  was considered statistically significant.

### 2.6. Molecular docking simulation

The structure of proteins cyclooxygenase-5 (COX-5; PDB code: 1EQG) and human antimicrobial peptide LL-37(residues 17-29) was collected from Research Collaboratory for Structural Bioinformatics Protein Data Bank (RCSB PDB) (Adams et al.). Energy minimization of ligands was accomplished in Open Babel (version 2.4.0) (O'Boyle et al.). The docking approach was validated by re-docking the native ligands into the binding pockets of the corresponding proteins. The docking of the target protein-ligand was performed using Auto-Dock Vina (version 1.1.2) (Trott and Olson). Docking simulations were performed with a grid whose centers were positioned at the active site of the individual proteins and whose dimensions were 25,00, 25,00, and 25.00 Å. Discovery Studio Client 2019 has examined the interactions between ligands and macromolecular targets. In this docking technique, several ligand conformations were generated, and the final energy refinement of the ligand posture took place. The Dock score of the best poses docked into the target protein was determined for the test compound.

## 3. Results and Discussion

### 3.1. Chemistry

The formation of the novel compound was confirmed by  $^1\text{H}$  NMR (Fig. 1). The formation of  $-\text{CH}=\text{N}$  was confirmed by a singlet at  $\delta$  8 ppm.

White solid; Yield: 74.00%;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz): $\delta$  7.415-7.449 (1H, ddd,  $J = 7.2, 4.8, 1.2$  Hz), 7.680-7.714 (1H, ddd,  $J = 8.0, 3.2, 1.6$  Hz), 7.876-7.918 (1H, ddd,  $J = 15.6, 7.6, 1.2$  Hz), 7.993-8.013 (d,  $J = 8.0$ , Hz), 8.054-8.097 (1H, ddd,  $J = 15.6, 8.0, 2.0$  Hz), 8.144-8.164 (d,  $J = 8.0$  Hz), 8.620-8.631 (d,  $J = 4.4$  Hz), 8.697 (1H, S), 8.738-8.749(1H d  $J=4.4$ ), 12.506 (1H S).



A fungus strain known as *Candida albicans* was utilized. The growth of the fungus was carried out in a Sabouraud dextrose agar (SDA) medium, and testing was carried out in a Sabouraud dextrose broth (SDB) media. The cultures and the viable count were performed using the same method that was used in the antibacterial tests, except for the temperature, which should be kept at  $28 \pm 1^\circ\text{C}$  for approximately 72 hours. For the antifungal experiments, the same concentration of the test chemical, solvent (DMSO), and standard (ketoconazole) as was employed in the prior preparations was utilized. The antifungal activity of synthesized (*E*)-*N'*-(pyridine – 2 – ylmethylene) picolinohydrazide demonstrated a minimal zone of inhibition at a higher concentration of the tested sample, i.e., 9.5 mm which is almost comparable to that of the standard (ketoconazole: 9.0 millimeters). Table 1 presents the results of measurements made on the antifungal activity of (*E*)-*N'*-(pyridine – 2 -ylmethylene) picolinohydrazide. Fig. 2 depicts typical images of kirby-bauer disk diffusion technique. Similar behavior of antimicrobial activity was reported for pyridine-based hydrazide compounds in the literature (Rollas, Gulerman and Erdeniz; Vicini et al.; Halli, Sumathi and Kinni).

**Table 1:** Antimicrobial activity (zone of inhibition) of as-synthesized compound by disc diffusion method

Organisms name	Concentration ( $\mu\text{g/ml}$ )	Mean ZOI (mm)
<i>S.aureus</i>	32	18.2
<i>E.coli</i>	32	No zone of inhibition
<i>C. albicans</i>	32	9.5




Fig. 2: Photographic images of antimicrobial activities by zone of inhibition of *(E)-N'-(pyridin-2-ylmethylene) picolinohydrazide*; viz., *S. aureus* (a-a2); *E. coli*(b-b2) and *C. albicans* (c-c2)

### 3.3 *In vivo* anti-inflammatory activity

The synthesised *(E)-N'-(pyridin-2-ylmethylene) picolinohydrazide* was coded as NN. The results are provided in Fig. 3. A progressive increment of paw thickness was observed and was greater for the control group at four hours indicating the inflammation induced by 1 % carrageenan. However, with the increase in the dosage of the drug, the paw edema decreased indicating the anti-inflammatory activity at 4 hours. The edema further decreased with increasing 24 hours and remained almost equal. At 4 hours the control group showed 0.36 ml of paw edema, group II with control drug showed 0.23 ml, whereas, the groups III to VI, with increased concentrations of drug showed average of 0.35, 0.31, 0.28 and 0.26 ml of paw edema, respectively, confirming their anti-inflammatory activity.





Fig. 3: Anti-inflammatory activity of (*E*)-*N'*-(pyridin-2-ylmethylene) picolinohydrazide coded as NN, by carrageenan induced paw edema method.

### 3.4. Molecular Docking Studies

The docking study was performed using AutodockVina (version 1.1.2), which is a flexible docking program, that can take into consideration the flexibility of the ligand as well as the flexibility of the active site residues of COX 5 (Trott and Olson). Docking tests were carried out to evaluate the potential binding manner of the significantly active chemical against the COX-5 (PDB code: 1EQG) proteins, which served as the target of the investigation. Before beginning the docking tests on the test compound, the root mean square deviation (RMSD) of the X-ray pose and re-docked pose of the co-crystallized ligand in the target protein were analyzed. These data were also obtained in the protein data bank ([www.rcsb.org](http://www.rcsb.org)). The fact that the RMSD was calculated to be 0.20 Å suggested that the docking methodology may be trusted for the docking experiments of the test chemical. After careful inspection of the co-crystal ligands (3D and 2D) poses (Fig. 4), it was found

that celecoxib revealed four hydrogen bond interactions within the target, with the maximum docking score being -4.8 kcal/mol and the energy is 60.10 kcal/mol. The following amino acid residues actively engaged in the formation of hydrogen bonds between celecoxib and amino acid residues such as ALA-443, ARG-185, and PHE-504. In addition to the hydrogen bond, celecoxib interacted with the amino acid residues through two aromatic bonds. The interaction between the pyridine moiety of celecoxib and the pyrazole part of celecoxib was taking place through ASP-393. In addition, the active compound also exhibited four aromatic bond interactions with the amino acid residue ILE:442, ASP:393, PHE:404, and ALA:443 through the pyridine moiety, as shown in Fig 5 and 6. The compound exhibited a docking score of -4.3 kcal/mol and docking energy of -42.10 kcal/mol, which is comparatively less than those of the standard drug celecoxib. Relationships between the structure and the activity of the newly synthesised hydrazine hydride: *((E)-N'-(pyridine - 2 - ylmethylene) picolinohydrazide)* (possible inhibitors of COX 5), a docking study demonstrated that the inhibitory activity of *((E)-N'-(pyridine - 2 - ylmethylene) picolinohydrazide)* towards COX 5 is dependent on (i) hydrogen bonds between the pyridine groups of the synthesized ligands and the catalytic residues of the binding site, (ii)  $\pi$ cation/anion interactions, (iii)  $\pi$ -alkyl interactions, and (iv) formation of carbon-hydrogen bond, (v)  $\pi$ - $\pi$  bond that stabilizes the interaction with the active site.




**Fig. 4.** COX 5 - hormone interactions; the hormone is stabilized mainly by Amide (i) hydrogen bonds between the pyridine groups of the synthesized ligands and the catalytic residues of the binding site, (ii)  $\pi$ cation/anion interactions, (iii)  $\pi$ -alkyl interactions, and (iv) formation of carbon-hydrogen bond, (v)  $\pi$ -  $\pi$  bond that stabilizes the interaction with the active site.

The potent inhibitory activity of the synthesized compound: (*E*)-*N'*- (pyridine – 2 - ylmethylene)picolinohydrazide) as new antimicrobial agents prompted us to study the docking of these derivatives inside the active sites of the human LL 37 (17-29) protein. The human LL-37 (hLL-37), also known for its ability to self-assemble, is a cleavage product of the hCAP-18 protein. It is recognized to play a vital part in the body's first line of defense against microbial infections (Xhindoli et al.). LL-37, also known as hLL-37, has a role in replication and transcription, and it catalyzes the formation of negative supercoils in the circular DNA of microbes. Additionally, LL-37 (hLL-37) is a known target for antimicrobial drugs because its inhibition results in the death of microbes (Xhindoli et al.).

In our attempt to explain the activity of the above-synthesized compound, we docked within the active site of LL-37 (hLL-37) protein (residues 17-29). The binding pocket of the enzyme includes the following residues LYS 2, Arg7, and Arg3. Docking studies revealed that the compound showed two H-bonds between the pyridine groups of the synthesized ligands and

the catalytic residues of the binding site,  $\pi$ cation/anion interactions,  $\pi$ -alkyl interactions, and formation of carbon-hydrogen bond and  $\pi$ -  $\pi$  bond that stabilizes the interaction with the active site (Fig. 5). Docking results indicated that LYS 2, Arg7, and Arg3 play important roles in the activity. The explanation of the biological activity of the new compound may be attributed to the possibility of inhibition of these three amino acids.



**Fig. 5.** LL 37 (17-29) - hormone interactions; the hormone is stabilized mainly by Amide (i) hydrogen bonds between the pyridine groups of the synthesized ligands and the catalytic residues of the binding site, (ii)  $\pi$ cation/anion interactions, (iii)  $\pi$ -alkyl interactions, and (iv) formation of carbon-hydrogen bond, (v)  $\pi$ -  $\pi$  bond that stabilizes the interaction with the active site.

#### 4. Conclusion

In the present study, we synthesized a novel anti-inflammatory and antimicrobial agent: *(E)*-*N'*-(pyridine - 2- ylmethylene) picolinohydrazide. The in vitro anti-inflammatory study of the synthesized compound revealed that are significant inhibitors among the other series of hydrazine hydrides reported in the literature. The *in vitro* antibacterial experiment revealed that the synthesized compound is effective at a higher dose (32 g/ml) against

*Staphylococcus aureus* (zone of inhibition 19.5 mm) and *Candida albicans* (zone of inhibition 9.5 mm) strains. The anti-inflammatory activity was confirmed by the reduced paw edema at 4 hours in the groups administered with the synthesized drug. *In silico* molecular docking study also evaluates the efficiency of interactions with target molecules. Based on our findings, we may conclude that this is a promising path for the development of new antibacterial and anti-inflammatory drugs.

## References

- Abdelmonsef, Aboubakr H, et al. "A Search for Antiinflammatory Therapies: Synthesis, in Silico Investigation of the Mode of Action, and in Vitro Analyses of New Quinazolin-2, 4-Dione Derivatives Targeting Phosphodiesterase-4 Enzyme." *Journal of Heterocyclic Chemistry* 59.3 (2022): 474-92. Print.
- Adams, PLR, et al. "Using the Pair Angle Distribution Function for Analysing Protein Structure." *Acta Crystallogr. Found. Adv* 77 (2021): C574. Print.
- Alaa, A-M, Hussein I El-Subbagh, and TakehisaKunieda. "Lewis Acid-Promoted Transformation of 2-Alkoxy pyridines into 2-Aminopyridines and Their Antibacterial Activity. Part 2: Remarkably Facile C–N Bond Formation." *Bioorganic & medicinal chemistry* 13.16 (2005): 4929-35. Print.
- Altundas, Aliye, SelcukAyvaz, and ElifLogoglu. "Synthesis and Evaluation of a Series of Aminocyanopyridines as Antimicrobial Agents." *Medicinal chemistry research* 20.1 (2011): 1-8. Print.
- Bharadwaj, S Shashidhar, et al. "Design, Synthesis and Pharmacological Studies of Some New Quinoline Schiff Bases and 2, 5-(Disubstituted-[1, 3, 4])-Oxadiazoles." *New Journal of Chemistry* 41.16 (2017): 8568-85. Print.
- Bijo, Mathew, et al. "Synthesis, Characterisation of Some 2-Azetidinone Derivatives from 2-Aminopyridine and Evaluation of Their Antimicrobial Activity." *Der Pharma Chemica* 2.6 (2010): 238-42. Print.
- Bompart, Jacques, et al. "Synthèse Et Activité Anti-Bactérienne D'acides Dihydro-4, 7 Éthyl-7 Oxo-4 Thiéno [2, 3-B] Pyridine Carboxyliques-5 li." *European journal of medicinal chemistry* 23.5 (1988): 457-64. Print.
- Brideau, C, et al. "A Human Whole Blood Assay for Clinical Evaluation of Biochemical Efficacy of Cyclooxygenase Inhibitors." *Inflammation Research* 45.2 (1996): 68-74. Print.
- Chaban, Taras I, et al. "Synthesis, Molecular Docking and Biological Properties of Novel Thiazolo [4, 5-B] Pyridine Derivatives." *Acta Chimica Slovenica* 67.4 (2020): 1035-43. Print.
- Chengjian Lou, Marong Fang, Shan Ye, Zhangly Fang, Nashwa Amin, YiliChen. "Fluoxetine protects against inflammation and promotes autophagy in mice model of post-traumatic stress disorder" *Behavioural Brain Research* 433 (2022): 114004.
- de Candia, Modesto, et al. "Synthesis and Biological Evaluation of Direct Thrombin Inhibitors Bearing 4-(Piperidin-1-Yl) Pyridine at the P1 Position with Potent Anticoagulant Activity." *Journal of Medicinal Chemistry* 56.21 (2013): 8696-711. Print.
- Desai, NC, et al. "Synthesis and Characterization of Oxazine Bearing Pyridine Scaffold as Potential Antimicrobial Agents." *Synthetic Communications* 47.24 (2017): 2360-68. Print.
- Du, Huan, et al. "Design, Synthesis, Crystal Structure and in Vitro Antimicrobial Activity of Novel 1, 2, 4-Triazolo [1, 5-a] Pyrimidine-Containing Quinazolinone Derivatives." *Molecular Diversity* 25.2 (2021): 711-22. Print.
- Gallucci, Stefania, and Polly Matzinger. "Danger Signals: Sos to the Immune System." *Current opinion in immunology* 13.1 (2001): 114-19. Print.

Guoqiand Li, LetiDuan, Fengbiao Yang, Liang Yang, Yajun Deng, Yue Yu, Yanlong Xu, Yinian Zhang. "Curcumin suppress inflammatory response in traumatic brain injury via p38/MAPK signaling pathway" *Phytotherapy Research* 36 (2022): 1326-1337.

Gürsoy, A., et al. "Synthesis of Some New Hydrazone-Hydrazone, Thiosemicarbazides, Thiadiazoles, Triazoles and Their Derivatives as Possible Antimicrobials." *Die Pharmazie* 45.4 (1990): 246-50. Print.

Halli, MB, RB Sumathi, and MallikarjunKinni. "Synthesis, Spectroscopic Characterization and Biological Evaluation Studies of Schiff's Base Derived from Naphthofuran-2-Carbohydrazone with 8-Formyl-7-Hydroxy-4-Methyl Coumarin and Its Metal Complexes." *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 99 (2012): 46-56. Print.

Kamat, Vinuta, et al. "Pyridine-and Thiazole-Based Hydrazides with Promising Anti-Inflammatory and Antimicrobial Activities Along with Their in Silico Studies." *ACS omega* 5.39 (2020): 25228-39. Print.

KhanYusufzai, Samina, et al. "Design, Characterization, in Vitro Antibacterial, Antitubercular Evaluation and Structure-Activity Relationships of New HydrazinylThiazolylCoumarin Derivatives." *Medicinal Chemistry Research* 26.6 (2017): 1139-48. Print.

Kovala-Demertzi, Dimitra. "Recent Advances on Non-Steroidal Anti-Inflammatory Drugs, Nsaids: Organotin Complexes of Nsaids." *Journal of Organometallic Chemistry* 691.8 (2006): 1767-74. Print.

Lauretta, UmeakuChinyeluNkiru and EzechinwoyeChiemelie. "Antibacterial Effect of Honey and ChromolaenaOdorata on Escherichia Coli and Staphylococcus Aureus Isolated from Wound Infection." *Acta Scientific Microbiology* 4.8 (2021): 158-64. Print.

Leal, Bruno, et al. "Antibacterial Profile against Drug-Resistant Staphylococcus Epidermidis Clinical Strain and Structure-Activity Relationship Studies of 1h-Pyrazolo [3, 4-B] Pyridine and Thieno [2, 3-B] Pyridine Derivatives." *Bioorganic & medicinal chemistry* 16.17 (2008): 8196-204. Print.

Lon, Hoi-Kei, Dongyang Liu, and William J Jusko. "Pharmacokinetic/Pharmacodynamic Modeling in Inflammation." *Critical Reviews™ in Biomedical Engineering* 40.4 (2012). Print.

Mamolo, Maria Grazia, et al. "Synthesis and Antimycobacterial Activity of [5-(Pyridin-2-Yl)-1, 3, 4-Thiadiazol-2-Ylthio] Acetic Acid Arylidene-Hydrazone Derivatives." *Il Farmaco* 56.8 (2001): 587-92. Print.

Metobo, Sammy E, et al. "Design, Synthesis, and Biological Evaluation of Novel Tricyclic Hiv-1 Integrase Inhibitors by Modification of Its Pyridine Ring." *Bioorganic & medicinal chemistry letters* 16.15 (2006): 3985-88. Print.

Mizushima, Y, and M Kobayashi. "Interaction of Anti-Inflammatory Drugs with Serum Proteins, Especially with Some Biologically Active Proteins." *Journal of Pharmacy and Pharmacology* 20.3 (1968): 169-73. Print.

Nabel B. Ayrim, Asim A. Balakit, Souad J. Lafta. "Synthesis, Biological Activity, and Molecular Docking Studied of New Substituted Hydrazones" *Journal Of Kufa For Chemical Sciences* 2 (2022) 582-605.

Najafi, Zahra, et al. "Synthesis and in Vitro Cytotoxic Activity of Novel Triazole-Isoxazole Derivatives." *Journal of Heterocyclic Chemistry* 52.6 (2015): 1743-47. Print.

O'Boyle, Noel M, et al. "Open Babel: An Open Chemical Toolbox." *Journal of cheminformatics* 3.1 (2011): 1-14. Print.

Rakha, TH, et al. "Synthesis, Characterization, Dft and Biological Studies of (Z)-N'-(2-Oxoindolin-3-Ylidene) Picolinohydrazone and Its Co (Ii), Ni (Ii) and Cu (Ii) Complexes." *Journal of Molecular Structure* 1062 (2014): 96-109. Print.

Rollas, Sevim, NehirGulerman, and HabibeErdeniz. "Synthesis and Antimicrobial Activity of Some New Hydrazones of 4-Fluorobenzoic Acid Hydrazone and 3-Acetyl-2, 5-Disubstituted-1, 3, 4-Oxadiazolines." *Il Farmaco* 57.2 (2002): 171-74. Print.

- RoukiatouTraoré, CheiknaZongo, ArounaOuedraogo,,Mahamadi Sore Emmanuel Sampo, BoubacarYaro, Yves Traoré, and and Aly Savadogo. "In Vitro Antimicrobial Activity of Crude Extracts from VetiveriaNigritana (Benth.) Stapf, Mitragynalnermis (Willd.)
- Kuntze, KalanchoeCrenata (Andr.) Haw. Against Methicillin-Resistant Staphylococcus Aureus." *Acta Scientific Microbiology* 4.5 (2021): 56-67. Print.
- Sarva, Santhisudha, et al. "Synthesis, Antibacterial and Anti-Inflammatory Activity of Bis (Indolyl) Methanes." *Chinese Chemical Letters* 27.1 (2016): 16-20. Print.
- Trott, Oleg, and Arthur J Olson. "AutodockVina: Improving the Speed and Accuracy of Docking with a New Scoring Function, Efficient Optimization, and Multithreading." *Journal of computational chemistry* 31.2 (2010): 455-61. Print.
- Undare, Santosh S, et al. "One-Pot Synthesis and in Vivo Biological Evaluation of New Pyrimidine Privileged Scaffolds as Potent Anti-Inflammatory Agents." *Research on Chemical Intermediates* 42.5 (2016): 4373-86. Print.
- Vane, John R, and Regina M Botting. "Mechanism of Action of Nonsteroidal Anti-Inflammatory Drugs." *The American journal of medicine* 104.3S1 (1998): 2S-8S. Print.
- Mechanism of Action of Aspirin-Like Drugs*. Seminars in arthritis and rheumatism. 1997. Elsevier. Print.
- Vicini, Paola, et al. "Hydrazones of 1, 2-Benzisothiazole Hydrazides: Synthesis, Antimicrobial Activity and Qsar Investigations." *European journal of medicinal chemistry* 37.7 (2002): 553-64. Print.
- Ward, Peter A, and Alex B Lentsch. "The Acute Inflammatory Response and Its Regulation." *Archives of surgery* 134.6 (1999): 666-69. Print.
- Weigand, Markus A, et al. "The Systemic Inflammatory Response Syndrome." *Best Practice & Research Clinical Anaesthesiology* 18.3 (2004): 455-75. Print.
- Whitehouse, MW. "Some Biochemical and Pharmacological Properties of Anti-Inflammatory Drugs." *Fortschritte der Arzneimittelforschung/Progress in Drug Research/Progrès des recherches pharmaceutiques* (1965): 321-429. Print.
- Xhindoli, Daniela, et al. "The Human Cathelicidin LI-37—a Pore-Forming Antibacterial Peptide and Host-Cell Modulator." *Biochimica et BiophysicaActa (BBA)-Biomembranes*1858.3 (2016): 546-66. Print.
- Zabrodskii, Pavel F. "The Effects of A7n-Acetylcholine, B2-Adrenergic Receptors Agonists, Nf-Kb, Stat3 Inhibitors on the Implementation of the Cholinergic Anti-Inflammatory Pathway in Sepsis." *Acta Scientific Microbiology* 4.7 (2021): 16-20. Print.

## Abbreviations

NSAIDs: Non-steroidal Anti-inflammatory Drugs; Cyclooxygenase 5: COX 5; Thin-layer chromatography:TLC; DMSO: Dimethyl Sulfoxide; NMR : Nuclear Magnetic Resonance; Ppm: Parts per million; SDB: Saboraud dextrose broth (SDB); RT : Room Temperature; MSD: Mean Standard Deviation; RCSB PDB : Structural Bioinformatics Protein Data Bank; THF: Tetrahydrofuran, SDA: Sabouraud dextrose agar