

## Studies on Diversity of Some Local Jamun (*SyzygiumCuminiSkeels*) Genotypes

### ABSTRACT

The jamun (*SyzygiumcuminiSkeels*) is an important indigenous minor fruits of tropical and sub-tropical region and it is tall, handsome-evergreen tree of India. The western dry tract of West Bengal has the greater diversity of different fruit crops including jamun. The variation in different jamun genotypes in this zone is exclusively reported. Thus the present experiment entitled “Studies on diversity of some local jamun genotypes(*SyzygiumCuminiSkeels*)” has been undertaken to express the diversity of jamun fruits within the available genotype resources during the period of 2019 to 2021 under the Department of Horticulture & Postharvest Technology, PalliSikshaBhavana (Institute of Agriculture), Visva-Bharati, Sriniketan, West Bengal. The experiment was conducted by selecting 15 different jamun genotypes from Sriniketan and adjoining villages (viz. Andrews Pally, Rathindra Pally, Ballavpur, BahadurpurPashchim, Sonajhuri) and jamun fruits were brought to record their physical as well as bio-chemical character at laboratory of Department. Present study reveals a wide and significant variations in physical and bio-chemical characters of fruits has been observed among 15 different jamun genotypes. Among all genotypes G-14 was superior on the basis of fruit weight (3.16g), fruit diameter (17.46mm), and greater pulp content (2.34g) and average total sugar content (16.26%) content. Owing to moderate fruit size (21.89mm in length and 15.64mm in diameter)G-4 genotype is also superior on the basis of higher TSS (25.90□B), total sugar (13.65%) and reducing sugar (8.23%). So, the overall findings of the present study conclude that genotypes G-14 and G-4 was-are superior over the 15 local jamun genotypes.

**Key words:** Jamun, Genotypes, Fruit Physico- chemical Properties.

Formatted: Font: 12 pt

Formatted: Font: Italic

### INTRODUCTION

Jamun/ Indian blackberry/ black plum (*SyzygiumcuminiSkeels*.), is an important evergreen tropical and minor fruit which family belongs to ismyrtaceae and other common

names of jamun ~~is~~ are java plum/ ~~K~~Kalajam/ ~~p~~Phalinda/ Rajamun/ ~~d~~Diabetiesfibre/ ~~r~~Rose apple/ ~~p~~Pitanga cherry. It is believed to be indigenous to India, Burma, Ceylon and Andaman Islands (Zeven and de wet., 1982). Jamun was introduced from India and Tropical Asia to Southern Africa for its edible and attractive fruits. Jamun is widely distributed in India, Malaysia, Australia and USA. Jamun is grown mostly tropical and sub-tropical countries in the world. India rank 2<sup>nd</sup> in production of jamun in the world after China. In India, jamun grown almost throughout the country, leading producer states ~~is~~ are Maharashtra, Uttar Pradesh, Tamilnadu, Gujarat and others. Maharashtra is the leading producer of jamun in India. Jamun is very important fruit plants of both food and medicinal value. Jamun fruits have high demand for table and for the preparation of wide range of products like squash, R.T.S., syrup, jam, jellies, juice, vinegar etc (Patilet *et al.*, 2012). Jamun plants have ornamental value, inflorescence is emerged from leaf axils or branch lets and floral parts arise in the acropetal successions. Jamun flowers are bisexual and mostly light yellow in color and flowering period is between first week of March to April and bear fruits from May to July. In jamun cross pollination is ocured and mode of pollination is honeybees, house flies and wind, ~~and dropping is a serious problem of flowers and fruits~~ Flower and fruit drop is a serious problem in jamun (Singh *et al.*, 2019). The medicinal value of seeds of jamun (alkaloids- jambosin and glycosides- jambolin or antimellin) which retard the conversion of starch into sugars and its seeds powder quickly reduce the quantity of sugar in urine, so it's very useful for diabetic patients. Ripe fruits of jamun are highly juicy, with a pleasant, slightly bitter, astringent taste and consumed as a fresh or in processed products. It is also used to make wine in Phillipines. Wood of jamun is resistance to water and termite attacks, is used to install motors in the wells and being fast growing tree, it provide excelled fire wood and charcoal to the rural population in India (Chaudhary and mukhopadhyay., 2012) and it's wood is also used for the railway sleepers. In terms of medicinal value, jamun has astringent, stomachic, carminative, anti-scorbutic and diuretic properties (Singh, I.S. 2001).

---

## MATERIALS AND METHODS

---

The experiment was directed by selecting ~~different~~ 15 different jamun genotypes from Sriniketan and adjoining villages (*viz.* Andrews Pally, Rathindra Pally, Ballavpur, Sonajuhri, BahadurpurPashchim) and for the physical and biochemical estimation brought the fruits at the laboratory of Department of Horticulture and Postharvest Technology,

Formatted: Font: Italic

PalliSikshaBhavana (Institute of Agriculture), Visva-Bharati, Sriniketan, West Bengal. For the present experiment two or more mature and healthy fruits from the each different jamun plant has been collected and the physical parameters of different fruits have been noted and the biochemical or quality characters of fruits have been studied. Observation on genotypes viz., fruit length (mm), fruit diameter (mm), fruit weight (g), fruit volume (ml), seed length (mm), seed diameter (mm), seed weight (g), pulp content (g), Total soluble sugars ( $^{\circ}$ Brix), Acidity (%) Reducing sugar (%) and total sugar (%) content were recorded from randomly selected fruits from accession. The fruit length was measured with the help of digital vernier caliper from the tip of the fruit to bottom of the fruit and the average fruit length was calculated by the dividing total fruit length by 15 and it expressed in mm (Millimeter). The diameter of the fruit (15) taken from randomly chosen healthy fruits, it was measured by the help of digital vernier caliper and expressed in mm (Millimeter). The fruit volume was measured by water displacement system and average of fruit volume of 15 fruits was observed and expressed in ml (Milliliter). The weight was measured using electronic weighing balance and the calculation of the average fruit weight by dividing total fruit weight by 15 and expressed in gram (g). The seed length and seed diameter was measured by using digital vernier caliper and it expressed in mm (Millimeter). The weight of the seed was measured with the help of electronic weighing balance and results were expressed in gram (g). The pulp content was measured by the deducting average seed weight from the average fruit weight of corresponding fruits and expressed in gram (g). The total soluble solids (TSS) of pulp of the fruits were measured by the Erma Hand Refractometer (0-32  $\square$ B) in terms of degree brix ( $\square$ B) at room temperature. The refractometer was calibrated by the distilled water before the use of refractometer. Bio-chemical assessment of the fruits like the titratable acidity (TA), reducing sugar and total sugar (TS) was done by method suggested by Association of official Analytical chemists. The sources of plant material and the age of tree were different. Thus the studies on variation of such plants did not come under the normal statistical design. The each plant was considered as separate treatment and they were compared for different traits with their mean value of each parameter. Comparison was done on the basis of mean value and average value of parameters studied.

---

## RESULTS AND DISCUSSIONS

---

The analysis of data on physical characters of fruits and seeds of 15 different local jamun genotypes is presented in table 1. The obtained results are presented and discussed as follows. The maximum fruit length has been observed in G-4 (21.89mm). Higher fruit length was also noted in G-13 (20.69mm), G-12 (19.19mm), G-6 (18.46mm). Significantly minimum fruit length was observed in G-2 (4.99mm) followed by the G-8 (5.02mm) and average fruit length was noted as 15.17mm. These present findings are very close to that reported by Swamy *et al.* 2017. The maximum fruit weight has been observed in G-12 & G-14 (3.16g) genotypes. Higher fruit weight was also noted in G-13 (2.38g), G-4 (2.61g), G-5 (2.58g) and G-7 (2.46g). However, minimum fruit weight was observed in G-8 (0.44g) genotype followed by G-2 (0.48g), G-3 (1.06g). Average fruit weight of jamun genotype was noted as 1.89g. The maximum fruit diameter was recorded in G-14 (17.46mm) followed by G-12 (17.33mm), G-1 (16.27mm), G-13 (15.47mm) & so on. While minimum fruit diameter has been observed in G-2 (3.04mm) which was closely followed by G-8 (3.83mm). The average fruit diameter was noted as 12.31mm. Significantly maximum fruit volume has been observed in G-13 (3.28ml) followed by the G-4 (3.20ml), G-12 (3.15ml), G-14 (2.90ml). But minimum fruit volume was noted in G-2 (0.35ml) and also in G-8 (0.41ml) and the average fruit volume was noted as 2.05ml. The maximum seed length was observed in G-4 (16.24mm) followed by the G-7 (15.64mm), G-6 (14.50mm), G-5 (14.11mm), G-13 (13.51mm). While minimum seed length was observed in G-2 (1.99mm) and closely followed by G-8 (2.42mm) & in G-10 (4.20mm). In the present experiment average seed length of 9.70mm was observed. The maximum seed diameter has been found in G-7 (10.46mm) and significantly at par with in G-6 (10.31mm), G-13 (9.34mm), G-4 (9.19mm), G-5 (8.62mm). Significantly minimum seed diameter was observed in G-2 (1.83mm) followed by the G-8 (1.94mm), 3.10mm in G-10 & in G-3 (3.14mm) and the average seed diameter was noted as 6.88mm. The maximum seed weight has been recorded in G-7 (1.12g) which was closely followed by the G-6 (1.03g). While minimum seed weight was observed in G-8 (0.016g) out of 15 genotypes and also 0.026g in G-2, 0.41g in G-15 present. The average seed weight 0.63g was recorded in the present study. The present findings are in consonance with that proposed by Patel *et al.* 2005.

**Table 1: Physical characters of fruits and seeds**

<b>Genotypes</b>	<b>Fruit length(mm)</b>	<b>Fruit diameter(mm)</b>	<b>Fruit weight(g)</b>	<b>Fruit volume(ml)</b>	<b>Seed length(mm)</b>	<b>Seed diameter(mm)</b>	<b>Seed weight(g)</b>
G-1	17.65	16.27	2.24	2.60	9.28	7.88	0.74
G-2	4.99	3.04	0.48	0.35	1.99	1.83	0.026
G-3	12.42	12.29	1.06	1.22	4.69	3.14	0.60
G-4	21.89	15.34	2.61	3.20	16.24	9.19	0.84
G-5	18.45	12.96	2.58	2.60	14.11	8.62	0.80
G-6	18.46	14.38	2.16	2.71	14.50	10.31	1.03
G-7	15.64	10.46	2.46	2.22	15.64	10.46	1.12
G-8	5.02	3.83	0.44	0.41	2.42	1.94	0.016
G-9	14.51	13.50	1.16	1.30	9.11	7.24	0.49
G-10	10.77	8.66	1.46	1.08	4.20	3.10	0.52
G-11	13.39	12.76	1.42	1.05	10.73	8.03	0.48
G-12	19.19	17.33	3.16	3.15	11.68	8.24	0.86
G-13	20.69	15.47	2.80	3.28	13.51	9.34	0.78
G-14	18.43	17.46	3.16	2.90	9.07	7.15	0.82
G-15	16.11	11.03	1.28	2.81	8.34	6.83	0.41
<b>Mean</b>	15.17	12.31	1.89	2.05	9.70	6.88	0.63

Pulp and bio- chemical characters of fruits of 15 different local jamun genotypes are presented in Table 2. Highest pulp content was noted in G-14 (2.34g) which was closely followed by the G-12 (2.30g). Significantly lowest pulp content was found in G-8 (0.42g). Out of 15 genotypes, in 7 genotypes pulp content was found below 1.00g and others above 1.00g. The average pulp content of 1.25g was recorded for 15 genotypes. The present results were partially supplemented by Swamy *et al.* 2017. The observance of the data revealed that G-4 possessed significantly maximum TSS (25.90 □B) followed by the G-13 (25.40 □B), G-6 (24.20 □B), G-5 (23.30 □B). Minimum TSS was noted in G-11 (12.60 □B) out of 15 genotypes and average TSS was noted 18.08 □B. The findings were almost similar with that of Verma *et al.* 2019. Significantly maximum acidity has been observed (1.42%) in G-10 and also in the G-14 (1.17%). Minimum acidity was observed in G-5 and G-7 (0.55%). Out of 15 genotypes only in two genotypes recorded acidity above 1.00% and in others 13 genotypes below 1.00%. The average acidity of pulp of the jamun fruits was recorded 0.82%. Srivastava *et al.*, (2010) were found variation in acidity which is very close to the present findings.

**Table 2: Pulp and bio- chemical characters of fruits**

Genotypes	Pulp content(g)	TSS(□B)	Acidity (%)	Total sugar (%)	Reducing sugar (%)
G-1	1.50	13.4	0.72	12.50	11.56
G-2	0.45	19.2	0.91	16.26	11.11
G-3	0.46	13.6	0.68	13.69	9.85
G-4	1.77	25.9	0.85	13.65	8.23
G-5	1.70	23.3	0.55	16.86	9.85
G-6	1.13	24.2	0.72	15.87	9.38
G-7	1.34	21.3	0.55	14.66	9.70
G-8	0.42	16.3	0.70	15.38	10.36
G-9	0.67	15.0	0.74	18.86	10.01
G-10	0.94	17.2	1.42	15.50	12.82
G-11	0.94	12.6	0.91	16.83	11.12
G-12	2.30	13.8	0.70	13.66	9.52
G-13	2.02	25.4	0.78	14.70	9.85
G-14	2.34	14.4	1.17	16.26	10.36
G-15	0.87	15.6	0.96	13.98	10.52
<b>Mean</b>	1.25	18.08	0.82	15.24	10.28

Highest reducing sugar has been recorded in G-10 (12.82%) and also observed in G-1 (11.56%), G-11 (11.12%) & G-2 (11.11%). While minimum reducing sugar was found in G-4 (8.23%). The average reducing sugar of 15 genotypes was noted 10.28%. Kaur and Bal (2015) reported that variation in reducing sugar which are partially similar to the present findings. Out of 15 genotypes maximum total sugar has been found in G-9 (18.86%) and also followed by the genotypes G-5 (16.86%), G-11 (16.83) & G-2, G-4 (16.26%). Significantly minimum total sugar was noted in G-1 (12.50%) genotype. The average total sugar 15.24% was observed in present study. The present experiment was more close to the Alamet *al.* 2020.

## CONCLUSION

In the present experiment a wide and significant variations in fruit physical characters as well as quality characters was observed among 15 different jamun genotypes. From the findings of the present study it can be concluded that genotype G-14 was superior on the basis of higher fruit weight, fruit diameter and greater pulp content and average total sugar content. Owing to moderate fruit size G-4 genotype is also superior on the basis of higher total soluble solids, total sugar and reducing sugar.

## REFERENCES

- Alam., A., Mani., S., Mitra and Bauri, F.K. 2020. Flowering, Fruiting and Physio-chemical properties of Jamun (*Syzygiumcumini*Skeels) grown in Nadia district of West Bengal. *Advance Bioresearch*.**11**(5): 01-05.
- AOAC, 1990. Official Methods of Analysis. Association of Official Analytical Chemists, Washington, DC.
- Chaudhary, B. and Mukhopadhyay, K. 2012. *Syzygiumcumini* (L.)Skeels: A potential source of nutraceuticals. *Internatonal Journal of Pharm. and Biological Sciences*.**2** (1): 46-53.
- Kaur, M., &Bal, J.S. 2015. An evaluation of jamun (*Syzygiumcumini*Skeels) germplasm for conservation of elite one.I) F, 342.

- Patel, V.B., Pandey, S.N., Singh, S.K. and Das, B. 2005. Variability in jamun (*Syzygiumcumini*Skeels) accessions from Uttar Pradesh and Jharkhand. *Indian Journal of Horticulture*.**62**(3): 244-247.
- Patil, S. S., Thorat, R. M. and Rajasekaran, P. 2012. Utilization of jamun fruit (*Syzygiumcumini*) for production of red wine.*Journal of Advanced Laboratory Research in Biology*.**3**(3): 200-203.
- Singh, I.S. 2001.Minor fruits and their uses.*Indian journal of horticulture*.**58**: 178-182
- Singh, S., Singh, S.P., Singh,V and Shikha, K. 2019. Studies on floral biology, fruit set and fruit drop of different genotypes of jamun (*Syzygiumcumini*Skeels). *The Pharma Innovation Journal*.**8** (1): 558-561.
- Srivastava, V., Rai, P.N. and Kumar, P. 2010. Studies on variability in physic- chemical characters of different accessions of jamun (*Syzygiumcumini*Skeels).*Pantnagar Journal of Research*.**8**(1): 139-142.
- Swamy, G.S.K., Anushma, P.L. and Jagdeesh, R.C. 2017.Morphological characterization of elite Jamun (*Syzygiumcumini*Skeels) genotypes.*International Journal of Minor Fruits, Medicinal and Aromatic Plants*. **3**(1): 09-15.
- Verma, R.S., Lata, R., Ram, R.B., Prakash, S., Kumar, V., Verma, S.S., Pal, H. and Singh, R. 2019. Evaluation of physico-chemical attributes of different genotypes of jamun (*Syzygiumcumini* L. Skeels.) fruits. *Journal of Pharmacognosy and Phytochemistry*.**8**(5): 480-482.
- Zeven, A.C. and De wet, J.M.J. 1982. In: Dictionary of Cultivated Plants and their Regions of diversity: excluding most ornamentals, forest trees and lower plants. Centre for Agricultural Publishing and Documentation (pudoc). Wageningen. Pp. 1-11.