

Current Status and Trend on the Adoption of fish feed additives for Sustainable Tilapia aquaculture production: A Review.

Abstract

During fish feed production, feed additives are regarded as an essential component of the fish diet. The use of feed additives in fish feed is essential for boosting the growth and immunity of cultured fish. The aim of this study was therefore to make a comprehensive review of the Current Status and Trend in the Adoption of these fish feed additives for Sustainable Tilapia aquaculture production. An exploratory research design was used in the study to review and comprehend various peer-reviewed papers published in national and international publications on issues surrounding fish feed additives currently used in Tilapia aquaculture production and the current and future status of essential fish feed additives in use. Feed additives are ingredients that are added in small quantities to fish diets to enhance nutrients and preserve the feed. Preservatives, binders, feeding stimulants, and food colourants constitute common feed additives. With increased growth in aquaculture, the use of additives must be promoted in order to optimize the production of farmed fish. The aim of feed additives in fish feed is to promote healthier and faster fish growth in cultured fish. As a result, fish diets should incorporate the right quantities of feed additives. The conclusions drawn succinctly suggest that fish feed with no feed additives is comparable to "a vehicle without fuel" and should be regarded as poor feed when fed to fish in a culture system. And the end result will be a poor fish growth rate, less resistance to diseases, and hence low productivity in an aquaculture production system.

Keywords: Fish feeds, Feed additives, ingredient, oil, colorant, digestibility, attractants, stimulants

1.0. Introduction

Tilapia fish the world's most diverse intensively farmed fish species have recently hit unprecedented levels of production. In 2030, tilapia output is expected to be approximately 10.8 million metric tonnes based on FAO projection (FAO, 2022). The tilapia sector is now increasing at a 4% annual pace (Oglend, 2020). Disease outbreaks, particularly Streptococcus infection in red Tilapia and slow growth rate pose a significant concern to tilapia production (Hernández et al., 2009). To overcome these increasing challenges, the industry has developed special commercial fish feeds with different feed additives that are both essential and nonessential to the growth rate and disease resistance level of tilapia. Similarly, a variety of novel strains of Red tilapia (Behrends et al., 1982) and **genetically modified farmed Tilapia (GIFT)** (Oliveira et al., 2021) has been introduced. **The new strains of red tilapia have appealing colours, quicker development rates, and high market demand** (Fitzgerald, 1979). The availability of high-quality feeds with all-inclusive feed additives will be critical in realizing the predicted expansion of the tilapia farming industry. Feed additives are compounds that are added to a fish diet or feed ingredients in trace proportions to enhance or preserve it. Preservatives, binders, feeding stimulants, and food colorants are all common feed additives. To meet specific requirements of fish feeds according to (Désiré Adéyèmi et al., 2022) an ingredient or combinations are added to the basic fish feed mix in small levels. In formulated fish feeds attractants, flavors, and digestive aids are added to make the fish meal more appealing, appetizing, and digestible to by the cultured fish. Binding materials are used to keep fish feed from disintegrating in the surrounding culture system (Yossa et al., 2021). Feed free of nutritional components are additives added to diets for purposes besides supplying nutrients (Hossain et al., 2019). These substances, in substantial part, have little or

no nutritive value, but they are essential components of fish feeds, boosting pellet durability, diet safety, diet flavor, and fish health status, and their effect on the end quality of the feed. According to (Tacon&Metian, 2015) Feed binders, carotenoid supplementation, medicines and antibiotics, hormones, antifungals, antioxidants, fiber, flavorings, and water are examples of non-nutritive feed components added to fish feeds. In relation to Adéyèmi et al., (2020) the primary objective of feed additives in fish feeds is to ensure healthier and quicker fish growth, which leads to increased productivity. Therefore, fish feeds should be incorporated with feed additives in the required proportions. There are several types of additives used in the formulation of fish feeds which can be categorized as Essential additives, Non-essential rather than growth-promoting substances additives, and Supplementary additives. The aim of this study was therefore to make a comprehensive review of the Current Status and Trend in the Adoption of these fish feed additives for Sustainable Tilapia aquaculture production.

2.0. Material and Methods

The exploratory research design was used in this article review to evaluate and understand different peer-reviewed papers published in national and international publications journals on matters related to fish feed additives used in global aquaculture production and the current and future status of the crucial fish feed additives on a commercial scale. A comparative critical exploratory approach was used to assess the authors' contributions based on the journals and articles compilations made herein. A total of 98 articles from peer-reviewed journals were used as sources of literature for this study.

3. Results of the findings.

3.0. Types of fish feed additives used in Tilapia feed production

There are three different additives used for the purposes of tilapia fish feed production. These can be categorized as:

1. Essential fish feed additives,
2. Non-essential rather than growth-promoting substances fish feed additives, and
3. Supplementary additives (Auxiliary additives).

3.1. Essential fish feed additives

Essential fish feed additives are added in smaller doses to improve the nutritional value of the fish diet and promote healthy growth. Their extended absence in the fish diet may result in nutritional deficiency diseases, especially in red tilapia (Hernández et al., 2009; Wang et al., 2019). Vitamins, minerals, and so forth are equally vital in fish feed. For instance, fish health management practices used by Kenyan fish farmers include the use of prepared fish feeds with appropriate amounts of fish additives such as calcium and vitamin premix, as well as probiotics in *Oreochromis niloticus* production (Opiyo et al., 2018; Syanya & Mathia 2022) is worth emulation. Notably commonly used essential fish feed additives in aquaculture are not limited to the one discussed below.

3.1.1 Vitamin premix

According to, Rohani et al., (2023) vitamin premixes are concentrations in which continuous forms of required vitamins are blended with basal feed. Vitamin E therapy improves Nile tilapia growth through improved muscle development. Similarly, the reproductive efficiency and egg and larvae quality of Nile tilapia given varying amounts of vitamin C improved significantly. (Sarmiento et al., 2018). Choline chloride is not included in the vitamin premixes since it has been demonstrated to reduce the stability of some vitamins such as water-soluble vitamins. Added amounts ranging from 0.5 to 4% of the diet. Vitamins are vital dietary components that participate in cell metabolism and physiological and biochemical changes in the body of the fish. Vitamins are categorized as water-soluble and fat-soluble. Water-soluble vitamins include vitamin B complex members such as thiamine, riboflavin, pyridoxine, pantothenic acid, niacin, biotin, folic acid, and vitamin B12 (Waagbø, 2010) Similarly water-soluble essential factors alongside the vitamins with a key role in fish growth includes: choline, ascorbic acid; and other vitamins with minimal value to fish nutrition.

However, Thiamine is required for a good digestion system, growth, and fertility in fishes including Tilapia. It is equally required for appropriate neural tissue development. According to Prabhu et al., (2019) deficiency of Thiamine and other mineral elements in Salmon was observed to cause decreased glucose breakdown, low appetite in fish, poor growth, and increased vulnerability to fish stress. Similarly, Kumar et al., (2022) observed that Carps fed thiamine-deficient diets developed skin bulging and subcutaneous hemorrhage.

Vitamins A, D, E, and K are examples of fat-soluble vitamins also referred to as phyloquinone that act as fish feed additives. Vitamin E is the most common of fat-soluble vitamins due to its key role as an antioxidant in fish. Vitamin E helps to maintain normal blood capillary permeability and heart muscle integrity of the fish (Baker & Davies, 1997).

According to (Rohani et al., 2023) Vitamin E treatment was reported to improve Nile tilapia growth through enhanced muscle development. However, so far, no vitamin D requirements for fish have been reported, though some study has been done in this field utilizing highly purified diets and controlled studies with young developing fish (Brown & Robinson, 1992; Fraser, 2018; Lock et al., 2010). Therefore, vitamin premix in fish feed as the additive is not an exception for sustainable growth and biochemical functioning of culture fish.

3.1.2. Inorganic Minerals.

Major minerals that are considered with a wide range of use in the aquaculture sector include Calcium, phosphorus, sodium, molybdenum, chlorine, magnesium, iron, selenium, iodine, manganese, copper, cobalt, and zinc are all considered required for fish physiological functioning (Antony Jesu Prabhu et al., 2016). These mineral elements can be supplemented with fluorine and chromium, for the case of tilapia fish species both of which have been found to be essential elements when incorporated with feed as additives for their enhanced growth. According to Kord et al., (2022) In the diet of Tilapia Fry and fingerlings Calcium, phosphorus, sodium, potassium, iron, manganese, magnesium, copper, chloride, iodine, cobalt, and zinc are considered necessary minerals to be incorporated in the feeds. Therefore, their inefficiency may result in nutritional deficiency disease outbreaks in the culture system. When trace minerals like copper, zinc, cobalt, iodine, and manganese are supplied in the fish diet as additives, the growth of the fish is generally enhanced (Sørensen, 2012). Similarly, Calcium and phosphorus are required for bone and exoskeleton development in most fishes and crustaceans, while sodium, potassium, magnesium, and chloride are involved in osmoregulation (Antony Jesu Prabhu et al., 2016). Biochemical enzymes contain magnesium, manganese, and zinc as cofactors (Yu et al., 2021a). There is a need to incorporate most of the essential mineral elements as feed additives when formulating fish feed for aquaculture production as reported by the findings of most of these authors.

3.1.3 Fish oils

Fish oil (FO) has traditionally been the primary lipid source that has been used as fish feed additives due to its high concentration of n-3 long-chain polyunsaturated fatty acids (n-3 LC-PUFA) such as docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), which are essential for fish growth and immune booster during production (Peng et al., 2008). Fish oil when incorporated in the fish feeds provides dietary energy and the required essential fatty acids for the cultured fish, especially Tilapia species. According to Lin & Shiau, (2007), there is a significant effect on growth and non-specific immunological responses on the use of a fish oil-corn oil dietary combination on grouper, (*Epinephelus malabaricus*). Fish oil is therefore added at 2- 3% as additives in Tilapia fish feeds during feed formulation in order to improve growth and food conversion ratio as well as the palatability of the feeds by the fish. Similarly, according to Karapanagiotidis et al., (2022) fish oils high in polyunsaturated fatty acids (PUFA) such as cod liver oil, sardine oil, squid oil, and clam oil are commonly used as feed additives in the aquaculture sector. The feeds containing fish oil as additives are considered highly palatable by fish and promote FCR and digestibility of the feeds by the cultured fish.

3.1.4. Phospholipids as fish feed additives.

For fish growth, phospholipids provide choline, phosphorus, and necessary fatty acids that need to be provided through dietary feeds. Dietary phospholipids improve growth performance indirectly by

increasing CCK levels, which stimulate pancreatic secretion by increasing chylomicron levels (Luo et al., 2010). Similarly, Phospholipids, such as phosphatidylcholine, are necessary for fish growth and long-term survival. Khorshidi et al., (2022) reported that common carp (*Cyprinus carpio*) diet containing 1-2% soybean lecithin boosts accelerated growth and enhances the feed conversion ratio. This too is practically possible for tilapia fish species. Further phospholipids are physiologically significant in the transfer of lipids in the body (Wu et al., 2022). Many plant and animal sources have phospholipids, although not all of them have large quantities of certain phospholipids. Vegetable phospholipids, do not have sufficient long-chain polyunsaturated fatty acids (LC-PUFAs) such as EPA and DHA that are vital, especially in Tilapia fish farming. Therefore, due to this deficiency, Soya phospholipids are the most commonly used phospholipid source in red Tilapia fish feed additives during fish feed formulation. This is due to their low (phosphatidylcholine; a kind of phospholipid) concentration and omega-3 fatty acid levels. Therefore, for optimal growth and survival as well as prevention of fish skeletal deformities phospholipids are a requirement in the fish diet. Lack of sufficient dietary phospholipids limits lipoprotein synthesis in enterocytes, leading to impaired transport of lipids to the tissues in carps (Khorshidi et al., 2022). Similarly, according to (Luo et al., 2010) Japanese sea bass (*Lateolabrax japonicus* L.) reared in freshwater dietary phospholipids increase cholesterol and triglyceride mobilization from the gut to the hepatopancreas, hemolymph, and muscle. Phospholipids therefore in the fish meal when used as an additive are able to influence lipid deposition, resulting in enhanced lipid accumulation in the fish. Based on these findings Phospholipids stand out as the most essential additive when formulating Tilapia fish feeds in order to record outstanding growth during the culture period.

3.1.5. Fatty Acids:

Lipids (fats) are high-energy nutrients that should make up about 15% of fish diets. Their primary role in the diet is to provide essential fatty acids (EFA) and to transport fat-soluble vitamins. The most important required lipids for fish feeding are omega 3 and omega 6 fatty acids, often known as n-3 and n-6. (Hoseinifar et al., 2017) Similarly (Hearn et al., 1987) reported that to improve growth, 1% of highly unsaturated fatty acids (HUFA) such as eicosatetraenoic acid (20:5 W3) and docosahexaenoic acid (22:6 W3) can be added to fish feed. Freshwater fish such as Tilapia do not require long-chain highly unsaturated fatty acids, but they do require linolenic acid (18:3-n-3) in amounts ranging from 0.5 to 1.5% of the feed formulated (Hoseinifar et al., 2017) This is because Freshwater fishes are unable to synthesize this fatty acid, thus it must be provided as additives within the fish feed during the feeding process. Several freshwater fish however may ingest this fatty acid and use enzyme systems to extend to a long-chain polyunsaturated fatty acid.

3.1.6. Cholesterol:

Cholesterol is a nutritionally important steroid in the diet of fish. The addition of 0.1- 0.5% cholesterol to a fishmeal boosts growth and survival (Deng et al., 2014). The prawn head waste meal contains cholesterol that should be incorporated to fish feeds during formulation. And because bony fish can synthesize cholesterol, it is not necessary to supplement dietary cholesterol for their health (Deng et al., 2013), however recent studies that have suggested that adding cholesterol and soybean meal-based additives to rainbow trout (*Oncorhynchus mykiss*) formulated feeds as a protein source will increase their growth performance. Similarly, (Deng et al., 2014) observed that the Nonspecific immunity of rainbow trout (*Oncorhynchus mykiss*) can be improved by consuming 0.6-1.2% cholesterol. Nevertheless, our understanding of alternative physiological functions that cholesterol can play to enhance fish health under stress circumstances is limited, despite the finding made by (Xu et al., 2018) that cholesterol is tightly connected to tilapia fish stress via the hypothalamic-pituitary-interregal (HPI) axis when reared under brackish waters with high salinity. In relation to this finding Tilapia fish feed should incorporate cholesterol as a feed additive to boost the immunity and stress endurance of the fish in the culture systems even though freshwater fishes do not necessarily require cholesterol.

3.2. Non-essential but growth-promoting substances fish feed additives

Non-essential growth promoter substances are feed Ingredients that are sourced from plants and animals, single-cell proteins, and certain synthetic chemicals which can be utilized as feed supplements to promote faster fish growth and higher yield. Non-essential growth promoters do not cause deficiency illness if they are not included in the fish diet meal (Opiyo et al., 2019; Xu et al., 2021). However, when supplemented with fish feed as additives they have a more recommended significant role in fish growth. They are attractants and growth boosters either from Plant or animal materials, as well as single-cell proteins, antibiotics, drugs, and so on are examples worth mentioning.

3.2.1 Antibiotics:

In aquaculture, Antibiotics are administered to combat infections caused by bacteria in either shrimp or fish farms. Antibiotics may be utilized to cure bacterial-related infections and diseases, but they are usually used to prevent infections by treating the water or fish before the disease emerges (Romero et al., 2012). Antibiotics, according to Zhou et al., (2021), are often observed to increase growth in juvenile fish rather than adult fishes, especially when incorporated in fish feeds as an additive. Therefore, Antibiotics perform well in fish diets prepared with vegetable proteins. Antibiotics result in the Decrease or elimination of pathogen activity from the fish culture system and Remove microorganisms that create poisons that inhibit growth. (Romero et al., 2012). The use of antibiotics sometimes encourages the development of helpful bacteria such as probiotics that synthesize nutrients for the fish, decrease bacteria that compete for nutrition with the host, and Improve the intestine's absorption capacity. Based on these findings we strongly recommend the use of antibiotics as an additive in fish feeds however in smaller concentrations since the Massive use of antibiotics in aquaculture production has been outlawed by most EU countries.

3.2.2. Hormones:

Hormones are utilized in aquaculture for artificial reproduction in hatcheries and sex reversal. Artificial reproduction keeps the production units such as hatcheries constantly producing fingerlings (Yamazaki, 1976). Hormones are employed in sex reversal when the male and female growth rates differ significantly and are able to gain different weights and sizes at maturity. Several natural and synthetic hormones have also been utilized in aquaculture for instance spawning induction, sex reversal, mono-sex population production, and growth promotion. (Higgs et al., 1982; Shore & Shemesh, 2003) Growth hormones, such as thyroid hormone, gonadotropin, prolactin, insulin, and different steroids are the hormones responsible for fish growth (Higgs et al., 1982). Growth hormones such as androgen, estrogens, and progestogens, as well as non-steroidal hormones such as thyroxin, are utilized as growth promoters (Yamazaki, 1976).

3.2.3. Drugs:

Arsenicals and sulpha medicines when cooperated with fish feeds promote growth in Tilapia species specifically. The arsenicals that can be used as additives during feed formulation include 3-nitro-4-hydroxy phenyl arsenic acid, para-amino-phenyl arsenic acid, and sodium salt (Bijoy et al., 2018; Opiyo et al., 2018). Sulphonamides are among the sulpha medicines used. Medicines work as an immune promoter and for growth in juvenile fish, improving their overall health and biochemical activities. However, according to most literature on drug usage in aquaculture, the precise method of action of these medicines in fish is unknown, their actions appear to be comparable to those of antibiotics (Zhou et al., 2021). Therefore, it's prudent enough to incorporate Veterinary medicines and drugs as additives during fish feed formulation due to the associated benefits to the fish cultured.

3.2.4. Enzymes in fish diet

Protease, amylase, lipase, esterase, cellulase, xylanase, and urease are commonly used enzymes in the aquaculture sector to promote the growth of the culture fish species (Khorshidi et al., 2022). These enzymes are administered to fish as an additive since they can boost nutritional intake, improve the uptake of nutrients in the gastrointestinal tract, and accelerate fish maturation. According to Wangkahart et al., (2022) enzymes used as additives in fish feeds improve feed digestibility in fish hence being digested effectively. An enzyme in the fish body is also utilized to digest complex carbohydrates, collagen in the skin and bones, and other feed ingredients by the fish. Enzymes should be handled within optimum temperatures of between 40 °C to 50 °C. (Wiszniewski et al., 2022)

Therefore Temperatures exceeding 65^oc denature enzymes in the feed and hence become of no value to the fish. As a result, enzyme supplements are frequently applied to meals after pelleting.

3.2.5. Probiotics and Prebiotics

Probiotics are live microorganisms' dietary additives that promote fish development by influencing the intestinal microbiota flora density. Probiotics can be a single species or a combination of microorganisms of various species. The supplements contain bacteria that colonize the intestines and kill harmful microbes. Enabling the fish to avoid wasting metabolic energy fighting against harmful pathogens. After pelleting, probiotics must be introduced to the diets(Cano-Lozano et al., 2022; Guimarães et al., 2019). Similarly, Prebiotics are nondigestible fibers that benefit the host's health by specifically encouraging the development and activity of certain genera of microorganisms in the colon, primarily lactobacilli and bacteria in Nile Tilapia post-larvae diet (de Araújo et al., 2018)

3.2.6. Synbiotics and phytobiotics

Synbiotics were first proposed by Gibson where he incorporated, prebiotics with probiotics to form Symbiotic compounds (Mohammadi et al., 2022). The survivability and adoption of live microbiological dietary supplements within the gastrointestinal system were promoted by selectively encouraging an expansion of symbiotic benefits to the fish. The photobiotic vial is defined as a plant-derived nutrient infusion to feed in arrangements to promote aquatic species' performance. Herbs, spices, and other plants' leaves, roots, tubers, and fruits are all utilized as pytobiotics. Pytobiotics are commonly utilized to improve growth and yield in fish and prawn cultures (Saeed et al., 2014). All these feed additives ought to be included in the Tilapia fish diet however in small proportions due to environmental concerns about the Phyto therapeutic nature of photobiotic.

3.3. Supplementary additives (Auxiliary additives).

Auxiliary fish feed additives are small ingredients applied to feeds during formulation (Owatari et al., 2022). They serve as ingredients to enhance the physical appearance of fish feed. This, in turn, aids in increased digestibility of the feeds by the fish and hence culminating in increased feed efficiency. These additives can also be considered supplementary additives. They include and are not limited to Feed coloration, binders, molasses, fats, attractants, and so on.

A range of substances is added to fish diets to protect water-soluble nutrients from dissolving into the culture systems before the feeds are consumed by the fish and can also promote nutrient effective utilization by cultured fish. Binders, antioxidants, mould inhibitors, antibacterial agents, attractants, growth promoters, mendicants, colors, and other ingredients are frequently used as supplementary additives in GIFT tilapia fish farming (Xu et al., 2021). The active ingredient selected and used should not be hazardous to the cultured fish species or the farmer and end fish consumers. Similarly, Van Doan et al., (2021)In their finding, while supplementing pineapple peel powder and Lactobacillus plantarum as disease-resistance substances in Nile tilapia (*Oreochromis niloticus*) feeds. The author reported that the supplementary additives used should not react with the feed ingredients and negatively affect the nutritional value of the feed. They further concluded that these additives should not reduce the desirable qualities of the fish feeds produced by affecting their taste, appearance, flavour, and texture. This shows that the best Auxiliary additives should have minimal impact on the quality and nutritional values of the fish feeds as well as the source of material used should be readily available to an ordinary farmer in adequate quantities and at an affordable price.

3.3.1 Fish Feed Colourants and Pigments.

Fish feeds colorants are commonly considered to be of Plant and animal origin. Currently, there are roughly 300 pigments found in plants and animals used in both finfish and shellfish aquaculture as pigment supplementary additives. Tuan Harith et al., (2022a)Reported that Carotenoids and astaxanthins can be found in both natural and synthetic sources and that the most significant pigment classes are xanthophyll and carotenoids. Mostly xanthophyll is found in plants, while carotenoid pigments are present in crustaceans and fish (Tuan Harith et al., 2022b). Additionally, these pigments have been equally observed to increase fish growth and survival. The main role of these colorations is to ensure the correct pigmentation of cultivated organisms. The colour of red tilapia fish (Basavaraja & Raghavendra, 2017; Wohlfarth et al., 1990) has a significant impact on market pricing and demand since the consumers have a perception of Red tilapia being similar to most colored marine species in

the market due to its red coloration. Carotenoids are the most significant of the substances that give color to the Red tilapia Fish as well as crustaceans that cannot produce colors, but they may change the compounds by oxidation. The effect of dietary astaxanthin enrichment on the colour and development of red tilapia, *Oreochromis spp* is well documented as having a significant effect on growth as well (Tuan Harith et al., 2022a). The addition of carotenoids to broodstock diets was observed to cause a decrease in maturation time, an increase in egg quantity, improved egg hatchability, and larval survival (Wan-Mohtar et al., 2021). The right dosage of Fish Feed colorants is of wide economic benefit to the aquaculture sector and therefore its usage needs to be encouraged during the fish feed formulation stage since fish farmers in the rural setting have no idea of the proper usage and supplement required dosage for these fish feed additives even though they could be aware of the economic and nutritional benefits values of the additives to the cultured fish.

3.3.2. Binders:

Fish feed must be robust enough just to sustain typical handling and transportation without degradation. Moreover, fish feed must be water-stable. For this reason, Binders are added to fish feeds during formulation to increase their stability in water without being disintegrated before the feed is consumed by the fish. Most plant Starches are used as basic feed components during preparation and formulation which act as a binder in feed. Binder ingredients include agar-agar, carboxymethylcellulose (CMC), bentonite, guar gum, lignin sulfate, plaster of Paris, polyvinyl alcohol, sodium alginate, and wheat gluten (Désiré Adéyèmi et al., 2022).

Binders are required to provide the desired water stability to the feeds, preventing the disintegration of the fish feeds into the culture water before the fish consumes the feeds. The binding strength and cost influence binder selection and inclusion levels (Hossain et al., 2019). The commonly used binders in aquaculture are Gelatinised starch from tapioca, wheat flour, and rice flour which are considered readily available and cost-effective to the common fish farmer and hence sustainable. Recommended levels of binder's guar gum, gum-acacia, are 1-2%; gelatin, collagen, carrageenan, and agar 2-5%; and wheat gluten (10-12%) and wheat flour and tapioca starch could be used in relatively high levels (Flefil et al., 2022). A mixture of some of these binders may be considered more effective and economical to use. Similarly, to some extent, chitin and chitosan can be used in larval feed meals during larval rearing.

3.3.3. Molasses

The addition of molasses to fish feed as an additive often aids in the smooth pelleting process of fish feed. The palatability of feeds is also improved due to the presence of molasses. Similarly, it's a good source of energy for fish in feed (Khanjani et al., 2021). However, Molasses (with a C/N ratio of 1:2) was observed to raise the pH of pond water, promoting the production of beneficial biofloc in the pond (Alfiansah et al., 2022). Molasses also enhanced carbohydrates and proteins in bio-flocs and kept beneficial bacteria in abundance, resulting in low inorganic nutrient availability to the fish. As a result, molasses is appropriate for Tilapia and shrimp farming to boost growth rate and production.

3.3.4. Chemo-attractants and feeding stimulants

These are chemicals that stimulate eating patterns in animals and enhance feed consumption. The most crucial are free amino acids and nucleotides. Several finfish species are known to require special feeding stimulants in general, L-amino acid, glycine-betaine, inosine, or inosine-5-phosphate mixtures are regarded as 'universal feeding stimulants' for fish (Straus et al., 2013).

4.0. Current Emerging trend on the use of additives in fish feed in aquaculture.

Fish feed additives have increasingly been used in different aquaculture fish production systems depending on whether finfish/shellfish species are under culture. Different authors have reviewed these issues as well as conducted research on the current usage scale of additives in fish feed production, and their effect on the quality of aquaculture products in the market. Based on the study done by Lin et al., (2022) the novel elements such as benzyl paraben incorporated in fish feeds as additives promoted growth and acted as antioxidant and lipid metabolism of Nile tilapia (*Oreochromis niloticus*). This is an indication that some vital compounds can be incorporated in fish feeds during

formulations as feed additives on alleviating toxicity in farmed fish. Similarly, fin fishes are considered valuable and less expensive sources of omega-3 fatty acids and vital mineral elements in the human body (Shenouda, 1980). There is an urgent need to conduct additional research on novel fish feed additives, such as the inclusion of herbs in fish feeds, which reduce feed expenses, increase nutrient utilization, and prevent the remaining effects of hormones and antibiotics on fish tissues, which in turn have impacts on people who consume them. According to Yu et al., (2021b) Neuropeptide Y used as a feed additive has a significant effect of stimulating and promoting the growth of Tilapia (*Oreochromis niloticus*), especially when fed on a low fish meal diet. This is an indication that the presence of feed additives in fish meals promotes tilapia growth, improves their resistance to disease, and provides therapeutic benefits as compared to standard diets.

Probiotics, prebiotics, phytogenic ingredients, immunological stimulants, enzymes, hormones, mycotoxin binders, organic acids, and other functional feed additives are the optimum feed additives currently used in aquaculture. (Cano-Lozano et al., 2022; de Araújo et al., 2018). The authors observed that these additives have an influence on the growth of Tilapia fish species and recommended these feed additives be used in aquaculture for fish-formulated diets. In developing countries such as not limited to Kenya Tanzania Thailand India, the aquaculture industry is rapidly expanding. (Basavaraja, 2015; Hishamunda & Ridler, 2006) at an equal rate, fish feed demand is on the high rise, and to improve the quality of fish feed produced locally for aquaculture use feed additives are mandatory to be in cooperation. The purpose of using feed additives in aquaculture is to improve fish feed palatability, efficiency, and profitability. However, due to the biodiversity loss of most organisms that can be used as vital additives in fish feeds during formulation, environmental degradation, and the release of feed additive residues into the environment, a moral judgment option on feed additives has become compulsory by law. Further studies focused on boosting fish production through additives will allow the right fish feed. According to Gabriel, (2019); Poolsawat et al., (2021) certain studies on enzymes, herbal feed additives, and probiotics utilized as additives in fish feed have been reviewed. These feed additives enhance feed flavor and texture, feed color, and feed nutritional quality, as well as feed digestibility and, eventually, production efficiency.

Researchers such as Dang et al., (2021); Mamdouh et al., (2021) investigated whether feed additives in Tilapia fish diets can act as chemoprevention, mitigating the toxicity risk of various contaminants and detoxifying some active toxic substances in the fish feed. And therefore, having a clear overview of the internal action pathways of various protective feed additives may provide novel alternative treatments for the feeds during storage. Similarly studying fish nutritional requirements sought to be done with a lot of caution and hence there is a need to look for revolutionary feed additives and supplements that ensure cheap feed costs, and maximum digestibility, but few detrimental consequences on fish and the environment (Désiré Adéyèmi et al., 2022; Syed et al., 2022). Similarly, Genschick et al., (2021) in their work assessed how high-quality feed is prepared from acceptable and readily available feed additives. Beneficial feed additives improve not only the growth and yield of the fish but also the health function of the fish. Prebiotics, probiotics, seaweeds, microalgae, enzymes, organic acids, mycotoxin binders, phytobiotic substances, and yeasts are among the beneficial feed additives. Similarly, Zhou et al., (2021) evaluated the use of antibiotics in Tilapia fish farming feeds to control viral illnesses and increase growth performance, which is a common practice. Recently, the precautionary use of antibiotics and chemotherapy treatments was condemned, leading to their general ban in the global fish markets. The use of beneficial feed additives as an alternative to antibiotics in aquaculture diets is being encouraged among most fish farmers (Romero et al., 2012). The current study provides a comprehensive and practical consolidation of previous studies on various feed additives used in aquaculture, especially for Tilapia fish production with examples including probiotics, prebiotics, synbiotics, immunostimulants, organic acids, nucleotides, and medicinal plants.

In addition, according to El-Kady et al., (2022) the use of probiotics, prebiotics, and synbiotics for the sustainable growth of tilapia fish species has been evaluated. The authors highlight and discuss the effects of probiotic, prebiotic, or synbiotic administration on growth performance, stress tolerance, intestinal microbiota, immune response, and health of Nile Tilapia (*Oreochromis niloticus*). Harikrishnan et al., (2020) also analyzed how a Dried lemon peel-enriched diet improves

the antioxidant activity, growth, and metabolism of *Labeorohita*. Dried lemon peels can therefore be used to modulate the immune antioxidant of Tilapia as well. A similar finding was reported by (Syed et al., 2022) that Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) fed on different levels of Aloe vera extract as feed additives in a closed aquaculture system enhanced growth. Gabriel et al., (2017) observed that a fish diet containing Aloe vera (*Liliaceae*) as feed additives genetically enhances sex reversal in Farmed Nile Tilapia Fingerlings.

The use of Aloe vera in fish feeds has no adverse effect on the health of the fish, farmer and it's considered environmentally sustainable. This study, therefore, recommends the use of Aloe vera in sex reversal and growth promoters in the production of all male Tilapia fish species in order to regulate the prolific breeding nature of the fish for sustainable aquaculture production. Plant compounds prove to be environmentally and user-friendly when used in aquaculture feed production as a basic feed additive. Furthermore, plant extract additives are readily available and cheap to a common fish farmer this as a result will lower the cost of feed production and hence lowers the cost of feeds as well. Based on the study's findings, Syed et al., (2022) established that Aloe Vera fortification at 400g/kg diet is essential to enhancing the growth performance and metabolism of Nile Tilapia (*Oreochromis niloticus*). Another study by Ojha et al., (2014) examined how the fish feed in cooperation with *Petalium murex* as an additive play a role in improving the survival, metabolism, and immunity of *Labeorohita* (Hamilton, 1822) fingerlings. Their finding shows that *Petalium murex* extracts feed additives have a significant effect on promoting the growth performance of *Labeorohita*, fingerlings in addition to improving the fish's metabolism and disease resistance. The authors, therefore, recommended a dose of (0.08 gm/100 gm diet) in the diet *Labeorohita* feed during formulation. This Review study, therefore, makes a strong recommendation for the use of *Petalium murex* as growth promoter feed additives for Tilapia fish farming and may indeed be recommended for commercial aquaculture production globally.

The impact of an ethanolic extract of *Mucuna pruriens* on the growth, biochemistry, and sex reversal of Nile Tilapia (*Oreochromis niloticus*), and the spermatogenic influence of ethanolic extract *mucuna pruriens* was evaluated by the authors (Etta et al., 2009; Mukherjee et al., 2015). Their findings showed a significant effect on the growth, metabolism, and hematological parameters of Nile Tilapia (*Oreochromis niloticus*) fingerlings. The male population was also enhanced by *Mucuna* additives in the diet. This is a clear indication that *Mucuna pruriens* is the best-fit feed additive to be used in the commercial production of aquaculture. Furthermore, it has the potential to boost the fish's digestion and resistance. Furthermore, Nuhu Ozovehe (2013) reported in his study on the effects of different amounts of Moringa oleifera leaf meal diet on African catfish *Clarias gariepinus* growth performance, hematological indices, and biochemical enzymes that the growth performance of juvenile *Clarias gariepinus* significantly improved when fed feeds containing Moringa oleifera leaf additives. **Based on these findings, the effect of Moringa oleifera on growth performance, hematological effects, and biochemical activities in the bodies of various tilapia species such as red tilapia, Nile tilapia, tilapia Zilli, Oreochromis Mozambique, and other tilapia strains should be explored and documented. A similar finding was made by Saha et al., (2011) in their study on the Evaluation of the nutritive Value of Water Hyacinth (*Eichhornia crassipes*) Leaf Meal in compound diets for Rohu, Labeo fingerling diets following fermentation with two microbial isolates derived from the gut microbiome. Their findings geared towards the recommendation that 40% of fish meals can be substituted with water hyacinth leaf meal fermented with fish gut bacteria, as a feed additive with no adverse effect on different fish species. This finding when successfully implemented will make cost-effective formulated fish feed with high nutritional additives. The health and nutrition properties of Mucuna seed meal and its intake in the diet of *Clarias gariepinus* as feed additives are quite promising. Ayorinde et al., (1985); Okomoda et al., (2017) both concluded that toasting Mucuna seed meals significantly improved the nutritional quality of the fish feeds and palatability hence, allowing better consumption of the feed by the African catfish *C. gariepinus* at an inclusion level of 200g/Kg. In relation to these studies, this review paper recommends the adoption of Mucuna seed meal toasted to Tilapia fish farming to enhance the nutritional value of the fish feeds.**

Dietary cooperation of carrots (*Daucus carota*) used as colorant pigmentation in swordtail (*Xiphophorus helleri*) significantly enhanced the orange colouring of the body, especially when supplemented with dietary spinach at 1.3g/100g of the fish diet (Mukhopadhyay, 2021)

Based on the findings of this author this study recommends that since artificial carotenoid is costly, natural carotenoid sources like carrots and spinach can be incorporated into the fish diet as pigmentation coloration additives to improve the body color of Red Tilapia and O. Mozambique as well. The impact of the substitution of sunflower oil for fish oil in the diet of juvenile *Catla catla* (Ham) on growth performance and feed utilization has also been studied (Mukhopadhyay & Rout, 1996; SP et al., 2015). The results of the author clearly show that sunflower oil may be largely 50% substituted with no harmful effect on the growth performance of Indian Carp (*catla catla*). This study further recommends a similar study to be done on different species of Tilapia to optimize the right formulation dosage for sunflower oil as additives in Tilapia fish feed production. Similarly, Liu et al., (2019) evaluated the significance of oils on the nutritional value of fillets in Nile tilapia and ascertained that oils are inexpensive providers of essential fatty acids. The utilization of food oils in precise amounts as required by different fish species and not limited to fin fishes only but even crustaceans for optimal growth and muscle and body mass index development. These oils in fish diets are advantageous to fish due to their antibacterial, anti-oxidative, and growth-promoting properties, as well as their potential to improve feed palatability, digestibility, and binding capacity. This study strongly recommends the use of oils and fats as essential additives during fish feed formulation for sustainable aquaculture production.

5.0. Conclusion.

In conclusion, Essential fish feed additives, growth-boosting but non-essential fish feed additives, and auxiliary additives currently are the most common fish feed additives used in aquaculture production. The reviewed literature spells out clearly their importance in fish health ranching from promoting growth to sex reversal immune boosters in most culture fish species inclusive of Tilapia. Preservatives, binders, feeding stimulants, and food colorants are major feed additives that have been reported to promote healthy and quicker fish growth, resulting in increased production and yield. Therefore, it's essential that all locally formulated fish feeds should adhere to the inclusion levels of both essential, non-essential, and auxiliary fish feed additives in the fish diet. Lastly, feed additives are added to fish feed to improve growth and disease resistance levels and make fish feeds more appealing, palatable, and digestible- attractants, tastes, and increased digestibility. This summarily indicates that fish feed with No feed additives is like "a vehicle without fuel" and in the ordinary scenario is considered as NO feed when given to fish in a culture system. If additives are not included then the fish should better be fed on natural feeds such as copepods, artemia, phytoplankton, zooplankton, and algae. That explains how vital fish feed additives are to cultured fish for sustainable aquaculture production.

REFERENCES

- Adéyèmi, A. D., Kayodé, A. P. P., Chabi, I. B., Odouaro, O. B. O., Nout, M. J. R., & Linnemann, A. R. (2020). Screening local feed ingredients of Benin, West Africa, for fish feed formulation. July 2020, 100386 *Aquaculture Reports*, 17 July 2020 <https://doi.org/10.1016/j.aqrep.2020.100386>
- Alfiansah, Y. R., Harder, J., Slater, M. J., & Gärdes, A. (2022). Addition of Molasses Ameliorates Water and Bio-Floc Quality in Shrimp Pond Water. *Tropical Life Sciences Research*, 33(1), 121. <https://doi.org/10.21315/TLSR2022.33.1.8>

- Antony Jesu Prabhu, P., Schrama, J. W., & Kaushik, S. J. (2016). Mineral requirements of fish: A systematic review. *Reviews in Aquaculture*, 8(2), 172–219. <https://doi.org/10.1111/RAQ.12090>
- Ayorinde, F. O., Grissom, F. E., Afolabi, O. A., Oshuntogun, B. A., Adewusi, S. R., Fapojuwo, O. O., & Oke, O. L. (1985). Preliminary Nutritional and Chemical Evaluation of Raw Seeds from *Mucuna solanei*: An Underutilized Food Source. *Journal of Agricultural and Food Chemistry*, 33(1), 122–124. <https://doi.org/10.1021/JF00061A035>
- Baker, R. T. M., & Davies, S. J. (1997). Modulation of tissue cc-tocopherol in African catfish, *Clarias gariepinus* (Burchell), fed oxidized oils, and the compensatory effect of supplemental dietary vitamin E. *Aquaculture Nutrition*, 3(2), 91–97. <https://doi.org/10.1046/J.1365-2095.1997.00078.X>
- Basavaraja, N. (2015). Recent developments in Indian aquaculture. *Fish Chim*, 34(12), 24–28.
- Basavaraja, N., & Raghavendra, C. H. (2017). Hormonal sex reversal in red tilapia (*Oreochromis niloticus* and *Oreochromis mossambicus*) and inheritance of body colour in *O. mossambicus* and red tilapia: implications for commercial farming. *Aquaculture International*, 25(3), 1317–1331. <https://doi.org/10.1007/S10499-017-0116-3/FIGURES/1>
- Behrends, L. L., Nelson, R. G., Smitherman, R. O., & Stone, N. M. (1982). BREEDING AND CULTURE OF THE RED □ GOLD COLOR PHASE OF TILAPIA. *Journal of the World Mariculture Society*, 13(1–4), 210–220. <https://doi.org/10.1111/J.1749-7345.1982.TB00028.X>
- Bijoy, V. M., Sabu, S., & Harikrishnan, M. (2018). Fish meal replacement with squilla (*Oratosquilla napa*, Latreille) silage in a practical diet for the juvenile giant freshwater prawn, *Macrobrachium rosenbergii* de man, 1879. *Aquaculture International*, 26(5), 1229–1245. <https://doi.org/10.1007/S10499-018-0280-0/TABLES/10>
- Brown, P. B., & Robinson, E. H. (1992). Vitamin D studies with channel catfish (*Ictalurus punctatus*) reared in calcium-free water. *Comparative Biochemistry and Physiology -- Part A: Physiology*, 103(1), 213–219. [https://doi.org/10.1016/0300-9629\(92\)90265-R](https://doi.org/10.1016/0300-9629(92)90265-R)
- Cano-Lozano, J. A., Villamil Diaz, L. M., Melo Bolivar, J. F., Hume, M. E., & Ruiz Pardo, R. Y. (2022). Probiotics in tilapia (*Oreochromis niloticus*) culture: Potential probiotic *Lactococcus lactis* culture conditions. *Journal of Bioscience and Bioengineering*, 133(3), 187–194. <https://doi.org/10.1016/j.jbiosc.2021.11.004>
- Dang, Z. C., Arena, M., & Kienzler, A. (2021). Fish toxicity testing for the identification of thyroid-disrupting chemicals. *Environmental Pollution*, 284. <https://doi.org/10.1016/j.envpol.2021.117374>
- de Araújo, E. R. L., Barbas, L. A. L., Ishikawa, C. M., de Carla Dias, D., Sussel, F. R., de Almeida Marques, H. L., & Tachibana, L. (2018). Prebiotic, probiotic, and synbiotic in the diet of Nile tilapia post-larvae during the sex reversal phase. *Aquaculture International*, 26(1). <https://doi.org/10.1007/s10499-017-0201-7>
- Deng, J., Bi, B., Kang, B., Kong, L., Wang, Q., & Zhang, X. (2013). Improving the growth performance and cholesterol metabolism of rainbow trout (*Oncorhynchus mykiss*) fed soyabean meal-based diets using dietary cholesterol supplementation. *British Journal of Nutrition*, 110(1), 29–39. <https://doi.org/10.1017/S0007114512004680>
- Deng, J., Zhang, X., Long, X., Tao, L., Wang, Z., Niu, G., & Kang, B. (2014). Effects of dietary cholesterol supplementation on growth and cholesterol metabolism of rainbow trout (*Oncorhynchus mykiss*) fed diets with cottonseed meal or rapeseed meal. *Fish Physiology and Biochemistry*, 40(6), 1827–1838. <https://doi.org/10.1007/S10695-014-9971-2>

- Désiré Adéyèmi, A., Polycarpe Kayodé, A. P., Morelle Atcheon, I., Bienvenue Chabi, I., Rob Nout, M. J., & Linnemann, A. R. (2022). Performance of sustainable fish feeds in Benin for *Clarias gariepinus* (Burchell 1822). *Egyptian Journal of Aquatic Research*. <https://doi.org/10.1016/j.ejar.2022.09.002>
- El-Kady, A. A., Magouz, F. I., Mahmoud, S. A., & Abdel-Rahim, M. M. (2022). The effects of some commercial probiotics as water additive on water quality, fish performance, blood biochemical parameters, expression of growth and immune-related genes, and histology of Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 546. <https://doi.org/10.1016/j.aquaculture.2021.737249>
- Etta, H. E., Basse, U. P., Eneobong, E. E., & Okon, O. B. (2009). Anti-spermatogenic Effects of Ethanol Extract of *Mucuna Urens*. *Journal of Reproduction and Contraception*, 20(3), 161–168. [https://doi.org/10.1016/S1001-7844\(09\)60021-3](https://doi.org/10.1016/S1001-7844(09)60021-3)
- FAO. (2022). *The state of world fisheries and Aquaculture, 2022, Towards Blue transformation*.
- Fitzgerald, W. (1979). The red-orange tilapia: a hybrid that could become a world favorite. *Fish Framing Int*, 6(1), 26–27.
- Flefil, N. S., Ezzat, A., Aboseif, A. M., & Negm El-Dein, A. (2022). Lactobacillus-fermented wheat bran, as an economic fish feed ingredient, enhanced dephytinization, micronutrient bioavailability, and tilapia performance in a biofloc system. *Biocatalysis and Agricultural Biotechnology*, 45. <https://doi.org/10.1016/J.BCAB.2022.102521>
- Fraser, D. R. (2018). Evolutionary Biology: Mysteries of Vitamin D in Fish. *Vitamin D: Fourth Edition*, 1, 13–27. <https://doi.org/10.1016/B978-0-12-809965-0.00002-1>
- Gabriel, N. N. (2019). Review on the progress in the role of herbal extracts in tilapia culture. In *Cogent Food and Agriculture* (Vol. 5, Issue 1). <https://doi.org/10.1080/23311932.2019.1619651>
- Gabriel, N. N., Qiang, J., Ma, X. Y., He, J., Xu, P., & Omoregie, E. (2017). Sex-Reversal Effect of Dietary Aloe vera (*Liliaceae*) on Genetically Improved Farmed Nile Tilapia Fry. *North American Journal of Aquaculture*, 79(1). <https://doi.org/10.1080/15222055.2016.1236046>
- Genschick, S., Mekkawy, W., Rossignoli, C., & Benzie, J. A. H. (2021). Growth performance of three strains of Nile tilapia (*Oreochromis niloticus*) on four different feeds in Western and Central Kenya. *Aquaculture Reports*, 20. <https://doi.org/10.1016/j.aqrep.2021.100701>
- Guimarães, M. C., Dias, D. de C., de Araujo, F. von A. P., Ishikawa, C. M., & Tachibana, L. (2019). PROBIOTIC bacillus subtilis AND lactobacillus plantarum IN DIET OF NILE TILAPIA. *Boletim Do Instituto de Pesca*, 45(1). <https://doi.org/10.20950/1678-2305.2019.45.1.252>
- Harikrishnan, R., Thamizharasan, S., Devi, G., Van Doan, H., Ajith Kumar, T. T., Hoseinifar, S. H., & Balasundaram, C. (2020). Dried lemon peel enriched diet improves antioxidant activity, and immune response and modulates immuno-antioxidant genes in Labeorohita against *Aeromonas sorbia*. *Fish & Shellfish Immunology*, 106, 675–684. <https://doi.org/10.1016/J.FSI.2020.07.040>
- HEARN, T. L., SGOUTAS, S. A., HEARN, J. A., & SGOUTAS, D. S. (1987). Polyunsaturated Fatty Acids and Fat in Fish Flesh for Selecting Species for Health Benefits. *Journal of Food Science*, 52(5), 1209–1211. <https://doi.org/10.1111/J.1365-2621.1987.TB14045.X>
- Hernández, E., Figueroa, J., & Iregui, C. (2009). Streptococcosis on a red tilapia, *Oreochromis sp.*, farm: A case study. *Journal of Fish Diseases*, 32(3), 247–252. <https://doi.org/10.1111/J.1365-2761.2008.00981.X>

- Higgs, D. A., Fagerlund, U. H. M., Eales, J. G., & McBride, J. R. (1982). Application of thyroid and steroid hormones as anabolic agents in fish culture. *Comparative Biochemistry and Physiology - Part B: Biochemistry And*, 73(1), 143–176. [https://doi.org/10.1016/0305-0491\(82\)90206-1](https://doi.org/10.1016/0305-0491(82)90206-1)
- Hishamunda, N., & Ridler, N. B. (2006). Farming fish for profits: A small step towards food security in sub-Saharan Africa. *Food Policy*, 31(5), 401–414. <https://doi.org/10.1016/J.FOODPOL.2005.12.004>
- Hoseinifar, S. H., Sun, Y. Z., & Caipang, C. M. (2017). Short-chain fatty acids as feed supplements for sustainable aquaculture: an updated view. *Aquaculture Research*, 48(4), 1380–1391. <https://doi.org/10.1111/ARE.13239>
- Hossain, M. I., Haque, N., Alam, A. S. A. F., Begum, H., & Mokhtar, M. Bin. (2019). Sinking and floating feeds formulations on the growth of mono-sex tilapia (Linnaeus, 1758): A comparative study. *Turkish Journal of Fisheries and Aquatic Sciences*, 19(1). https://doi.org/10.4194/1303-2712-v19_1_08
- Karapanagiotidis, I. T., Metsoviti, M. N., Gkalogianni, E. Z., Psafakis, P., Asimaki, A., Katsoulas, N., Papapolymerou, G., & Zarkadas, I. (2022). The effects of replacing fishmeal by *Chlorella vulgaris* and fish oil by *Schizochytrium sp.* and *Microchloropsis gaditana* blend on growth performance, feed efficiency, muscle fatty acid composition, and liver histology of gilthead seabream (*Sparus aurata*). *Aquaculture*, 561. <https://doi.org/10.1016/j.aquaculture.2022.738709>
- Khanjani, M. H., Alizadeh, M., Mohammadi, M., & Aliabad, H. S. (2021). The Effect of Adding Molasses at Different Times on the Performance of Nile *Tilapia (Oreochromis niloticus)* Raised in a Low-Salinity Biofloc System. *Annals of Animal Science*, 21(4), 1435–1454. <https://doi.org/10.2478/AOAS-2021-0011>
- Khorshidi, Z., Paknejad, H., Sodagar, M., Hajimoradloo, A., & Shekarabi, S. P. H. (2022). Effect of a commercial multi-effect additive (Biotronic® Top3) on growth performance, digestive enzymes, and intestinal barrier gene expression in common carp (*Cyprinus carpio*). *Aquaculture*, 560, 738588. <https://doi.org/10.1016/J.AQUACULTURE.2022.738588>
- Kord, M. I., Maulu, S., Srour, T. M., Omar, E. A., Farag, A. A., Nour, A. A. M., Hasimuna, O. J., Abdel-Tawwab, M., & Khalil, H. S. (2022). Impacts of water additives on water quality, production efficiency, intestinal morphology, gut microbiota, and immunological responses of Nile tilapia fingerlings under a zero-water-exchange system. *Aquaculture*, 547. <https://doi.org/10.1016/j.aquaculture.2021.737503>
- Kumar, S., J. Sándor, Z., Biró, J., Gyalog, G., Kumar Sinha, A., & De Boeck, G. (2022). Does nutritional history impact future performance and utilization of plant-based diet in common carp? *Aquaculture*, 551, 737935. <https://doi.org/10.1016/J.AQUACULTURE.2022.737935>
- Lin, H., Jia, Y., Han, F., Xia, C., Zhao, Q., Zhang, J., & Li, E. (2022). Toxic effects of waterborne benzyl paraben on the growth, antioxidant capacity and lipid metabolism of Nile tilapia (*Oreochromis niloticus*). *Aquatic Toxicology*, 248. <https://doi.org/10.1016/j.aquatox.2022.106197>
- Lin, Y. H., & Shiau, S. Y. (2007). Effects of a dietary blend of fish oil with corn oil on growth and non-specific immune responses of grouper, *Epinephelus malabaricus*. *Aquaculture Nutrition*, 13(2), 137–144. <https://doi.org/10.1111/J.1365-2095.2007.00458.X>
- Liu, Y., Jiao, J. G., Gao, S., Ning, L. J., Mchele Limbu, S., Qiao, F., Chen, L. Q., Zhang, M. L., & Du, Z. Y. (2019). Dietary oils modify lipid molecules and nutritional value of fillet in Nile tilapia: A

- deep lipidomics analysis. *Food Chemistry*, 277, 515–523. <https://doi.org/10.1016/j.foodchem.2018.11.020>
- Lock, E. J., Waagbø, R., Wendelaar Bonga, S., & Flik, G. (2010). The significance of vitamin D for fish: A review. *Aquaculture Nutrition*, 16(1), 100–116. <https://doi.org/10.1111/J.1365-2095.2009.00722.X>
- Luo, G., Xu, J., Teng, Y., Ding, C., & Yan, B. (2010). Effects of dietary lipid levels on the growth, digestive enzyme, feed utilization, and fatty acid composition of Japanese sea bass (*Lateolabrax japonicus* L.) reared in freshwater. *Aquaculture Research*, 41(2), 210–219. <https://doi.org/10.1111/J.1365-2109.2009.02319.X>
- Mamdouh, A. Z., Zahran, E., Mohamed, F., & Zaki, V. (2021). Nannochloropsis oculata feed additive alleviates mercuric chloride-induced toxicity in Nile tilapia (*Oreochromis niloticus*). *Aquatic Toxicology*, 238. <https://doi.org/10.1016/j.aquatox.2021.105936>
- Mohammadi, G., Hafezieh, M., Karimi, A. A., Azra, M. N., Van Doan, H., Tapingkae, W., Abdelrahman, H. A., & Dawood, M. A. O. (2022). The synergistic effects of plant polysaccharide and *Pediococcus acidilactici* as a synbiotic additive on growth, antioxidant status, immune response, and resistance of Nile tilapia (*Oreochromis niloticus*) against *Aeromonas hydrophila*. *Fish and Shellfish Immunology*, 120, 304–313. <https://doi.org/10.1016/j.fsi.2021.11.028>
- Mukherjee, D., Ghosal, I., & Chakraborty, S. B. (2015). Production of monosex Nile tilapia, *Oreochromis niloticus* using seed of *Mucuna pruriens*. *IOSR Journal of Pharmacy and Biological Sciences*, 10(1).
- Mukhopadhyay, P. K., & Rout, S. K. (1996). Effects of different dietary lipids on growth and tissue fatty acid changes in fry of the carp *Catla catla* (Ham). *Aquaculture Research*, 27(8), 623–630. <https://doi.org/10.1111/J.1365-2109.1996.TB01295.X/ABSTRACT>
- Mukhopadhyay, S. (2021). *SYNTHETIC AND SYSTEMS BIOLOGY METHODS FOR APPLICATION IN GENE CIRCUITS AND MICRO-GRAVITY-RELATED SPACE BIOLOGY*.
- Nuhu Ozovehe, B. (2013). Growth Performance, Haematological Indices and Some Biochemical Enzymes of Juveniles *Clarias gariepinus* (Burchell 1822) Fed Varying Levels of Moringa oleifera Leaf Meal Diet. <https://doi.org/10.4172/2155-9546.1000166>
- Oglend, A. (2020). Challenges and opportunities with aquaculture growth. <https://doi.org/10.1080/13657305.2019.1704937>, 24(2), 123–127. <https://doi.org/10.1080/13657305.2019.1704937>
- Ojha, M., Chadha, N., Saini, V., Damroy Director, S., Sawant, P., & Damroy, S. (2014). Role of *Petalium murex* in the enhancement of growth, metabolism and immunity of Labeorohita (Hamilton, 1822) fingerlings. 70 ~ *International Journal of Fauna and Biological Studies*, 1(6), 70–77.
- Okomoda, V. T., Tihamiyu, L. O., & Akpan, I. S. (2017). Nutritional evaluation of toasted *Mucuna utilis* seed meal and its utilization in the diet of *Clarias gariepinus* (Burchell, 1822). <http://dx.doi.org/10.1080/10454438.2017.1278733>, 29(2), 167–182. <https://doi.org/10.1080/10454438.2017.1278733>
- Oliveira, L. K., Pilz, L., Furtado, P. S., Ballester, E. L. C., & Bicudo, Á. J. de A. (2021). Growth, nutritional efficiency, and profitability of juvenile GIFT strain of Nile tilapia (*Oreochromis niloticus*) reared in biofloc system on graded feeding rates. *Aquaculture*, 541. <https://doi.org/10.1016/j.aquaculture.2021.736830>

- Opiyo, M. A., Jumbe, J., Ngugi, C. C., & Charo-Karisa, H. (2019). Different levels of probiotics affect the growth, survival, and body composition of Nile tilapia (*Oreochromis niloticus*) cultured in low-input ponds. *Scientific African*, 4. <https://doi.org/10.1016/J.SCIAF.2019.E00103>
- Opiyo, M. A., Marijani, E., Muendo, P., Odede, R., Leschen, W., & Charo-Karisa, H. (2018). A review of aquaculture production and health management practices of farmed fish in Kenya. *International Journal of Veterinary Science and Medicine*, 6(2), 141–148. <https://doi.org/10.1016/j.ijvsm.2018.07.001>
- Owatari, M. S., da Silva, L. R., Ferreira, G. B., Rodhermel, J. C. B., de Andrade, J. I. A., Dartora, A., & Jatobá, A. (2022). Body yield, growth performance, and hematological evaluation of Nile tilapia fed a diet supplemented with *Saccharomyces cerevisiae*. *Animal Feed Science and Technology*, 293, 115453. <https://doi.org/10.1016/j.anifeeds.2022.115453>
- Peng, S., Chen, L., Qin, J. G., Hou, J., Yu, N., Long, Z., Ye, J., & Sun, X. (2008). Effects of replacement of dietary fish oil by soybean oil on growth performance and liver biochemical composition in juvenile black seabream, *Acanthopagrus schlegelii*. *Aquaculture*, 276(1–4), 154–161. <https://doi.org/10.1016/J.AQUACULTURE.2008.01.035>
- Poolsawat, L., Yang, H., Sun, Y. F., Li, X. Q., Liang, G. Y., & Leng, X. J. (2021). Effect of replacing fish meal with enzymatic feather meal on growth and feed utilization of tilapia (*Oreochromis niloticus* × *O. aureus*). *Animal Feed Science and Technology*, 274. <https://doi.org/10.1016/J.ANIFEEDSCI.2021.114895>
- Prabhu, P. A. J., Lock, E. J., Hemre, G. I., Hamre, K., Espe, M., Olsvik, P. A., Silva, J., Hansen, A. C., Johansen, J., Sissener, N. H., & Waagbø, R. (2019). Recommendations for dietary level of micro-minerals and vitamin D3 to Atlantic salmon (*Salmo salar*) parr and post-smolt when fed low fish meal diets. *PeerJ*, 2019(5). <https://doi.org/10.7717/PEERJ.6996>
- Rohani, M. F., Tarin, T., Hasan, J., Islam, S. M. M., & Shahjahan, M. (2023). Vitamin E supplementation in diet ameliorates the growth of Nile tilapia by upgrading muscle health. *Saudi Journal of Biological Sciences*, 30(2). <https://doi.org/10.1016/j.sjbs.2023.103558>
- Romero, J., Gloria, C., & Navarrete, P. (2012). Antibiotics in Aquaculture – Use, Abuse and Alternatives. *Health and Environment in Aquaculture*. <https://doi.org/10.5772/28157>
- Saeed, F., Arshad, M. U., Pasha, I., Naz, R., Batool, R., Khan, A. A., Nasir, M. A., & Shafique, B. (2014). The nutritional and phyto-therapeutic potential of papaya (*Carica Papaya* Linn.): An overview. *International Journal of Food Properties*, 17(7), 1637–1653. <https://doi.org/10.1080/10942912.2012.709210>
- Saha, S., Ray, A. K., Sümbülü, S., Ununun, Y., Bağırşagından, B., İki, İ. E., Tüürü, B., Sonrası, F., Rohu, Y., Rohita, L., Besinsel, D., & Değerlendirilmesi, K. (2011). Evaluation of Nutritive Value of Water Hyacinth (*Eichhorniacrassipes*) Leaf Meal in Compound Diets for Rohu, *Labeorohita* (Hamilton, 1822) Fingerlings after Fermentation with Two Bacterial Strains Isolated from Fish Gut. *Turkish Journal of Fisheries and Aquatic Sciences*, 11, 199–207. <https://doi.org/10.4194/trjfas.2011.0204>
- Sarmiento, N. L. A. F., Martins, E. F. F., Costa, D. C., Mattioli, C. C., da Costa Julio, G. S., Figueiredo, L. G., Luz, M. R., & Luz, R. K. (2018). Reproductive efficiency and egg and larvae quality of Nile tilapia fed different levels of vitamin C. *Aquaculture*, 482, 96–102. <https://doi.org/10.1016/j.aquaculture.2017.08.035>
- Shenouda, S. Y. K. (1980). Theories of protein denaturation during frozen storage of fish flesh. *Advances in Food Research*, 26(C), 275–311. [https://doi.org/10.1016/S0065-2628\(08\)60320-1](https://doi.org/10.1016/S0065-2628(08)60320-1)

- Shore, L. S., & Shemesh, M. (2003). Naturally produced steroid hormones and their release into the environment. *Pure and Applied Chemistry*, 75(11–12), 1859–1871. <https://doi.org/10.1351/PAC200375111859>
- Sørensen, M. (2012). A review of the effects of ingredient composition and processing conditions on the physical qualities of extruded high-energy fish feed as measured by prevailing methods. *Aquaculture Nutrition*, 18(3), 233–248. <https://doi.org/10.1111/J.1365-2095.2011.00924.X>
- SP, D., G, R., K, R., G, V., & R, S. (2015). Effect of Fish Oil Substitution with Sunflower Oil in Diet of Juvenile Catlacatla (*Ham*) on Growth Performance and Feed Utilization. *Journal of Fisheries & Livestock Production 2015 3:3*, 3(3), 1–3. <https://doi.org/10.4172/2332-2608.1000144>
- Straus, D. L., Bowker, J. D., Bowman, M. P., Carty, D. G., Mitchell, A. J., Farmer, B. D., & Ledbetter, C. K. (2013). Safety of feed treated with 17 α -methyltestosterone (17MT) to larval Nile Tilapia. *North American Journal of Aquaculture*, 75(2). <https://doi.org/10.1080/15222055.2012.758211>
- Syanya F.J, & Mathia W.Munala. (2022). *Moving towards sustainable Aquaculture for Rural sustainability and development in Kenya; A case of Vihiga county.*
- Syed, R., Masood, Z., Ul Hassan, H., Khan, W., Mushtaq, S., Ali, A., Gul, Y., Jafari, H., Habib, A., Ishaq Ali Shah, M., Gabol, K., Gul, H., & Ullah, A. (2022). Growth performance, hematological assessment and chemical composition of Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) fed different levels of Aloe vera extract as feed additives in a closed aquaculture system. *Saudi Journal of Biological Sciences*, 29(1), 296–303. <https://doi.org/10.1016/J.SJBS.2021.08.098>
- Tacon, A. G. J., & Metian, M. (2015). Feed matters: Satisfying the feed demand of aquaculture. *Reviews in Fisheries Science and Aquaculture*, 23(1), 1–10. <https://doi.org/10.1080/23308249.2014.987209>
- Tuan Harith, Z., MohdSukri, S., Remlee, N. F. S., Mohd Sabir, F. N., & Zakaria, N. N. A. (2022a). Effects of dietary astaxanthin enrichment on enhancing the colour and growth of red tilapia, *Oreochromis sp.* *Aquaculture and Fisheries*. <https://doi.org/10.1016/j.aaf.2022.06.001>
- Tuan Harith, Z., MohdSukri, S., Remlee, N. F. S., Mohd Sabir, F. N., & Zakaria, N. N. A. (2022b). Effects of dietary astaxanthin enrichment on enhancing the colour and growth of red tilapia, *Oreochromis sp.* *Aquaculture and Fisheries*. <https://doi.org/10.1016/J.AAF.2022.06.001>
- Van Doan, H., Lumsangkul, C., Hoseinifar, S. H., Tongsiri, S., Chitmanat, C., Musthafa, M. S., El-Haroun, E., & Ringo, E. (2021). Modulation of growth, innate immunity, and disease resistance of Nile tilapia (*Oreochromis niloticus*) culture under biofloc system by supplementing pineapple peel powder and *Lactobacillus plantarum*. *Fish and Shellfish Immunology*, 115, 212–220. <https://doi.org/10.1016/j.fsi.2021.06.008>
- Waagbø, R. (2010). Water-soluble vitamins in fish ontogeny. *Aquaculture Research*, 41(5), 733–744. <https://doi.org/10.1111/J.1365-2109.2009.02223.X>
- Wangkahart, E., Kersanté, P., Lee, P. T., Sanbut, O., Nontasan, S., & Chantiratikul, A. (2022). Effect of Kera-Stim@50, a feed additive containing free amino acid mix on growth, antioxidant and immune responses, digestive enzymes, and fatty acid composition in Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 551. <https://doi.org/10.1016/j.aquaculture.2021.737874>
- Wang, L. mei, Bu, H. yi, Song, F. biao, Zhu, W. bin, Fu, J. jun, & Dong, Z. jie. (2019). Characterization and functional analysis of slc7a11 gene, involved in skin color differentiation in the red tilapia. *Comparative Biochemistry and Physiology -Part A : Molecular and Integrative Physiology*, 236. <https://doi.org/10.1016/j.cbpa.2019.110529>

- Wan-Mohtar, W. A. A. Q. I., Taufek, N. M., Thiran, J. P., Rahman, J. F. P., Yerima, G., Subramaniam, K., & Rowan, N. (2021). Investigations on the use of exopolysaccharide derived from the mycelial extract of *Ganoderma lucidum* as a functional feed ingredient for aquaculture-farmed red hybrid Tilapia (*Oreochromis sp.*). *Future Foods*, 3. <https://doi.org/10.1016/J.FUFO.2021.100018>
- Wiszniewski, G., Jarmolowicz, S., Hassaan, M. S., Soady, M. R., Kamaszewski, M., Szudrowicz, H., Terech-Majewska, E., Pajdak-Czaus, J., Wiechetek, W., & Siwicki, A. K. (2022). Beneficial effects of dietary papain supplementation in juvenile sterlet (*Acipenser ruthenus*): Growth, intestinal topography, digestive enzymes, antioxidant response, immune response, and response to a challenge test. *Aquaculture Reports*, 22. <https://doi.org/10.1016/j.aqrep.2021.100923>
- Wohlfarth, G. W., Rothbard, S., Hulata, G., & Zweigman, D. (1990). Inheritance of red body coloration in Taiwanese tilapias and in *Oreochromis mossambicus*. *Aquaculture*, 84(3–4), 219–234. [https://doi.org/10.1016/0044-8486\(90\)90088-5](https://doi.org/10.1016/0044-8486(90)90088-5)
- Wu, J. Y., Feng, L., Wu, P., Liu, Y., Ren, H. M., Jin, X. W., Jiang, J., Kuang, S. Y., Li, S. W., Tang, L., Zhang, L., Zhou, X. Q., & Jiang, W. D. (2022). Modification of beneficial fatty acid composition and physicochemical qualities in the muscle of sub-adult grass carp (*Ctenopharyngodon idella*): The role of lipids. *Aquaculture*, 561. <https://doi.org/10.1016/j.aquaculture.2022.738656>
- Xu, C., Li, E., Xu, Z., Su, Y., Lu, M., Qin, J. G., Chen, L., & Wang, X. (2018). Growth and Stress Axis Responses to Dietary Cholesterol in Nile Tilapia (*Oreochromis niloticus*) in Brackish Water. *Frontiers in Physiology*, 9(MAR), 254. <https://doi.org/10.3389/fphys.2018.00254>
- Xu, W., Mawolo, P. Y., Gao, J., Chu, L., Wang, Y., Nie, Z., Song, L., Shao, N., Gao, J., Xu, P., & Xu, G. (2021). Effects of supplemental effective microorganisms in feed on the growth, immunity, and appetite regulation in juvenile GIFT tilapia. *Aquaculture Reports*, 19. <https://doi.org/10.1016/J.AQREP.2020.100577>
- Yamazaki, F. (1976). Application of Hormones in Fish Culture. *Journal of the Fisheries Research Board of Canada*, 33(4), 948–958. <https://doi.org/10.1139/F76-122>
- Yossa, R., Greiling, A. M., Basiita, R. K., Sakala, M. E., Baumgartner, W. A., Taylor, A., & Gatlin, D. M. (2021). Replacing fishmeal with a single cell protein feedstuff in Nile tilapia *Oreochromis niloticus* diets. *Animal Feed Science and Technology*, 281. <https://doi.org/10.1016/J.ANIFEEDSCI.2021.115089>
- Yu, X., Xin, Y., Cui, L., Jia, J., Yuan, X., Fu, S., Zhang, J., Sun, C., Miao, X., & Li, W. (2021a). Effects of neuropeptide Y as a feed additive on stimulating the growth of tilapia (*Oreochromis niloticus*) fed low fish meal diets. *Peptides*, 138. <https://doi.org/10.1016/j.peptides.2021.170505>
- Yu, X., Xin, Y., Cui, L., Jia, J., Yuan, X., Fu, S., Zhang, J., Sun, C., Miao, X., & Li, W. (2021b). Effects of neuropeptide Y as a feed additive on stimulating the growth of tilapia (*Oreochromis niloticus*) fed low fish meal diets. *Peptides*, 138, 170505. <https://doi.org/10.1016/J.PEPTIDES.2021.170505>
- Zhou, M., Yu, S., Hong, B., Li, J., Han, H., & Qie, G. (2021). Antibiotics control in aquaculture requires more than antibiotic-free feeds: A tilapia farming case. *Environmental Pollution*, 268. <https://doi.org/10.1016/J.ENVPOL.2020.115854>