

[Original Research](#)

“Impact of organic nutrient sources on growth and yield of rice varieties (*Oryza sativa* L.) and yield validation using SPSS model”

ABSTRACT

A field experiment was conducted during Kharif 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P) to determine the “**Impact of organic nutrient sources on growth and yield of rice varieties (*Oryza sativa* L.) and Yield Validation Using SPSS Model**”. The experiment was laid out in Randomized Block Design comparing of 9 treatment which include 3 varieties RNR-15048, BPT-5204 AND NDR-359, 4 organic manures, Vermicompost 10(t/ha) and Panchagavya 6%+ Vermicompost 5(t/ha)+ Panchagavya 3%. Whose effect is observed in varieties rice. The treatments consisted of organic nutrient sources and rice varieties. [Panchagavya 6%+Vermicompost 10t/ha, Panchagavya 3%+Vermicompost 5t/ha]. and the rice varieties [RNR-15048, BPT- 5204, NDR-359]. whose effect is observed in paddy. The results observed that the treatment with application of NDR-359 + vermicompost 5t/ha + panchagavya 3% recorded higher plant height, number of tillers/hill, plant dry weight(g), CGR, RGR and yield parameter number of panicles/hill, number of grains/panicles, number of grains, test weight, grain yield, straw yield and harvest index. Treatment 9 has shown 44.62% increase over predicted yield whereas there was 28.89% increase in treatment 8 over predicted yield through SPSS model. From the basis of one year experimentation, it is concluded that Panchagavya 3% + Vermicompost 5(t/ha) was found more productive for the variety NDR-359.

Keywords: organic nutrients; varieties; growth; yield; yield validation using SPSS Model

Introduction

Rice (*Oryza sativa* L.) is one of the most important staple food crops as it helps to sustain two thirds of the world's population **Kahani et al., (2015)**. India is an important centre of rice cultivation. Rice is a nutritional staple food which provides instant energy as its most important component is carbohydrate (starch), it also provides 27% of dietary energy supply, 20% of dietary protein and 3% of dietary fat. It can contribute nutritionally significant amounts of thiamine, riboflavin, niacin and zinc to diet, but smaller amounts of other micronutrients.

In Asia, rice is a major source of calories, especially for the poor, which eat from 50 to 80 percent of the daily calories from it. In Asia, where it is a staple of culture and tradition, and over 90% of the world's rice is grown and eaten. While it is true that rice is not a tropical plant, it is believed to belong with moist, humid areas. China (32.7%), India (26.0%), Thailand (5.3%), Myanmar (4.8%), the Philippines (2.8%), Brazil (2.0%), and Japan (1.9%) are the top 10 suppliers of rice in the world (**Anonymus, 2012**). To name a few other uses, the 85% of the rice is also used in cerelac, snack foods, brewed beverages, flour, oil, syrup, and religious rituals.

Organic recycling and organic farming could survive the dual purpose of minimising pollution and utilising waste in your farm from for increasing the soil productivity. **Kenchaiah (1997)** observed a positive influence of farm manure on rice plant tiller number per hill and dry weight production with increasing a yield and yield. **Moorthy(1994)** obtained higher plant growth attributes and grain yield in low land rice applied with *Sesbania aculeate*. The plant growth characters and yield were marked influenced by poultry manure and panchagavya so waste composed can increase in the yield of many crop like rice groundnut sorghum pearl millet and taint condition crops,**Somasundaram et al., (2003)**.

Ayurveda uses the term "panchagavya" to describe a fermented product formed from five components obtained from cows, including milk, urine, dung, and bile. clarified butter and curd. In India, the contribution of foliar panchagavya to the production of numerous plantation crops has received considerable attention (**Selvaraj, 2003**). Often used for horticultural and agricultural crops, Panchagavya is a well-known foliar nutrition made by organic producers in Tamil Nadu (**Swaminathan et al., 2007**). The goal of the current study was to test the hypothesis that foliar panchagavya administration will have on various physiological parameters, yield.

Panchagavya, an organic product has potential to play the role in promoting immunity growth and providing immunity in plant system. The use of organic liquid such as panchagavya results in higher growth, yield and quality of crops. Different species of insect pests were reported to be associated with spinach in different areas of the country. Usually, the management of the pests is insecticides oriented but the problems associated with synthetic chemicals with., development of pest resistance objectionable pesticides residue and higher cost etc, has necessitated development of new control methods. Several plants and its products are known to be potential resources. (**Choudhary et al.,2014**). **Beulah et al., (2002)** concluded that the beneficial microorganisms

from panchagavya and their presence in the rhizospheres environment of the root zone influence the plant growth and crop yield.

Vermicompost has been considered as a soil additive to reduce the use of mineral fertilizers because it provides required nutrient amounts, increases cation exchange capacity and improves water holding capacity (**Tejada and Gonzaler, 2009**). Vermicompost not only increases yield of rice but can also substitute chemical fertilizer to some extent (**Sharma *et al.*, 2008; Guera, 2010**).

Vermicomposting is an effective biological process for conversion of organic wastes into a stable end product, where in microbial activity plays an essential role. Earthworms are mainly responsible for fragmentation and conditioning of the substrate, increasing surface area for microbial activity and significantly altering biological activity of the process (**Domynguez 2004**). Vermicompost is homogenous, with desirable aesthetics and possess plant growth hormones and high levels of soil enzymes, while enhancing microbial population and tending to hold more nutrients over longer periods without adverse impacts on the environment. In India, about 320 million tonnes of agricultural wastes are generated annually. The vermicompost is a rich source of beneficial microorganisms and nutrients (**Pant 2008**). Increase in crop yield, soil nutrients status and nutrients uptake was reported due to application of vermicompost.

MATERIALS AND METHODS:

The experiment was conducted during *Kharif* season of 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of the field constituting a part of central gangetic alluvium is neutral and deep. The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH 7.8), low level of organic carbon (0.51%), available N (78.9 kg/ha), available P (32.88 kg/ha) and available K (385.10 kg/ha). Nutrient sources were Panchagavya and vermicompost to fulfilled with Nitrogen, Phosphorus, Potassium respectively. The treatments were RNR-15048+ Vermicompost 10 (t/ha), RNR-15048+ Panchagavya-6%, RNR-15048+ Vermicompost 5 (t/ha) + Panchagavya-3%, BPT -5204 + Vermicompost 10 (t/ha), BPT-5204 + Panchagavya-6%, BPT-5204+ Vermicompost (5t/ha) + Panchagavya-3% , NDR-359+ Vermicompost 10 (t/ha), NDR-359 + Panchagavya-6%, NDR-359 Vermicompost 5 (t/ha) + Panchagavya - 3%. The experiment was laid out in Randomized Block Design, with 9 treatments replicated thrice. The observations were recorded for plant height (cm), number of tillers/hill, plant dry weight(g), Crop growth rate (g/m²/day), Relative growth rate (g/g/day), Number of effective tillers, number of panicles/hill, number of grains/panicles, number of grains, test weight (g), grain yield (t/ha), straw yield (t/ha) and harvest index (%). The collected data was subjected to statistical analysis by analysis of variance (ANOVA) was used statically to examine these variables using SPSS (Statistical Product and Service Solutions). The sum of weather parameters and the sum product of several weather parameters were utilised to determine the correlation coefficient. Many regressions have been carried out using the dependent variable (yield) and the independent variables (time, sum, and sum products depending on meteorological conditions). The regression equation has been generated using the regression formula.

3. RESULTS AND DISCUSSION

Growth parameters

Plant height :

The data revealed that significant and higher plant height (112.6 cm) was observed in treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%], and there was no statistically significant difference at par value.

The results of the present study are also in line with **Emily (2003)** who reported that, the plant height, leaf area, leaf area index, total chlorophyll content and dry matter production increased significantly due to spraying panchagavya 4 %. **Vennila and Jayanthi (2008)** revealed that application of 100% recommended dose of fertilizer along with panchagavya spray (2%), increased plant height of crops with vermicompost application had also been reported by **Thanunathan *et al.*, (2001)**.

Number of Tillers/plant :

The data revealed that treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%] recorded significant and maximum number of tillers/hill (12.7) and there was no statistically significant difference at par value.

An increase in the number of productive tillers could be attributed to the improvement in soil physical and chemical properties and better availability of nutrients by the addition of fertilizer,vermicompost.Similar findings were also reported by **Kamakar *et al.*, (2010)** increase the numbers of effective tillers.

Plant dry weight (g):

Results revealed that treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%] recorded significantly higher plant dry weight (52.3 g).which was superior to all the treatment and treatment T₇ with the application of [NDR-359 + Vermicompost(10t/ha)].

The probable reason for higher dry matter production might the application of vermicompost stimulated the plant growth due the higher microbial activity and soil reaction and large portion of nitrogen in vermicompost in organic fractions and fermented solution of panchagavya contains various salts rich in N, P, K, S and micronutrients in plant available form. Hence, availability of these nutrients to plants helps in the higher dry matter production in plants. The results were also found to be similar with findings of **Shrimal and Khan (2017)** and **Gopal *et al.*, (2007)**

Crop Growth Rate (g/m²/day):

The data recorded that during 80-100DAS that treatment T₉ [ZnO (900 ppm)+ Boron (0.5%)] was highest crop growth rate (26.44 g/m²/day). which was superior to all the treatment and treatment T₇ with the application of [BPT-5204 + Panchagavya 6%]

Releative Growth Rate (g/g/day):

The data revealed that during 80-100,treatment T₈ [NDR-359 + Panchagavya 6%], and there was no stastically at par value.

Yield parameter:**Number of effective tillers/hill:**

The data showed that treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%] recorded significantly number of effective tillers/hill (12.7/hill) and there was no stastically at par value.

Number of panicle/hill:

The data showed that treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%] recorded significantly higher number of panicle/hill (7.6/hill) and there were no stastically at par value.

Number of grain/panicle :

The data recorded that significant and higher number of grains/panicle (114.7), and there were no stastically at par value.

Test weight (g):

Significant and maximum test weight (21.8) was recorded in treatment T₉[NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%]. However, treatment T₇ [NDR-359 + Panchagavya 6%] was stastically at par with the treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%].

Grain yield(t/ha):

Significant and higher grain yield (5.64 t/ha) was obtained in treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%], and there were no stastically at par value.

Straw yield (t/ha):

Significant and higher stover yield (7.34 t/ha) was obtained in treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%], and there were no stastically at par value.

Harvest index (%):

Significant and higher harvest index (41.63 %) was obtained in treatment T₉ [NDR-359 + Vermicompost(5t/ha) + Panchagavya 3%], which was superior to all the treatment and treatment T₆ with the application of [BPT-5204 + Vermicompost (5 t/ha)+ Panchagavya 3%], T₃ application of RNR-15048+Vermicompost(5t/ha) + Panchagavya 3%.

When the ten randomly selected plants in each plot were harvested, the number of panicles (productive tillers) per hill, panicle length, total number of grains, number of filled grains per panicle, and thousand grains weight were noted (**Hemalatha et al., 2000**).

The higher increase in the yield has been reported to be associated with the release of macro and micro nutrients during the course of microbial decomposition. Organic matter also functions as source of energy for soil micro flora which brings about the transformation of other nutrients held in soil or applied through other means, in a form that is readily utilized by growing plants which helped in increase of seed yield. The results were in accordance with **Jadhav et al., (2011)** yield attributes which eventually helped in increase in seed yield. The results were found to be similar with **Vimalendran and Wahab (2013)**.

The maximum grain and straw yield was due to marked improvement in dry matter accumulation, yield attributes and greater nutrient content and their uptake by rice crop. These findings are in direct conformity with that of **Barik et al., (2008)**.

Yield validation using SPSS model

The multi-regression analysis using SPSS has been employed for the estimation of rice yield. The regression for SPSS model is $Y = 15.266 + (0.292 \times Z21 \text{ of prediction year}) + (0.066 \times \text{time})$ Here, Z21 is the sum product of minimum temperature. The yield obtained in treatment 9 with NDR-359 + Vermicompost (5 t/ha) + Panchagavya 3% (5.6 t/ha) showed 44.62% increase over the predicted yield through SPSS model (2.88 t/ha).

Table 1. Impact of Organic nutrient sources and growth attributes of rice varieties.

| SI No | Treatment | 100 DAT | | | 80-100 DAT | |
|--------------------|---|-------------------|------------------------|-------------|-----------------------------|---------------|
| | | Plant height (cm) | Number of tillers/hill | Dry weight | CGR (g/m ² /day) | RGR (g/g/day) |
| 1 | RNR-15048 + Vermicompost 10(t/ha) | 108.5 | 10.8 | 50.0 | 25.93 | 0.012 |
| 2 | RNR-15048 + Panchagavya 6% | 107.4 | 10.4 | 49.3 | 25.65 | 0.012 |
| 3 | RNR-15048+Vermicompost(5t/ha) + Panchagavya 3% | 111.2 | 11.8 | 51.1 | 24.22 | 0.011 |
| 4 | BPT-5204 + Vermicompost 10(t/ha) | 107.9 | 10.5 | 49.7 | 25.89 | 0.012 |
| 5 | BPT-5204 + Panchagavya 6% | 105.2 | 10.3 | 49.5 | 26.44 | 0.012 |
| 6 | BPT-5204 + Vermicompost(5t/ha) + Panchagavya 3% | 110.5 | 11.5 | 50.7 | 26.20 | 0.011 |
| 7 | NDR-359 + Vermicompost 10(t/ha) | 111.8 | 12.1 | 51.9 | 25.47 | 0.011 |
| 8 | NDR-359 + Panchagavya 6% | 109.7 | 11.2 | 50.6 | 26.44 | 0.013 |
| 9 | NDR-359 + Vermicompost(5t/ha) + Panchagavya 3% | 112.6 | 12.7 | 52.3 | 22.78 | 0.009 |
| F test | | S | S | S | S | S |
| SEm(±) | | 0.20 | 0.13 | 0.12 | 0.47 | 0.0003 |
| CD (P=0.05) | | 0.61 | 0.39 | 0.37 | 1.41 | 0.009 |

Table 2. Impact of Organic nutrient sources and yield attributes of rice varieties

| SL. NO | TREATMENTS | No. Of effective tillers/hill | No.of grains/pa nicle | No.of panicles/ hill | Test weight (g) | Grain yield (t/ha) | Stover yield (t/ha) | Harvest index(%) |
|---------------|---|--------------------------------------|------------------------------|-----------------------------|------------------------|---------------------------|----------------------------|-------------------------|
| 1 | RNR-15048 + Vermicompost 10(t/ha) | 10.8 | 105.0 | 6.7 | 16.2 | 4.40 | 6.40 | 39.84 |
| 2 | RNR-15048 + Panchagavya 6% | 10.4 | 111.0 | 7.2 | 14.7 | 3.99 | 5.90 | 40.15 |
| 3 | RNR-15048+Vermicompost(5t/ha) + Panchagavya 3% | 11.8 | 104.7 | 7.4 | 18.3 | 4.94 | 6.93 | 41.55 |
| 4 | BPT-5204 + Vermicompost 10(t/ha) | 10.5 | 110.7 | 6.9 | 15.9 | 4.15 | 6.18 | 40.19 |
| 5 | BPT-5204 + Panchagavya 6% | 10.3 | 101.3 | 7.3 | 15.1 | 3.82 | 5.70 | 40.27 |
| 6 | BPT-5204 + Vermicompost(5t/ha) + Panchagavya 3% | 11.5 | 104.3 | 7.4 | 17.4 | 4.77 | 6.84 | 40.97 |
| 7 | NDR-359 + Vermicompost 10(t/ha) | 12.1 | 105.3 | 6.6 | 21.3 | 5.14 | 7.10 | 37.86 |
| 8 | NDR-359 + Panchagavya 6% | 11.2 | 98.0 | 7.1 | 19.7 | 4.67 | 6.56 | 41.60 |
| 9 | NDR-359 + Vermicompost(5t/ha) + Panchagavya 3% | 12.7 | 114.7 | 7.6 | 21.8 | 5.64 | 7.34 | 41.63 |
| | F test | S | S | S | S | S | S | S |
| | SEm(±) | 0.13 | 1.24 | 0.08 | 0.21 | 0.09 | 0.07 | 0.51 |
| | CD (P=0.05) | 0.39 | 3.70 | 0.24 | 0.63 | 0.26 | 0.20 | 1.23 |

CONCLUSION

Based on the previous findings, it can be concluded that the use of NDR-359 Vermicompost 5 (t/ha) and Panchagavya 3% has enhanced the characteristics of growth and yield of the rice variety NDR-359 while also proving to be profitable. Further studies are required to confirm the findings as they are based on only one season.

REFERENCE

- Anonymous. 2014.** Advances in nutrient dynamics in soil - plant - Atmosphere system for improving nutrient use efficiency. Training manual, IISS, Bhopal. p 70.
- Barik, A. K.; Raj, A. and Saha. R. K. 2008.** Yield performance, economics and soil fertility through organic sources (vermicompost) of nitrogen as substitute to chemical fertilizers in wet season rice. *Crop Research (Hisar)*, 36: 1/3, 4-7.
- Beaulah, A., 2002.** Growth and development of moringa (*Moringa oleifera* Lan.) under organic and inorganic systems of culture. Ph.D. Thesis, Tamil Nadu Agriculture University, Coimbatore.
- Chaudhary, R. S., Anchal Das, and Patnaik, U.S., 2004,** Organic farming for vegetable production using vermicompost and FYM in Kokriguda watershed of Orissa. *Ind. J. Soil Conservation*. 31(2): 203-206.
- Choudhary, K.M., Patel, M.M. and Pagar, R.D. 2014.** Effect of foliar application of panchagavya and leaf extracts of endemic plants on groundnut (*Arachis hypogaea* (L)). *Agricultural Research Communication Centre*, 37(2): 223 –226.
- Dass, A. a.** Irrigation, spacing and cultivar effects on net photosynthetic rate, dry matter partitioning and productivity of rice under system of rice intensification in Mollisol of northern India. *Experimental Agriculture*. 2013; 49(4), 504-523.
- Dominguez, J. (2004).** State of the art and new perspectives on vermicomposting research. In: *Earthworm Ecology*, 2nd ed., ed. C. A. Edwards, 401-424. Boca Raton, FL: CRC Press.
- Emily, A.C.S. 2003.** Standardization of organic production packages for *Withania somnifera* Dunal. M.Sc. Thesis, Tamil Nadu Agricultural University, Coimbatore.
- Gopal Lal Choudhary, Sharma SK, Kendra Pal Singh, Sanju Choudhary, Bazaya BR.** Effect of Panchagavya on Growth and Yield of Organic Black gram [*Vigna mungo* (L.) Hepper]. *International Journal of Current Microbiology and Applied Sciences*. 2017;6(10):1627-1632.
- Jadhav RP, Khafi HR, Raj AD.** Effect of Nitrogen and Vermicompost on Protein Content and Nutrients uptake in Pearl Millet [*Pennisetum glaucum* L.] R. Br. *Emend Stuntz*. *Agricultural Science Digest*. 2011;31(4):319-321.
- Kahani, F. and Hittalmani, S. (2015).** Genetic Analysis and Traits Association in F2 Intervarietal Populations in Rice Under Aerobic Condition. *Rice Research: Open Access Journal*. 3(4): 2375-4338.

- Kamakar (2010).**, Influences of soil physical and chemical properties of vermicompost. *Agricultural Science Digest*. 2011;31(4):319-321.
- Kenchaiah, 1997**, organic farming studies in rice PhD thesis Tamilnadu Agriculture University Coimbatore.
- Khan U, Mishra B, Pachauri P and Kumar Y. 2006.** Effect of integrated nitrogen management on yield and nitrogen nutrition of irrigated rice (*Oryza sativa*). *Indian Journal of Agricultural Sciences* 76(3): 176–80.
- Kumar, B. Gupta, R.K and Bhandari, A.L (2008)** Soil fertility changes after long term application of organic manures and crop residues under rice-wheat System. *Journal of Indian Society of the Soil Science*. 56:80-85.
- Pant, S. R., and Yami, K. D. (2008)** Selective utilization of organic solid wastes by earthworm (*Eisenia foetida*), *Nepal Journal of Science and Technology*. 9: 99-104.
- Selvaraj, N. 2003.** Report on Organic Farming at Horticulture Research Station, Tamil Nadu Agricultural University, Ooty. pp: 2-5.
- Shrimal P, Khan TI.** Studies on the Effects of Vermicompost on Growth Parameters and Chlorophyll Content of Bengal Gram (*Cicer arietinum* L.) var. RSG896. *IOSR Journal of Environmental Science, Toxicology and Food Technology*. 2017;11(5):12-16.
- Singh, S., Singh, V., Khanna, R., Kumar, P., & Shukla, R. (2019).** Effect of nitrogen and sulphur fertilization on growth and yield of rice (*Oryza sativa* L.). *Progressive Research*, 14(1), 526-530.
- Singh RK, Dawson J, Srivastava N.** Effect of sources of nutrient on growth and yield of blackgram (*Vigna mungo* L.) Varieties in NEPZ of India. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(4):1064-1066.
- Somasundaram E and Amanullah M M. 2007.** Panchagavya on growth and productivity of crops: A review. *Green Farming* 1: 22–6.
- Somasundaram E, Sankaran N, Meena S, Thiyagarajan T M, Chandaragiri K and Pannerselvam S. 2003.** Response of greengram to varied levels of panchagavya (organic nutrition) foliar spray. *Madras Agricultural Journal* 90: 169–72.
- Sridhar, K., Srinivas, A., Kumar, A.K., Ramprakesh, T., & Rao, R.P.** Physiological growth parameters of rabi rice (*Oryza sativa* L.) under alternate wetting and drying irrigation with varied nitrogen levels. *International Journal Current Microbiology and Applied Sciences*. 2019;8, 01-15.

Suthar, S. (2007). Nutrients changes and biodynamic of epigeic earthworms *Perionyx excavatus* (Perrier) during recycling of some agricultural wastes, *Biores. Technol.* 98:1608- 1614.

Swaminathan, C., Swaminathan, V. and Vijayalakshmi, K. 2007. Panchagavya - Boon to organic farming, International Book, Distributing Co., Lucknow.

Thanunathan, K., Imayavarambarn, V., Singaravel, R. and Kandasamy, S. 2001. Effect of fly ash on growth, yield and nutrient uptake of sesame. *Sesame and Safflower Newsletter*, 16.

Vennila, C. and Jayanthi, C. 2008. Response of okra to integrated nutrient management. *Journal of Soils and Crops*, 18: 36-40

Vimalendran L, Wahab K. Effect of foliar spray of panchagavya on yield attributes, yield and economic of baby corn. *Journal of Agronomy*. 2013;12(2):109-11.

