

Original Research Article

Performance of tomato genotypes for growth, yield and quality attributes under eastern dry zone of Karnataka, India

Abstract

Field experiments were conducted in 2015-16 at Dept. of Vegetable Science, College of Horticulture, Bengaluru, Karnataka. The main aim was to assess some of growth, earliness, yield and quality attributes of 200 tomato genotypes. These 200 genotypes collected from AVRDC, Taiwan, IIVR, Varanasi, NBPGR, New Delhi and SAU of India and some Research centers of Karnataka. The parameters like Days to first flowering tomato genotype IIHR-2195 took least number of days for anthesis (15.00) after transplanting. Days to 50 per cent flowering minimum (21.00) days after transplanting in Akshaya hence, this superior this genotype useful in heterosis breeding program. In case of growth parameters like plant height, maximum plant height noticed in genotype (13P3) 202.33 cm. The number of branches per plant maximum was observed in genotype (AR-56) i.e. 11.33 branches per plant. Wide variability was observed for fruit width in tomato. The highest fruit width reported (H-86) 8.15cm. Wide variability was observed for tomato fruit length also, the length of tomato fruit maximum noticed in genotype (EC-321426) 7.45 cm. The maximum number of locules per fruit was observed in EC-501580 (13.00) and lowest was observed in ARTD-1 (2.00). Pericarp thickness varied from 7.42 mm (AOTD-10) to 1.05 mm (8). The highest number of fruits per plant (204.37) was recorded in EC-338717 and the least (11.66) was recorded in PKM-1. The highest average fruit weight was observed in H-86 (131.94 g) and the lowest in 73P2 (20.00 g). The genotype AR-56 (7.00) was recorded highest total soluble solids and the genotype ARTD-3 (2.67) recorded lowest total soluble solids. The range for fruit firmness was varied from 0.33 kg/cm² (83P1) to 6.25 kg/cm² (AOTD-18) with the overall mean of 1.33kg/cm². The total tomato fruit yield recorded per plant ranged between 0.39 kg (Roma) and 6.48 kg (4P1). This study gives information on the agronomic variation in the 200 tomato genotypes assessed which can invariably serve as an index to selecting putative parents in breeding for new tomato variety that will combine desirable agronomic characteristics with high yield and its attributing characters.

Keywords: Growth and yield, tomato, genotypes

Introduction

Introduction Tomato (*Solanum Lycopersicon* L.) is one of the most popular and widely grown vegetable in the world ranking second in importance only next to potato and ranked first in preserved and processed vegetables and in the international market there is great demand for tomato [11]. Tomato crop has wider adaptability, high yielding potential and multipurpose uses in fresh as well as processed food industries. Tomatoes are an excellent source of minerals, vitamins [2], antioxidants viz., lycopene and betacarotene which prevent cancer and other heart diseases [8]. Development of tomato genotypes of a promising nature has been important to the vegetable industry throughout the world. New bred varieties have enriched and advanced the agriculture of many countries. Evaluation of germplasm is of immense important in genetic improvement of the crop. For the selection of parents in hybridization, diversity among parents for the character of interest [5]. The production and productivity not only depends on cultural practices and area of cultivation but on high yielding genotypes which have good adaptability to the growing area [3]. Inclusion of genetically diverse parents in any breeding programme is essential to generate new variability and desirable recombinants. Hence, evaluation of tomato genotypes is very essential to see the performance of genotypes for their adaptability and agronomic performance like growth and yield traits to identify the potential genotype. Considering the above facts, the research has been planned with the following objectives to investigate the performance of different tomato genotypes in the Karnataka for its yield and related attributes.

Materials and Methods

Field experiment was conducted at vegetable block, College of Horticulture, UHS Campus, GKVK, Bengaluru. The experimental site is located at an altitude of 930 meters above mean sea level (MSL) and 13° N latitude and 77.370° E longitude in the Eastern Dry Zone of Karnataka (Zone-5). The soil of the experimental area was red sandy loam (Alfisol) with an uniform fertility having soil pH range 6 to 7.3. The material for the present study comprised a total of 200 genotypes which were procured from Indian Institute of Vegetable Research (IIVR),

Varanasi, Uttara Pradesh, Indian Institute of Horticultural Research (IIHR), Hessarghatta, Bengaluru and University of Agricultural Sciences, GKVK, Bengaluru (Table 1 and 2).

The seeds were sown in protrays containing 98 holes. Coir pith was used as growing media. The sown trays were stacked and covered with polythene for three days in order to get early as well as uniform germination. Trays were irrigated daily once or twice depending up on the temperature. After fifteen days of sowing the trays were drenched with 19:19:19 (NPK) at the concentration of 1g/lit in order to get good rooting as well as growth. The prophylactic sprays were taken against pest and diseases.

The field was brought to fine tilth by disc ploughing followed by harrowing and cross cultivation. Farm yard manure at the rate of 25 tonnes per hectare was also incorporated at the time of land preparation. Ridges and furrows were prepared at 60 cm spacing. The half dose of the nitrogen and full dose of phosphorus and potash at the rate of 150:150:150 kg (NPK) per hectare was applied at the time of planting. Twenty five days old seedlings were transplanted in the main field with a spacing of 60 cm between plants, on one side, half way up the ridges. Light irrigation was given at the time of planting. Subsequent irrigations were provided whenever it was required. Just prior to earthing up i.e. 30 days after transplanting, half of nitrogen was given as top dress. Regular weeding was carried out and staking was provided forty five days after transplanting [1].

The experiments were carried out in randomized complete block design (RCBD) with ten plants in each genotype. The experimental field was fairly leveled land with red sandy loam soil of uniform fertility status. The seeds of tomato genotypes were sown in January-15, in the protrays of 98 cells at nursery, department of vegetable science, College of Horticulture, Bengaluru. Four week old seedlings were transplanted in the main field at the spacing of (60x 60cm). All the package of practices recommended by UHS, Bagalkot were followed.

Five plants were selected and labeled at random from each replication in each treatment for recording the following observations and the average from these plants was worked out for the purpose of statistical computation (analysis). The details of observations recorded in each experiment and techniques adopted for the recording the observations were as follows.

Analysis of variance

Analysis of variance was computed for 13 characters. Highly Significant difference was observed among the genotypes for all the characters (Table.3) studied.

Per se performance

Two hundred genotypes collected from different parts of India were evaluated for the growth, yield and quality parameters during summer, season 2015-16. The *per se* performance is presented in the Table 5. The plant height ranged from 43.67 (9P4) to 202.33 cm (13P3) with a population mean of 74.54 cm (Table 4). The genotypic and phenotypic variances were 365.82 and 417.51, respectively. The number of branches per plant ranged from 2.00 (IIHR-2197) to 11.33 (AR-56) with an average of 6.34 branches per plant. The GV and PV were 2.06 and 2.52, respectively. Days to first flowering tomato genotype IIHR-2195 took least number of days for anthesis (15.00) after transplanting, while Sioux and Utkal Local-2 took maximum number of days (37.00). On an average, tomato genotypes took 25.27 days for appearance of first flower after transplanting in the main field.

Days to 50 per cent flowering The days taken for 50 per cent flowering was maximum in Sioux and A. Ahuti (44.00 days after transplanting) and minimum (21.00 days after transplanting) in Akshaya. The tomato genotypes in general took 31.45 days for the days to 50 per cent of the plants to flower after their transplanting in main field. Fruit width (cm) of the tomato shown Wide variability was observed for fruit width in tomato. The width of tomato fruit ranged between 2.50 (62P4) and 8.15 cm (H-86). The average fruit width was 4.41cm. Fruit length (cm) Wide variability was observed for tomato fruit length. The length of tomato fruit ranged between 2.00 cm (73P2) and 7.45 cm (EC-321426). The average fruit length was 4.21 cm. Number of locules per fruit tomato The maximum number of locules per fruit was observed in EC-501580 (13.00) and lowest was observed in ARTD-1 (2.00) with over all mean of 3.74. Pericarp thickness (mm) Pericarp thickness varied from 7.42 mm (AOTD-10) to 1.05 mm (8) with over all mean of 2.96 mm. Number of fruits per plant The highest number of fruits per plant (204.37) was recorded in EC-338717 and the least (11.66) was recorded in PKM-1. On an average, the tomato genotypes had 67.53 fruits per plant. Average fruit weight (g), highest average fruit weight was observed in H-86 (131.94 g) and the lowest in 73P2 (20.00 g) with over all mean of 48.47 g. Yield per plant (kg) The mean total tomato fruit yield recorded per plant was 2.96kg. The quantum of variation for this trait ranged between 0.39 kg (Roma) and 6.48 kg

(4P1). Total soluble solids ($^{\circ}$ Brix) genotype AR-56 (7.00) was recorded highest total soluble solids and the genotype ARTD-3 (2.67) recorded lowest total soluble solids with overall mean of 4.23. Fruit firmness (kg/cm^2) of tomato range for fruit firmness was varied from $0.33 \text{ kg}/\text{cm}^2$ (83P1) to $6.25 \text{ kg}/\text{cm}^2$ (AOTD-18) with the overall mean of $1.33 \text{ kg}/\text{cm}^2$ (Table 4).

Discussion

Vegetable improvement for yield and quality needs a sound knowledge on the genetic architecture of the crop and inheritance of economic characters are of great interest to the breeder. Tomato is one of the important and most demanded vegetable crop. Therefore, improvement of tomato productivity with quality and high nutritional value and resistance to diseases can play significant role in the overall production and nutritional security.

Growth parameters

With respect to the growth parameter the genotype 13P3 recorded highest plant height (202.33 cm) followed by Pant Polyhouse (138.33cm) at final harvesting stage [9], [4] and [6] indicated that, different genotypes recorded the different heights due to each genotype having different potential of growth. With regard to number of branches per plant, the genotype AR-56 (11.33) recorded maximum branches per plant at final harvesting stage and earliness characters like days to first flowering the genotype IIHR-2195 took least number of days (15.00) for anthesis after transplanting and in case of days to 50 per cent of flowering the genotype Akshaya took minimum (21.00) days after transplanting [7], [10], [9], [4] and [6] (Table.4). This indicated that the trait is conditioned by additive gene effects and simple selection would be rewarding in improvement of these earliness characters.

Yield parameters

Among yield and yield related attributes (Table. 4) yield per plant is very important trait as it is dependent character. The highest yield per plant was recorded in genotype 6.48 kg (4P1) followed by H-86 (6.45kg), PTR-6 (5.96kg) because increase in yield per plant in these germplasm was due to higher number of fruits per plant in 4P1 (108.31) followed by H-86

(48.85) and PTR-6 (79.80) [6] [4]. Maximum average fruit weight is observed in H-86 (131.94) [6],[4], [9], [10] and [7].

Quality parameters

Important quality parameters of tomato are TSS, fruit firmness (Table 4). Genotype AR-56 recorded maximum TSS of 7.00 °Brix [6] and [4], maximum fruit firmness was recorded in AOTD-18 (6.25kg/cm²), lowest number of locules per fruit was observed in ARTD-1 (2.00) and highest pericarp thickness was observed in genotype AOTD-10 (7.42mm) [10], [9] and [7].

Conclusion

Considering the mean performance, five superior genotypes for fruit yield may be utilized as parents after multi location, multi seasonal studies. The genotypes with superior quality traits viz for TSS, for firmness content. Hence, the identified superior genotypes should be utilized in further improvement studies through various breeding strategies.

References

- [1] Anonymous, 2019. www.nhb.gov.in. National Horticulture Board, Statistical data.
- [2] Akinfasoye J, Dotun A, Ogunniyan J, Ajayi E. O., (2011). Phenotypic relationship among agronomic characters of commercial tomato (*Lycopersicon esculentum* Mill.) hybrids. *America-Eurasia. J. Agro.*; 4(1):17-22.
- [3] Asiya K.R, Amarananjundeswara. H, Aravinda. J. S, Doddabasappa B, Veere Gowda R. (2017). Performance of garlic (*Allium sativum* L.) genotypes for growth and yield traits under Eastern Dry Zone of Karnataka. *J. Phar. Phyt.*; 12(1):213-216
- [4] Bharathkumar, M. V., (2014), Development of F₁ hybrids with resistance to early blight (*Alternaria solani*) in tomato (*Solanum lycopersicum* L.). *M. Sc. (Hort). Thesis*, Univ. Hortil. Sci., Bagalkot.
- [5] Harrington J. B., (1940), Yielding capacity of wheat crosses as indicated by bulk hybrid tests. *Canadian .J. Res.*; 18:578-584.

- [6] Jyothi, K., (2015). Inheritance of bacterial wilt resistance in tomato (*Solanum lycopersicum* L.), *Ph. D. Hort. Thesis*, Univ. Hort. Sci., Bagalkot.
- [7] Jaiprakashnarayan, R. P., (2007). Genetics of yield attributes and resistance to tomato leaf curl virus (ToLCV) and bacterial wilt in tomato (*Solanum lycopersicum* L.), *Ph. D Thesis*, Univ. Agric. Sci., Bengaluru.
- [8] Kaur C, Walia S, Nagal S, Walia S, Singh J, Singh B.B., (2013). Functional quality and antioxidant composition of selected tomato (*Solanum lycopersicon* L.) cultivars grown in Northern India. *LWT Food Sci. Tech.*; 50:139-145
- [9] Renuka, D. M., (2012) Half diallel analysis in a set of cherry tomato lines for increased growth, yield and quality parameters, *M. Sc. (Hort.) Thesis*, Univ. Hort. Sci., Bagalkot.
- [10] Shalini, M., (2009). Studies on heterosis and combining ability in tomato, (*Solanum lycopersicum* L.), *Ph. D. (Agri.) Thesis*, Univ. Agric. Sci., Dharwad.
- [11] Solieman THI, El-Gabry MAH, Abido AI. (2013). Heterosis, potence ratio and correlation of some important characters in tomato (*Solanum lycopersicum* L.). *Sci. Horti.*, 150:25-30.

Table 1. List of genotypes used in the present study

| Sl. No. | Name of the institute | Number of genotypes collected |
|---------|--|-------------------------------|
| 1 | NBPGR, New Dehli | 10 |
| 2 | IIHR, Hessarghatta, Bengaluru, Karnataka | 06 |
| 3 | IARI, RS, Katrain | 03 |
| 4 | IARI, New Delhi | 06 |
| 5 | IIPR, Kanpur | 02 |
| 6 | IIVR, Varanasi | 08 |
| 7 | PAU, Ludhiana | 09 |
| 8 | MPKV, Rahuri | 03 |
| 9 | TNAU, Coimbatore | 01 |
| 10 | KAU, Thrissur | 01 |
| 11 | UAS, Bengaluru | 01 |
| 12 | UAS, Raichur | 02 |
| 13 | HRS, Devihosur, Haveri, Karnataka | 20 |
| 14 | COH, Mudigere | 05 |
| 15 | COH, Mysuru | 73 |
| 16 | Dept. of Vegetable Science, COH, Bengaluru | 14 |

| | | |
|----|------------------------------|------------|
| 17 | Dept. of BCI, COH, Bengaluru | 27 |
| 18 | Ahmedabad local | 01 |
| 19 | Alahabad local | 03 |
| 20 | Guntur | 01 |
| 21 | Rajamundri | 01 |
| 22 | Madanapalli | 01 |
| 23 | Sakura Seeds Pvt, Ltd | 01 |
| 24 | Sungro Seeds Pvt, Ltd | 01 |
| | Total | 200 |

Table 2. Details of tomato genotypes and checks used in the experiment and their source of collection

| Sl.No. | Original Code | Field Code | Source of Collection |
|--------|---------------|------------|-----------------------------------|
| 1 | ARTD-1 | G-1 | HRS, Devihosur, Haveri, Karnataka |
| 2 | ARTD-2 | G-2 | HRS, Devihosur, Haveri, Karnataka |
| 3 | ARTD-3 | G-3 | HRS, Devihosur, Haveri, Karnataka |
| 4 | ARTD-4 | G-4 | HRS, Devihosur, Haveri, Karnataka |
| 5 | ARTD-5 | G-5 | HRS, Devihosur, Haveri, Karnataka |
| 6 | ARTD-6 | G-6 | HRS, Devihosur, Haveri, Karnataka |
| 7 | ARTD-7 | G-7 | HRS, Devihosur, Haveri, Karnataka |
| 8 | ARTD-8 | G-8 | HRS, Devihosur, Haveri, Karnataka |
| 9 | AOTD-9 | G-9 | HRS, Devihosur, Haveri, Karnataka |
| 10 | AOTD-10 | G-10 | HRS, Devihosur, Haveri, Karnataka |
| 11 | AOTD-11 | G-11 | HRS, Devihosur, Haveri, Karnataka |
| 12 | AOTD-12 | G-12 | HRS, Devihosur, Haveri, Karnataka |
| 13 | AOTD-13 | G-13 | HRS, Devihosur, Haveri, Karnataka |
| 14 | AOTD-14 | G-14 | HRS, Devihosur, Haveri, Karnataka |
| 15 | AOTD-15 | G-15 | HRS, Devihosur, Haveri, Karnataka |
| 16 | AOTD-16 | G-16 | HRS, Devihosur, Haveri, Karnataka |
| 17 | AOTD-17 | G-17 | HRS, Devihosur, Haveri, Karnataka |
| 18 | AOTD-18 | G-18 | HRS, Devihosur, Haveri, Karnataka |
| 19 | AOTD-19 | G-19 | HRS, Devihosur, Haveri, Karnataka |
| 20 | AOTD-20 | G-20 | HRS, Devihosur, Haveri, Karnataka |
| 21 | EC-321426 | G-23 | NBPGR, New Dehli |
| 22 | EC-326146 | G-31 | NBPGR, New Dehli |
| 23 | EC-338714 | G-33 | NBPGR, New Dehli |
| 24 | EC-338717 | G-35 | NBPGR, New Dehli |
| 25 | EC-338725 | G-37 | NBPGR, New Dehli |
| 26 | EC-338735 | G-38 | NBPGR, New Dehli |

| Sl.No. | Original Code | Field Code | Source of Collection |
|--------|-----------------------|------------|--|
| 27 | EC-9057 | G-39 | NBPGR, New Dehli |
| 28 | EC-357839 | G-40 | NBPGR, New Dehli |
| 29 | EC-357846 | G-42 | NBPGR, New Dehli |
| 30 | EC-362940 | G-43 | NBPGR, New Dehli |
| 31 | Punjab Varkha Bahar-1 | G-46 | PAU, Ludhiana |
| 32 | PNR-1 | G-47 | PAU, Ludhiana |
| 33 | S-12 | G-48 | PAU, Ludhiana |
| 34 | Punjab Varkha Bahar-2 | G-49 | PAU, Ludhiana |
| 35 | Hisar Lal | G-50 | PAU, Ludhiana |
| 36 | Punjab Upama | G-51 | PAU, Ludhiana |
| 37 | Punjab Chuhara | G-52 | PAU, Ludhiana |
| 38 | Punjab Kesari | G-53 | PAU, Ludhiana |
| 39 | Punjab Ratta | G-54 | PAU, Ludhiana |
| 40 | 52135 | G-57 | IIPR, Kanpur |
| 41 | Azad-T-1 | G-61 | IIPR, Kanpur |
| 42 | Kashi Sharad | G-62 | IIVR, Varanasi |
| 43 | Kashi Hemanth | G-63 | IIVR, Varanasi |
| 44 | Kashi Amrith | G-64 | IIVR, Varanasi |
| 45 | Kashi Anupama | G-65 | IIVR, Varanasi |
| 46 | H-86 | G-66 | IIVR, Varanasi |
| 47 | Kalyanpur Type | G-68 | IIVR, Varanasi |
| 48 | Arka Abha | G-69 | IIHR, Hessarghatta, Bengaluru, Karnataka |
| 49 | Arka Sourabh | G-70 | IIHR, Hessarghatta, Bengaluru, Karnataka |
| 50 | DT-10 | G-72 | IIVR, Varanasi |
| 51 | H-24 | G-73 | IIVR, Varanasi |
| 52 | PTR-4 | G-75 | UAS, Raichur |
| 53 | PTR-6 | G-76 | UAS, Raichur |
| 54 | Bhagyashree | G-77 | MPKV, Rahuri |
| 55 | Dhanashree | G-78 | MPKV, Rahuri |
| 56 | Pule Raja | G-79 | MPKV, Rahuri |
| 57 | Utkal Kumari | G-80 | Dept. of Vegetable Science, COH, Bengaluru |
| 58 | Utkal Deepti | G-81 | Dept. of Vegetable Science, COH, Bengaluru |
| 59 | Utkal Raja | G-82 | Dept. of Vegetable Science, COH, Bengaluru |
| 60 | Pragnya | G-83 | Dept. of Vegetable Science, COH, Bengaluru |
| 61 | Roma | G-84 | IARI, RS, Katrain |
| 62 | Best of All | G-85 | IARI, RS, Katrain |
| 63 | Sioux | G-86 | IARI, RS, Katrain |
| 64 | PKM-1 | G-87 | TNAU, Coimbatore |
| 65 | Akshaya | G-88 | KAU, Thrissur |
| 66 | Rajamundri Local | G-90 | Rajamundri |
| 67 | Guntur Local | G-91 | Gunturu |
| 68 | Arka Alok | G-92 | IIHR, Hessarghatta, Bengaluru, Karnataka |

| Sl.No. | Original Code | Field Code | Source of Collection |
|--------|-------------------|------------|--|
| 69 | Arka Vikas | G-93 | IIHR, Hessarghatta, Bengaluru, Karnataka |
| 70 | Arka Meghali | G-95 | IIHR, Hessarghatta, Bengaluru, Karnataka |
| 71 | Arka Ahuthi | G-96 | IIHR, Hessarghatta, Bengaluru, Karnataka |
| 72 | Madanapalli Local | G-98 | Madanapalli |
| 73 | Pusa Sheetal | G-99 | IARI, New Delhi |
| 74 | Pusa Sadabahar | G-100 | IARI, New Delhi |
| 75 | Pusa Rohini | G-101 | IARI, New Delhi |
| 76 | Pusa Gourava | G-102 | IARI, New Delhi |
| 77 | P-120 | G-103 | IARI, New Delhi |
| 78 | Pusa Ruby | G-104 | IARI, New Delhi |
| 79 | S-22 | G-105 | Sakura Seeds PVT, LTD |
| 80 | Navodaya | G-106 | Sungro Seeds PVT, LTD |
| 81 | Selection | G-107 | Ahmedabad local |
| 82 | C-10-2 | G-108 | Dept. of BCI, COH, Bengaluru |
| 83 | C-11-2 | G-109 | Dept. of BCI, COH, Bengaluru |
| 84 | C-20-1 | G-110 | Dept. of BCI, COH, Bengaluru |
| 85 | CO-3 | G-111 | Dept. of BCI, COH, Bengaluru |
| 86 | CLN-2026 | G-112 | Dept. of BCI, COH, Bengaluru |
| 87 | 41 | G-113 | Dept. of BCI, COH, Bengaluru |
| 88 | DVRT-2 | G-114 | Dept. of BCI, COH, Bengaluru |
| 89 | EC-13904 | G-116 | Dept. of BCI, COH, Bengaluru |
| 90 | C-1-4 | G-119 | Dept. of BCI, COH, Bengaluru |
| 91 | C-4-1 | G-120 | Dept. of BCI, COH, Bengaluru |
| 92 | EC-381263 | G-121 | Dept. of BCI, COH, Bengaluru |
| 93 | EC-501574 | G-123 | Dept. of BCI, COH, Bengaluru |
| 94 | EC-501580 | G-124 | Dept. of BCI, COH, Bengaluru |
| 95 | EC-501583 | G-126 | Dept. of BCI, COH, Bengaluru |
| 96 | EC-538404 | G-129 | Dept. of BCI, COH, Bengaluru |
| 97 | EC-538405 | G-130 | Dept. of BCI, COH, Bengaluru |
| 98 | EC-620383 | G-133 | Dept. of BCI, COH, Bengaluru |
| 99 | EC-620398 | G-134 | Dept. of BCI, COH, Bengaluru |
| 100 | EC-620401 | G-135 | Dept. of BCI, COH, Bengaluru |
| 101 | EC-620446 | G-136 | Dept. of BCI, COH, Bengaluru |
| 102 | EC-620464 | G-137 | Dept. of BCI, COH, Bengaluru |
| 103 | EC-620469 | G-138 | Dept. of BCI, COH, Bengaluru |
| 104 | EC-620470 | G-139 | Dept. of BCI, COH, Bengaluru |
| 105 | Monte Fevarate | G-140 | Dept. of BCI, COH, Bengaluru |
| 106 | Rio Grande | G-141 | Dept. of BCI, COH, Bengaluru |
| 107 | Angoorlata | G-143 | Dept. of BCI, COH, Bengaluru |
| 108 | Ageta-2 | G-144 | Dept. of BCI, COH, Bengaluru |
| 109 | 85 | G-145 | COH, Mysuru |
| 110 | 4 | G-148 | COH, Mysuru |

| Sl.No. | Original Code | Field Code | Source of Collection |
|--------|---------------|------------|----------------------|
| 111 | 80 | G-149 | COH, Mysuru |
| 112 | 200 | G-150 | COH, Mysuru |
| 113 | 8 | G-151 | COH, Mysuru |
| 114 | 1P2 | G-152 | COH, Mysuru |
| 115 | 2P2 | G-153 | COH, Mysuru |
| 116 | 3P2 | G-154 | COH, Mysuru |
| 117 | 4P1 | G-155 | COH, Mysuru |
| 118 | 5T5P6 | G-156 | COH, Mysuru |
| 119 | 6T6P8 | G-157 | COH, Mysuru |
| 120 | 7 | G-158 | COH, Mysuru |
| 121 | 8P3 | G-159 | COH, Mysuru |
| 122 | 9P4 | G-160 | COH, Mysuru |
| 123 | 10P6 | G-161 | COH, Mysuru |
| 124 | 11P4 | G-162 | COH, Mysuru |
| 125 | 12P1 | G-163 | COH, Mysuru |
| 126 | 13P3 | G-164 | COH, Mysuru |
| 127 | 14P6 | G-165 | COH, Mysuru |
| 128 | 15P4 | G-166 | COH, Mysuru |
| 129 | 16P2 | G-167 | COH, Mysuru |
| 130 | 17P5 | G-168 | COH, Mysuru |
| 131 | 18P3 | G-169 | COH, Mysuru |
| 132 | 19P8 | G-170 | COH, Mysuru |
| 133 | 20 | G-171 | COH, Mysuru |
| 134 | 21 | G-172 | COH, Mysuru |
| 135 | 23P4 | G-173 | COH, Mysuru |
| 136 | 25P2 | G-175 | COH, Mysuru |
| 137 | 27P2 | G-177 | COH, Mysuru |
| 138 | 28P2 | G-178 | COH, Mysuru |
| 139 | 29P4 | G-179 | COH, Mysuru |
| 140 | 30P2 | G-180 | COH, Mysuru |
| 141 | 33P2 | G-183 | COH, Mysuru |
| 142 | 34P2 | G-184 | COH, Mysuru |
| 143 | 35P2 | G-185 | COH, Mysuru |
| 144 | 36 | G-186 | COH, Mysuru |
| 145 | 37P2 | G-187 | COH, Mysuru |
| 146 | 38P2 | G-188 | COH, Mysuru |
| 147 | 40P4 | G-190 | COH, Mysuru |
| 148 | 43 | G-193 | COH, Mysuru |
| 149 | 44P2 | G-194 | COH, Mysuru |
| 150 | 45 | G-195 | COH, Mysuru |
| 151 | 46P5 | G-196 | COH, Mysuru |
| 152 | 47P2 | G-197 | COH, Mysuru |

| Sl.No. | Original Code | Field Code | Source of Collection |
|--------|------------------|------------|--|
| 153 | 48P4 | G-198 | COH, Mysuru |
| 154 | 51 | G-201 | COH, Mysuru |
| 155 | 53 | G-202 | COH, Mysuru |
| 156 | 54P3 | G-203 | COH, Mysuru |
| 157 | 55P2 | G-204 | COH, Mysuru |
| 158 | 56P2 | G-205 | COH, Mysuru |
| 159 | 58 | G-206 | COH, Mysuru |
| 160 | 59 | G-207 | COH, Mysuru |
| 161 | 61P4 | G-208 | COH, Mysuru |
| 162 | 62P4 | G-209 | COH, Mysuru |
| 163 | 63P3 | G-210 | COH, Mysuru |
| 164 | 64 | G-211 | COH, Mysuru |
| 165 | 65P5 | G-212 | COH, Mysuru |
| 166 | 66P1 | G-213 | COH, Mysuru |
| 167 | 70 | G-214 | COH, Mysuru |
| 168 | 71P2 | G-215 | COH, Mysuru |
| 169 | 72P2 | G-216 | COH, Mysuru |
| 170 | 73P2 | G-217 | COH, Mysuru |
| 171 | 74P5 | G-218 | COH, Mysuru |
| 172 | 75P3 | G-219 | COH, Mysuru |
| 173 | 76P1 | G-220 | COH, Mysuru |
| 174 | 77P1 | G-221 | COH, Mysuru |
| 175 | 78P4 | G-222 | COH, Mysuru |
| 176 | 83P1 | G-223 | COH, Mysuru |
| 177 | 84P1 | G-224 | COH, Mysuru |
| 178 | 86P2 | G-225 | COH, Mysuru |
| 179 | 67 | G-226 | COH, Mysuru |
| 180 | 68 | G-227 | COH, Mysuru |
| 181 | Nandhi | G-228 | UAS, Bengaluru |
| 182 | Pant Polyhouse-2 | G-229 | Alahabad local |
| 183 | AR-28 | G-230 | COH, Mudigere |
| 184 | Anaga | G-231 | Dept. of Vegetable Science, COH, Bengaluru |
| 185 | IIHR -2195 | G-233 | Dept. of Vegetable Science, COH, Bengaluru |
| 186 | AR-21 | G-234 | COH, Mudigere |
| 187 | AR-56 | G-235 | COH, Mudigere |
| 188 | AR-4 | G-237 | COH, Mudigere |
| 189 | IIHR – 2199 | G-239 | Dept. of Vegetable Science, COH, Bengaluru |
| 190 | IIHR – 2198 | G-240 | Dept. of Vegetable Science, COH, Bengaluru |
| 191 | IIHR – 2197 | G-242 | Dept. of Vegetable Science, COH, Bengaluru |
| 192 | AR-29 | G-244 | COH, Mudigere |
| 193 | IIHR – 2196 | G-246 | Dept. of Vegetable Science, COH, Bengaluru |
| 194 | Utkal Local-2 | G-247 | Dept. of Vegetable Science, COH, Bengaluru |

| Sl.No. | Original Code | Field Code | Source of Collection |
|---------------|----------------|------------|--|
| 195 | Pant Polyhouse | G-248 | Alahabad local |
| 196 | Pant-3 | G-249 | Alahabad local |
| 197 | IIHR – 2200 | G-250 | Dept. of Vegetable Science, COH, Bengaluru |
| 198 | IIHR - 2201 | G-251 | Dept. of Vegetable Science, COH, Bengaluru |
| 199 | H-24-1 | G-252 | Dept. of Vegetable Science, COH, Bengaluru |
| 200 | Solan-2 | G-254 | COH, Mysuru |
| Checks | | | |
| 1 | Arka Samrat | Check-1 | IIHR, Hessarghatta, Bengaluru, Karnataka |
| 2 | Arka Rakshak | Check-2 | IIHR, Hessarghatta, Bengaluru, Karnataka |
| 3 | 5105 | Check-3 | IIPR, Kanpur |

Table 3. Analysis of variance (ANOVA) for growth, yield and quality parameters in tomato

| Sl. No. | Source of variation | Mean sum of squares | | | | | |
|----------|--------------------------------|---------------------|------------|------------|---------------|-------------------------|--------|
| | | Blocks | Entries | (a) Checks | (b) Varieties | (c) Checks Vs Varieties | Error |
| | Degrees of freedom | 10 | 200 | 3 | 203 | 1 | 18 |
| A | Growth parameters | | | | | | |
| 1 | Plant height (cm) | 164.87 | 456.12 ** | 2502.06 ** | 487.40 ** | 2683.26** | 51.68 |
| 2 | Number of branches/plant | 1.35 | 2.74 ** | 7.34 ** | 2.78** | 2.58* | 0.46 |
| 3 | Days to first flowering | 4.53 | 20.63* | 8.53* | 20.49** | 17.04* | 1.64 |
| 4 | Days to 50 per cent flowering | 3.77 | 20.37** | 14.53** | 20.67** | 92.52** | 1.53 |
| B | Yield parameters | | | | | | |
| 5 | Fruit width (cm) | 0.46 | 0.87 ** | 10.96 ** | 0.98** | 2.89** | 0.14 |
| 6 | Fruit length (cm) | 0.59 | 0.97 ** | 7.99 ** | 1.05 ** | 1.96** | 0.20 |
| 7 | Number of locules per fruit | 0.33 | 2.16 ** | 2.03 ** | 2.22** | 15.60** | 0.10 |
| 8 | Pericarp thickness (mm) | 0.93 | 1.30 ** | 2.19 * | 1.32** | 1.87* | 0.36 |
| 9 | Fruits per plant | 313.58 | 1160.81 ** | 931.26 ** | 1155.28** | 502.04* | 106.79 |
| 10 | Average fruit weight (g) | 172.95 | 312.47 ** | 6337.77 ** | 372.16** | 319.86* | 62.41 |
| 11 | Yield per plant (kg) | 0.81 | 1.45 ** | 13.01 ** | 1.57** | 1.58* | 0.31 |
| C | Quality parameters | | | | | | |
| 12 | TSS (⁰ B) | 0.341 | 0.601 ** | 5.410 ** | 0.650 ** | 0.990 ** | 0.110 |
| 13 | Firmness (kg/cm ²) | 0.007 | 0.519 ** | 19.004 ** | 0.932 ** | 46.853 ** | 0.003 |

*Significant at 5 per cent, ** Significant at 1 per cent

Table 4. *Per se* performance of 200 tomato genotypes for growth, yield and quality attributes

| Sl. No. | Original code | Genotypes | Plant height (cm) | Number of branches/plant | Days to first flowering | Days to 50 per cent flowering | Fruit width (cm) | Fruit length (cm) | Number of locules per fruit | Pericarp thickness (mm) |
|---------|-----------------------|-----------|-------------------|--------------------------|-------------------------|-------------------------------|------------------|-------------------|-----------------------------|-------------------------|
| 1 | ARTD-1 | G-1 | 60.33 | 6.67 | 23.00 | 29.00 | 4.90 | 5.95 | 2.00 | 2.61 |
| 2 | ARTD-2 | G-2 | 65.67 | 4.00 | 24.00 | 30.00 | 4.95 | 6.50 | 2.00 | 4.56 |
| 3 | ARTD-3 | G-3 | 56.00 | 5.67 | 24.00 | 28.00 | 5.10 | 4.70 | 3.00 | 3.63 |
| 4 | ARTD-4 | G-4 | 53.00 | 5.67 | 23.00 | 31.00 | 5.55 | 5.20 | 3.00 | 4.17 |
| 5 | ARTD-5 | G-5 | 62.67 | 5.00 | 23.00 | 30.00 | 4.80 | 6.45 | 3.00 | 2.45 |
| 6 | ARTD-6 | G-6 | 63.67 | 3.67 | 23.00 | 31.00 | 4.50 | 4.60 | 2.00 | 4.49 |
| 7 | ARTD-7 | G-7 | 51.33 | 4.33 | 24.00 | 28.00 | 4.25 | 4.60 | 2.00 | 1.41 |
| 8 | ARTD-8 | G-8 | 48.33 | 4.00 | 22.00 | 27.00 | 5.10 | 6.45 | 2.00 | 4.80 |
| 9 | AOTD-9 | G-9 | 61.67 | 4.67 | 23.00 | 28.00 | 4.70 | 6.50 | 3.00 | 5.03 |
| 10 | AOTD-10 | G-10 | 58.33 | 4.33 | 24.00 | 30.00 | 4.95 | 6.45 | 2.00 | 7.42 |
| 11 | AOTD-11 | G-11 | 70.33 | 7.67 | 24.00 | 29.00 | 4.75 | 5.80 | 3.00 | 4.36 |
| 12 | AOTD-12 | G-12 | 59.67 | 6.00 | 25.00 | 30.00 | 4.50 | 4.45 | 3.00 | 2.81 |
| 13 | AOTD-13 | G-13 | 52.00 | 7.33 | 19.00 | 26.00 | 5.20 | 4.85 | 2.00 | 2.29 |
| 14 | AOTD-14 | G-14 | 45.67 | 7.33 | 22.00 | 27.00 | 4.55 | 4.55 | 3.00 | 2.72 |
| 15 | AOTD-15 | G-15 | 53.00 | 6.00 | 24.00 | 31.00 | 5.05 | 4.80 | 3.00 | 3.24 |
| 16 | AOTD-16 | G-16 | 84.00 | 4.67 | 20.00 | 27.00 | 4.90 | 4.70 | 4.00 | 3.25 |
| 17 | AOTD-17 | G-17 | 52.00 | 6.33 | 21.00 | 26.00 | 5.20 | 5.05 | 3.00 | 2.78 |
| 18 | AOTD-18 | G-18 | 47.00 | 5.00 | 18.00 | 31.00 | 4.80 | 4.75 | 3.00 | 4.02 |
| 19 | AOTD-19 | G-19 | 58.00 | 5.67 | 27.00 | 31.00 | 4.70 | 5.05 | 2.00 | 5.60 |
| 20 | AOTD-20 | G-20 | 56.33 | 5.67 | 26.00 | 31.00 | 4.45 | 4.60 | 3.00 | 5.92 |
| 21 | EC-321426 | G-23 | 55.00 | 6.00 | 26.00 | 31.00 | 7.45 | 7.45 | 4.00 | 6.11 |
| 22 | EC-326146 | G-31 | 78.67 | 6.33 | 21.00 | 26.00 | 4.00 | 5.05 | 2.00 | 2.61 |
| 23 | EC-338714 | G-33 | 66.33 | 4.67 | 23.00 | 27.00 | 4.25 | 3.75 | 3.00 | 3.15 |
| 24 | EC-338717 | G-35 | 110.00 | 7.33 | 19.00 | 35.00 | 3.30 | 3.15 | 2.00 | 2.81 |
| 25 | EC-338725 | G-37 | 96.00 | 6.33 | 33.00 | 32.00 | 3.85 | 3.55 | 3.00 | 6.05 |
| 26 | EC-338735 | G-38 | 58.33 | 5.00 | 27.00 | 33.00 | 4.90 | 3.90 | 7.00 | 1.31 |
| 27 | EC-339057 | G-39 | 66.67 | 5.00 | 17.00 | 23.00 | 3.35 | 3.25 | 2.00 | 1.71 |
| 28 | EC-357839 | G-40 | 83.67 | 6.00 | 15.00 | 22.00 | 4.65 | 4.50 | 5.00 | 2.20 |
| 29 | EC-357845 | G-41 | 130.00 | 6.67 | 16.00 | 22.00 | 4.75 | 3.95 | 3.00 | 4.77 |
| 30 | EC-362940 | G-43 | 78.33 | 9.00 | 18.00 | 23.00 | 4.70 | 3.00 | 6.00 | 3.94 |
| 31 | Punjab Varkha Bahar-1 | G-46 | 93.33 | 5.00 | 27.00 | 33.00 | 5.75 | 5.80 | 4.00 | 2.80 |

| Sl. No. | Original code | Genotypes | Plant height (cm) | Number of branches/plant | Days to first flowering | Days to 50 per cent flowering | Fruit width (cm) | Fruit length (cm) | Number of locules per fruit | Pericarp thickness (mm) |
|---------|-----------------------|-----------|-------------------|--------------------------|-------------------------|-------------------------------|------------------|-------------------|-----------------------------|-------------------------|
| 32 | PNR-1 | G-47 | 102.67 | 6.67 | 24.00 | 33.00 | 5.55 | 4.90 | 3.00 | 2.96 |
| 33 | S-12 | G-48 | 63.67 | 4.67 | 24.00 | 27.00 | 4.10 | 3.60 | 5.00 | 3.19 |
| 34 | Punjab Varkha Bahar-2 | G-49 | 56.67 | 5.67 | 23.00 | 29.00 | 5.70 | 4.55 | 4.00 | 3.16 |
| 35 | Hisar Lal | G-50 | 61.00 | 5.67 | 23.00 | 29.00 | 4.15 | 6.50 | 3.00 | 3.18 |
| 36 | Punjab Upama | G-51 | 73.67 | 4.67 | 22.00 | 27.00 | 4.60 | 5.45 | 3.00 | 2.45 |
| 37 | Punjab Chhuhar | G-52 | 75.00 | 6.33 | 24.00 | 28.00 | 4.20 | 6.50 | 3.00 | 3.40 |
| 38 | Punjab Kesari | G-53 | 71.67 | 3.67 | 23.00 | 27.00 | 4.90 | 4.35 | 4.00 | 2.03 |
| 39 | Punjab Ratta | G-54 | 69.33 | 6.67 | 27.00 | 33.00 | 5.35 | 6.20 | 2.00 | 4.48 |
| 40 | 52135 | G-57 | 98.00 | 6.67 | 22.00 | 27.00 | 4.45 | 3.80 | 3.00 | 2.55 |
| 41 | Azad-T-1 | G-61 | 80.33 | 6.67 | 31.00 | 37.00 | 4.30 | 3.80 | 5.00 | 2.55 |
| 42 | Kashi Sharad | G-62 | 96.00 | 6.33 | 24.00 | 31.00 | 5.35 | 5.45 | 5.00 | 6.74 |
| 43 | Kashi Hemanth | G-63 | 102.33 | 7.67 | 24.00 | 32.00 | 6.35 | 6.70 | 4.00 | 3.49 |
| 44 | Kashi Amrith | G-64 | 59.00 | 8.33 | 33.00 | 37.00 | 5.50 | 4.85 | 3.00 | 2.71 |
| 45 | Kashi Anupama | G-65 | 51.00 | 6.33 | 29.00 | 35.00 | 6.60 | 4.45 | 4.00 | 2.50 |
| 46 | H-86 | G-66 | 62.00 | 6.00 | 29.00 | 32.00 | 8.15 | 5.85 | 5.00 | 1.49 |
| 47 | Kalyanpur Type | G-68 | 109.33 | 7.67 | 27.00 | 31.00 | 7.20 | 5.05 | 3.00 | 4.05 |
| 48 | A Abha | G-69 | 72.67 | 5.67 | 27.00 | 33.00 | 5.60 | 4.35 | 5.00 | 2.23 |
| 49 | A Saurabh | G-70 | 78.67 | 7.00 | 27.00 | 34.00 | 3.25 | 3.10 | 5.00 | 2.91 |
| 50 | DT-10 | G-72 | 94.33 | 9.00 | 24.00 | 30.00 | 6.00 | 4.60 | 3.00 | 3.18 |
| 51 | H-24 | G-73 | 61.00 | 9.00 | 27.00 | 33.00 | 3.85 | 3.75 | 4.00 | 2.03 |
| 52 | PTR-4 | G-75 | 64.33 | 4.67 | 15.00 | 23.00 | 4.20 | 4.15 | 4.00 | 1.13 |
| 53 | PTR-6 | G-76 | 68.00 | 5.00 | 26.00 | 30.00 | 5.10 | 4.75 | 3.00 | 5.72 |
| 54 | Bhagyashree | G-77 | 58.33 | 6.00 | 22.00 | 28.00 | 6.55 | 5.60 | 6.00 | 3.69 |
| 55 | Dhanashree | G-78 | 72.33 | 6.33 | 27.00 | 30.00 | 4.85 | 4.25 | 3.00 | 2.40 |
| 56 | Pule Raja | G-79 | 105.67 | 8.00 | 25.00 | 29.00 | 4.95 | 5.40 | 3.00 | 4.10 |
| 57 | Utkal Kumari | G-80 | 61.67 | 5.33 | 25.00 | 33.00 | 5.20 | 3.95 | 5.00 | 2.97 |
| 58 | Utkal Deepti | G-81 | 64.00 | 6.33 | 25.00 | 31.00 | 3.45 | 3.65 | 3.00 | 2.30 |
| 59 | Utkal Raja | G-82 | 84.00 | 6.33 | 16.00 | 22.00 | 5.25 | 4.30 | 4.00 | 2.75 |
| 60 | Pragnya | G-83 | 63.67 | 6.00 | 25.00 | 33.00 | 4.40 | 4.05 | 3.00 | 2.75 |
| 61 | Roma | G-84 | 68.67 | 5.67 | 27.00 | 42.00 | 4.10 | 6.65 | 2.00 | 3.35 |
| 62 | Best of all | G-85 | 105.67 | 5.67 | 28.00 | 35.00 | 3.45 | 3.60 | 4.00 | 1.50 |
| 63 | Sioux | G-86 | 108.33 | 6.33 | 37.00 | 44.00 | 4.85 | 4.15 | 4.00 | 2.60 |

| Sl. No. | Original code | Genotypes | Plant height (cm) | Number of branches/plant | Days to first flowering | Days to 50 per cent flowering | Fruit width (cm) | Fruit length (cm) | Number of locules per fruit | Pericarp thickness (mm) |
|---------|----------------|-----------|-------------------|--------------------------|-------------------------|-------------------------------|------------------|-------------------|-----------------------------|-------------------------|
| 64 | PKM-1 | G-87 | 45.33 | 6.33 | 28.00 | 37.00 | 4.60 | 3.65 | 5.00 | 2.70 |
| 65 | Akshaya | G-88 | 49.33 | 4.00 | 15.00 | 21.00 | 5.60 | 3.95 | 3.00 | 3.15 |
| 66 | Rajamundri | G-90 | 69.67 | 6.33 | 28.00 | 34.00 | 4.70 | 3.75 | 4.00 | 3.84 |
| 67 | Guntur Local | G-91 | 66.00 | 5.33 | 25.00 | 34.00 | 5.10 | 3.70 | 5.00 | 2.00 |
| 68 | A Alok | G-92 | 48.33 | 6.00 | 27.00 | 34.00 | 5.85 | 5.35 | 3.00 | 2.00 |
| 69 | A Vikash | G-93 | 56.33 | 5.00 | 27.00 | 33.00 | 5.05 | 3.85 | 4.00 | 2.80 |
| 70 | A Meghali | G-95 | 78.67 | 5.33 | 23.00 | 33.00 | 4.70 | 4.10 | 3.00 | 2.25 |
| 71 | A Ahuthi | G-96 | 68.67 | 4.67 | 33.00 | 44.00 | 4.00 | 6.50 | 2.00 | 2.05 |
| 72 | Madanapalli | G-98 | 90.00 | 5.67 | 22.00 | 32.00 | 6.30 | 4.70 | 4.00 | 2.70 |
| 73 | Pusa Sheetal | G-99 | 73.00 | 4.00 | 29.00 | 36.00 | 5.05 | 5.00 | 3.00 | 3.44 |
| 74 | Pusa Sadabahar | G-100 | 56.67 | 3.33 | 25.00 | 33.00 | 3.80 | 5.00 | 5.00 | 3.14 |
| 75 | Pusa Rohini | G-101 | 77.00 | 4.33 | 24.00 | 32.00 | 5.80 | 5.25 | 3.00 | 3.37 |
| 76 | Pusa Gourava | G-102 | 60.33 | 5.00 | 17.00 | 29.00 | 4.70 | 5.55 | 2.00 | 4.50 |
| 77 | P-120 | G-103 | 61.67 | 3.00 | 23.00 | 28.00 | 5.00 | 4.35 | 5.00 | 2.37 |
| 78 | Pusa Ruby | G-104 | 70.00 | 5.67 | 23.00 | 27.00 | 4.50 | 3.60 | 7.00 | 2.25 |
| 79 | S-22 | G-105 | 59.67 | 4.33 | 24.00 | 31.00 | 5.50 | 4.45 | 9.00 | 4.20 |
| 80 | Navodaya | G-106 | 57.67 | 4.67 | 29.00 | 42.00 | 4.50 | 3.80 | 3.00 | 1.65 |
| 81 | Selection | G-107 | 74.00 | 7.33 | 15.00 | 32.00 | 4.80 | 3.85 | 4.00 | 2.50 |
| 82 | C-10-2 | G-108 | 73.00 | 5.00 | 18.00 | 28.00 | 4.95 | 3.60 | 4.00 | 2.82 |
| 83 | C-11-2 | G-109 | 54.00 | 5.00 | 26.00 | 33.00 | 3.85 | 3.20 | 5.00 | 2.71 |
| 84 | C-20-1 | G-110 | 70.67 | 4.67 | 26.00 | 32.00 | 5.95 | 5.35 | 5.00 | 4.75 |
| 85 | CO-3 | G-111 | 49.67 | 5.00 | 28.00 | 33.00 | 5.00 | 4.05 | 4.00 | 2.58 |
| 86 | CLN-2026 | G-112 | 63.67 | 5.33 | 25.00 | 29.00 | 3.90 | 4.80 | 3.00 | 2.10 |
| 87 | 41 | G-113 | 75.33 | 2.67 | 32.00 | 37.00 | 4.95 | 5.70 | 3.00 | 3.25 |
| 88 | DVRT-2 | G-114 | 71.67 | 4.33 | 27.00 | 33.00 | 4.85 | 3.75 | 6.00 | 1.50 |
| 89 | EC-13904 | G-116 | 86.67 | 6.00 | 18.00 | 25.00 | 4.35 | 3.65 | 4.00 | 2.34 |
| 90 | C-1-4 | G-119 | 60.33 | 8.33 | 32.00 | 44.00 | 4.95 | 4.00 | 4.00 | 2.80 |
| 91 | C-4-1 | G-120 | 110.67 | 7.67 | 28.00 | 33.00 | 3.10 | 3.40 | 3.00 | 2.80 |
| 92 | EC-381263 | G-121 | 114.33 | 7.67 | 27.00 | 33.00 | 3.20 | 4.00 | 3.00 | 3.00 |
| 93 | EC-501574 | G-123 | 101.67 | 6.67 | 19.00 | 27.00 | 5.80 | 4.30 | 5.00 | 2.12 |
| 94 | EC-501580 | G-124 | 74.67 | 9.00 | 27.00 | 33.00 | 5.15 | 4.40 | 13.00 | 4.11 |
| 95 | EC-501583 | G-126 | 81.67 | 10.00 | 33.00 | 39.00 | 3.90 | 3.10 | 4.00 | 2.13 |

| Sl. No. | Original code | Genotypes | Plant height (cm) | Number of branches/plant | Days to first flowering | Days to 50 per cent flowering | Fruit width (cm) | Fruit length (cm) | Number of locules per fruit | Pericarp thickness (mm) |
|---------|----------------|-----------|-------------------|--------------------------|-------------------------|-------------------------------|------------------|-------------------|-----------------------------|-------------------------|
| 96 | EC-538404 | G-129 | 65.00 | 4.67 | 26.00 | 32.00 | 6.15 | 4.80 | 8.00 | 1.90 |
| 97 | EC-538405 | G-130 | 61.67 | 5.00 | 28.00 | 32.00 | 5.50 | 5.35 | 5.00 | 4.16 |
| 98 | EC-620383 | G-133 | 77.67 | 7.00 | 25.00 | 29.00 | 3.20 | 3.50 | 2.00 | 3.38 |
| 99 | EC-620398 | G-134 | 50.67 | 4.33 | 24.00 | 30.00 | 4.05 | 5.35 | 2.00 | 3.33 |
| 100 | EC-620401 | G-135 | 66.33 | 6.00 | 25.00 | 31.00 | 5.20 | 5.20 | 3.00 | 1.99 |
| 101 | EC-620446 | G-136 | 54.00 | 3.67 | 23.00 | 28.00 | 5.05 | 5.80 | 3.00 | 3.00 |
| 102 | EC-620464 | G-137 | 56.67 | 5.33 | 27.00 | 37.00 | 4.85 | 5.00 | 2.00 | 2.55 |
| 103 | EC-620469 | G-138 | 57.00 | 5.33 | 24.00 | 28.00 | 4.40 | 5.80 | 3.00 | 4.11 |
| 104 | EC-620470 | G-139 | 59.33 | 4.67 | 27.00 | 33.00 | 4.10 | 4.25 | 4.00 | 1.50 |
| 105 | Monte Fevarate | G-140 | 55.33 | 4.67 | 26.00 | 32.00 | 4.15 | 4.90 | 2.00 | 2.67 |
| 106 | Rio Grande | G-141 | 67.00 | 7.33 | 33.00 | 37.00 | 3.30 | 3.45 | 6.00 | 2.26 |
| 107 | Angoorlata | G-143 | 76.67 | 4.33 | 25.00 | 33.00 | 3.95 | 3.70 | 3.00 | 2.61 |
| 108 | Ageta-2 | G-144 | 59.33 | 5.00 | 22.00 | 26.00 | 5.50 | 5.35 | 3.00 | 3.00 |
| 109 | 85 | G-145 | 96.00 | 7.00 | 27.00 | 32.00 | 4.20 | 4.10 | 3.00 | 2.50 |
| 110 | 4 | G-148 | 99.00 | 8.67 | 15.00 | 24.00 | 4.15 | 4.20 | 3.00 | 2.30 |
| 111 | 80 | G-149 | 54.67 | 5.67 | 28.00 | 33.00 | 4.95 | 3.25 | 6.00 | 1.56 |
| 112 | 200 | G-150 | 69.67 | 5.33 | 33.00 | 37.00 | 4.30 | 5.05 | 2.00 | 5.40 |
| 113 | 8 | G-151 | 57.33 | 7.33 | 29.00 | 33.00 | 5.05 | 4.80 | 4.00 | 1.05 |
| 114 | 1P2 | G-152 | 76.67 | 9.67 | 29.00 | 33.00 | 4.55 | 3.20 | 4.00 | 2.06 |
| 115 | 2P2 | G-153 | 95.67 | 7.00 | 24.00 | 29.00 | 4.75 | 4.40 | 5.00 | 1.62 |
| 116 | 3P2 | G-154 | 86.33 | 9.33 | 28.00 | 33.00 | 3.45 | 3.05 | 4.00 | 2.71 |
| 117 | 4P1 | G-155 | 85.33 | 9.00 | 28.00 | 33.00 | 5.80 | 4.65 | 4.00 | 2.16 |
| 118 | 5T5P6 | G-156 | 71.67 | 8.67 | 28.00 | 33.00 | 3.35 | 2.80 | 6.00 | 3.86 |
| 119 | 6T6P8 | G-157 | 69.33 | 7.00 | 24.00 | 28.00 | 4.05 | 4.50 | 3.00 | 2.18 |
| 120 | 7 | G-158 | 58.67 | 10.00 | 29.00 | 33.00 | 4.40 | 3.95 | 4.00 | 2.60 |
| 121 | 8P3 | G-159 | 49.67 | 6.33 | 18.00 | 33.00 | 4.75 | 3.40 | 5.00 | 3.70 |
| 122 | 9P4 | G-160 | 43.67 | 6.00 | 29.00 | 33.00 | 5.00 | 4.85 | 3.00 | 2.41 |
| 123 | 10P6 | G-161 | 82.00 | 7.00 | 17.00 | 23.00 | 3.60 | 3.45 | 3.00 | 3.17 |
| 124 | 11P4 | G-162 | 122.33 | 9.00 | 29.00 | 33.00 | 4.25 | 3.45 | 4.00 | 4.60 |
| 125 | 12P1 | G-163 | 77.67 | 6.00 | 30.00 | 37.00 | 4.00 | 3.50 | 3.00 | 5.56 |
| 126 | 13P3 | G-164 | 202.33 | 6.00 | 31.00 | 36.00 | 2.55 | 3.40 | 3.00 | 2.60 |
| 127 | 14P6 | G-165 | 60.00 | 6.67 | 28.00 | 33.00 | 3.45 | 3.20 | 2.00 | 3.10 |

| Sl. No. | Original code | Genotypes | Plant height (cm) | Number of branches/plant | Days to first flowering | Days to 50 per cent flowering | Fruit width (cm) | Fruit length (cm) | Number of locules per fruit | Pericarp thickness (mm) |
|---------|---------------|-----------|-------------------|--------------------------|-------------------------|-------------------------------|------------------|-------------------|-----------------------------|-------------------------|
| 128 | 15P4 | G-166 | 102.33 | 7.00 | 22.00 | 24.00 | 4.55 | 3.55 | 3.00 | 2.69 |
| 129 | 16P2 | G-167 | 61.33 | 5.67 | 28.00 | 33.00 | 3.00 | 2.60 | 4.00 | 3.79 |
| 130 | 17P5 | G-168 | 86.67 | 7.33 | 31.00 | 37.00 | 3.25 | 2.85 | 4.00 | 2.08 |
| 131 | 18P3 | G-169 | 115.33 | 5.67 | 29.00 | 33.00 | 3.70 | 3.45 | 3.00 | 3.42 |
| 132 | 19P8 | G-170 | 81.00 | 6.67 | 26.00 | 33.00 | 4.20 | 3.85 | 4.00 | 4.05 |
| 133 | 20 | G-171 | 76.33 | 6.67 | 17.00 | 24.00 | 3.55 | 3.35 | 5.00 | 3.02 |
| 134 | 21 | G-172 | 82.33 | 9.00 | 30.00 | 35.00 | 4.10 | 3.20 | 5.00 | 2.41 |
| 135 | 23P4 | G-173 | 105.67 | 8.00 | 24.00 | 28.00 | 3.85 | 3.60 | 6.00 | 2.54 |
| 136 | 25P2 | G-175 | 78.33 | 7.33 | 30.00 | 37.00 | 3.75 | 3.10 | 4.00 | 6.23 |
| 137 | 27P2 | G-177 | 96.00 | 8.33 | 24.00 | 30.00 | 5.45 | 4.30 | 3.00 | 4.29 |
| 138 | 28P2 | G-178 | 115.00 | 9.67 | 28.00 | 33.00 | 2.85 | 3.45 | 4.00 | 3.63 |
| 139 | 29P4 | G-179 | 57.33 | 5.67 | 29.00 | 33.00 | 3.45 | 3.55 | 5.00 | 4.88 |
| 140 | 30P2 | G-180 | 96.00 | 9.33 | 24.00 | 37.00 | 3.90 | 3.75 | 2.00 | 3.26 |
| 141 | 33P2 | G-183 | 92.67 | 7.00 | 30.00 | 36.00 | 3.75 | 3.35 | 3.00 | 1.53 |
| 142 | 34P2 | G-184 | 61.00 | 6.33 | 29.00 | 35.00 | 3.50 | 4.35 | 2.00 | 2.45 |
| 143 | 35P2 | G-185 | 68.00 | 8.00 | 16.00 | 25.00 | 5.35 | 3.10 | 6.00 | 2.59 |
| 144 | 36 | G-186 | 73.33 | 5.00 | 33.00 | 37.00 | 4.55 | 4.95 | 3.00 | 2.60 |
| 145 | 37P2 | G-187 | 71.67 | 8.00 | 24.00 | 30.00 | 4.55 | 3.65 | 5.00 | 2.50 |
| 146 | 38P2 | G-188 | 87.00 | 6.00 | 24.00 | 30.00 | 4.05 | 3.70 | 3.00 | 1.37 |
| 147 | 40P4 | G-190 | 91.33 | 5.00 | 30.00 | 37.00 | 3.05 | 3.00 | 4.00 | 2.75 |
| 148 | 43 | G-193 | 80.67 | 6.67 | 30.00 | 35.00 | 2.85 | 3.00 | 3.00 | 1.65 |
| 149 | 44P2 | G-194 | 86.33 | 7.33 | 29.00 | 34.00 | 4.10 | 3.30 | 3.00 | 2.25 |
| 150 | 45 | G-195 | 84.67 | 7.33 | 29.00 | 33.00 | 4.35 | 3.60 | 4.00 | 3.19 |
| 151 | 46P5 | G-196 | 66.00 | 5.33 | 30.00 | 33.00 | 4.15 | 3.90 | 7.00 | 1.50 |
| 152 | 47P2 | G-197 | 59.00 | 5.33 | 29.00 | 33.00 | 3.30 | 2.95 | 4.00 | 3.50 |
| 153 | 48P4 | G-198 | 59.00 | 6.33 | 30.00 | 34.00 | 3.65 | 3.90 | 3.00 | 2.30 |
| 154 | 51 | G-201 | 103.33 | 8.33 | 28.00 | 30.00 | 4.10 | 3.50 | 2.00 | 4.45 |
| 155 | 53 | G-202 | 63.67 | 7.33 | 29.00 | 33.00 | 2.70 | 3.75 | 2.00 | 3.87 |
| 156 | 54P3 | G-203 | 79.00 | 8.00 | 29.00 | 37.00 | 4.15 | 3.55 | 3.00 | 2.00 |
| 157 | 55P2 | G-204 | 108.67 | 8.33 | 28.00 | 33.00 | 3.85 | 3.25 | 7.00 | 2.50 |
| 158 | 56P2 | G-205 | 80.00 | 6.67 | 22.00 | 29.00 | 3.80 | 4.00 | 5.00 | 1.40 |
| 159 | 58 | G-206 | 62.67 | 7.00 | 22.00 | 28.00 | 4.90 | 3.85 | 4.00 | 4.50 |

| Sl. No. | Original code | Genotypes | Plant height (cm) | Number of branches/plant | Days to first flowering | Days to 50 per cent flowering | Fruit width (cm) | Fruit length (cm) | Number of locules per fruit | Pericarp thickness (mm) |
|---------|------------------|-----------|-------------------|--------------------------|-------------------------|-------------------------------|------------------|-------------------|-----------------------------|-------------------------|
| 160 | 59 | G-207 | 44.67 | 8.33 | 28.00 | 34.00 | 4.30 | 3.35 | 4.00 | 2.53 |
| 161 | 61P4 | G-208 | 58.33 | 9.00 | 29.00 | 34.00 | 3.40 | 3.60 | 3.00 | 2.37 |
| 162 | 62P4 | G-209 | 88.00 | 7.33 | 28.00 | 40.00 | 2.50 | 3.90 | 3.00 | 1.63 |
| 163 | 63P3 | G-210 | 70.67 | 9.00 | 22.00 | 28.00 | 4.05 | 2.60 | 4.00 | 1.13 |
| 164 | 64 | G-211 | 56.67 | 7.67 | 30.00 | 42.00 | 4.00 | 3.50 | 5.00 | 1.32 |
| 165 | 65P5 | G-212 | 87.00 | 9.33 | 23.00 | 28.00 | 5.30 | 2.20 | 5.00 | 3.10 |
| 166 | 66P1 | G-213 | 54.33 | 7.33 | 29.00 | 37.00 | 4.00 | 4.45 | 5.00 | 4.92 |
| 167 | 70 | G-214 | 52.33 | 8.33 | 24.00 | 30.00 | 3.10 | 3.60 | 2.00 | 3.28 |
| 168 | 71P2 | G-215 | 51.00 | 7.67 | 22.00 | 28.00 | 3.75 | 4.55 | 3.00 | 3.12 |
| 169 | 72P2 | G-216 | 50.33 | 8.33 | 29.00 | 33.00 | 5.05 | 3.15 | 6.00 | 2.98 |
| 170 | 73P2 | G-217 | 95.00 | 9.00 | 28.00 | 33.00 | 2.50 | 2.00 | 3.00 | 2.78 |
| 171 | 74P5 | G-218 | 51.00 | 8.00 | 22.00 | 29.00 | 3.95 | 3.05 | 4.00 | 1.60 |
| 172 | 75P3 | G-219 | 54.00 | 6.67 | 30.00 | 34.00 | 4.90 | 3.15 | 4.00 | 3.70 |
| 173 | 76P1 | G-220 | 92.00 | 4.00 | 29.00 | 33.00 | 3.90 | 2.95 | 4.00 | 2.44 |
| 174 | 77P1 | G-221 | 92.00 | 6.33 | 25.00 | 28.00 | 3.80 | 4.00 | 5.00 | 3.62 |
| 175 | 78P4 | G-222 | 113.33 | 6.33 | 29.00 | 33.00 | 2.70 | 2.30 | 3.00 | 3.62 |
| 176 | 83P1 | G-223 | 80.67 | 6.67 | 15.00 | 23.00 | 4.65 | 3.05 | 5.00 | 2.99 |
| 177 | 84P1 | G-224 | 60.33 | 5.33 | 29.00 | 33.00 | 3.85 | 5.05 | 4.00 | 1.66 |
| 178 | 86P2 | G-225 | 89.67 | 7.00 | 17.00 | 24.00 | 3.85 | 3.90 | 5.00 | 3.37 |
| 179 | 67 | G-226 | 82.00 | 5.67 | 29.00 | 36.00 | 2.85 | 3.30 | 2.00 | 2.21 |
| 180 | 68 | G-227 | 66.67 | 6.67 | 28.00 | 33.00 | 3.80 | 3.40 | 4.00 | 1.27 |
| 181 | Nandhi | G-228 | 65.33 | 3.33 | 17.00 | 33.00 | 3.45 | 3.85 | 2.00 | 2.20 |
| 182 | Pant Polyhouse-2 | G-229 | 107.33 | 6.33 | 24.00 | 27.00 | 4.70 | 4.75 | 2.00 | 1.62 |
| 183 | AR-28 | G-230 | 92.67 | 6.33 | 29.00 | 33.00 | 3.05 | 4.85 | 3.00 | 2.74 |
| 184 | Anaga | G-231 | 55.33 | 2.67 | 33.00 | 37.00 | 2.65 | 3.00 | 7.00 | 1.45 |
| 185 | IIHR-2195 | G-233 | 67.00 | 3.33 | 15.00 | 24.00 | 3.00 | 3.00 | 2.00 | 2.55 |
| 186 | AR-21 | G-234 | 96.67 | 5.67 | 15.00 | 25.00 | 5.20 | 4.00 | 6.00 | 1.69 |
| 187 | AR-56 | G-235 | 132.67 | 11.33 | 16.00 | 24.00 | 3.20 | 4.00 | 4.00 | 2.99 |
| 188 | AR-4 | G-237 | 109.33 | 7.00 | 23.00 | 28.00 | 3.65 | 4.00 | 5.00 | 1.62 |
| 189 | IHR-2199 | G-239 | 66.00 | 6.00 | 28.00 | 24.00 | 4.00 | 4.30 | 3.00 | 1.80 |
| 190 | IIHR-2198 | G-240 | 75.00 | 4.00 | 25.00 | 28.00 | 4.05 | 4.35 | 2.00 | 3.11 |
| 191 | IHR-2197 | G-242 | 59.00 | 2.00 | 23.00 | 27.00 | 4.10 | 5.05 | 3.00 | 4.15 |

| Sl. No. | Original code | Genotypes | Plant height (cm) | Number of branches/plant | Days to first flowering | Days to 50 per cent flowering | Fruit width (cm) | Fruit length (cm) | Number of locules per fruit | Pericarp thickness (mm) |
|----------------|----------------------|------------------|--------------------------|---------------------------------|--------------------------------|--------------------------------------|-------------------------|--------------------------|------------------------------------|--------------------------------|
| 192 | AR-29 | G-244 | 93.33 | 8.67 | 20.00 | 33.00 | 3.00 | 3.50 | 2.00 | 1.50 |
| 193 | IIHR-2196 | G-246 | 60.00 | 5.67 | 28.00 | 32.00 | 3.25 | 4.45 | 2.00 | 2.95 |
| 194 | Utkal Local-2 | G-247 | 98.00 | 5.67 | 37.00 | 43.00 | 3.95 | 3.30 | 4.00 | 2.15 |
| 195 | Pant Polyhouse | G-248 | 138.33 | 6.67 | 30.00 | 44.00 | 4.65 | 3.50 | 4.00 | 2.90 |
| 196 | Pant-3 | G-249 | 55.33 | 9.00 | 24.00 | 27.00 | 3.70 | 4.00 | 5.00 | 1.89 |
| 197 | IIHR-2200 | G-250 | 65.00 | 5.33 | 27.00 | 32.00 | 3.75 | 4.05 | 3.00 | 1.75 |
| 198 | IIHR-2201 | G-251 | 60.33 | 6.33 | 23.00 | 26.00 | 4.10 | 4.75 | 2.00 | 2.50 |
| 199 | H-24-1 | G-252 | 57.33 | 8.33 | 28.00 | 33.00 | 4.40 | 3.50 | 5.00 | 2.58 |
| 200 | Solan-2 | G-254 | 80.33 | 10.67 | 27.00 | 37.00 | 4.20 | 3.65 | 3.00 | 4.03 |

Table 5. Per se performance of 200 tomato genotypes for fruit weight, average yield and quality attributes

| Sl. No | Original code | Genotypes | Fruits per plant | Average fruit weight (g) | Yield per plant (kg/cm ²) | TSS (°Brix) | Firmness (kg/cm ²) |
|--------|-----------------------|-----------|------------------|--------------------------|---------------------------------------|-------------|--------------------------------|
| 1 | ARTD-1 | G-1 | 40.70 | 52.85 | 2.15 | 3.33 | 0.84 |
| 2 | ARTD-2 | G-2 | 34.00 | 69.05 | 1.49 | 2.83 | 2.41 |
| 3 | ARTD-3 | G-3 | 48.09 | 61.25 | 2.95 | 2.67 | 1.65 |
| 4 | ARTD-4 | G-4 | 37.97 | 54.38 | 2.06 | 2.83 | 2.81 |
| 5 | ARTD-5 | G-5 | 57.38 | 56.19 | 3.22 | 3.40 | 2.69 |
| 6 | ARTD-6 | G-6 | 46.40 | 64.71 | 3.00 | 3.07 | 3.37 |
| 7 | ARTD-7 | G-7 | 23.24 | 43.80 | 1.02 | 2.67 | 1.63 |
| 8 | ARTD-8 | G-8 | 41.93 | 63.17 | 2.65 | 3.03 | 1.43 |
| 9 | AOTD-9 | G-9 | 48.89 | 51.97 | 2.54 | 3.00 | 1.17 |
| 10 | AOTD-10 | G-10 | 64.24 | 56.98 | 3.66 | 2.70 | 1.46 |
| 11 | AOTD-11 | G-11 | 49.03 | 49.62 | 2.43 | 5.03 | 2.27 |
| 12 | AOTD-12 | G-12 | 75.62 | 42.50 | 3.21 | 3.67 | 1.84 |
| 13 | AOTD-13 | G-13 | 104.10 | 44.85 | 4.67 | 3.70 | 1.97 |
| 14 | AOTD-14 | G-14 | 78.87 | 43.64 | 3.44 | 3.80 | 1.53 |
| 15 | AOTD-15 | G-15 | 83.30 | 42.45 | 3.54 | 3.33 | 0.96 |
| 16 | AOTD-16 | G-16 | 72.14 | 57.70 | 4.16 | 3.67 | 0.84 |
| 17 | AOTD-17 | G-17 | 32.41 | 75.24 | 2.44 | 3.03 | 1.42 |
| 18 | AOTD-18 | G-18 | 75.80 | 52.84 | 4.01 | 3.67 | 6.25 |
| 19 | AOTD-19 | G-19 | 23.45 | 66.50 | 1.56 | 3.67 | 0.80 |
| 20 | AOTD-20 | G-20 | 46.05 | 62.53 | 2.88 | 3.67 | 2.96 |
| 21 | EC-321426 | G-23 | 20.08 | 94.85 | 1.90 | 3.47 | 1.70 |
| 22 | EC-326146 | G-31 | 57.89 | 34.16 | 1.98 | 5.00 | 1.01 |
| 23 | EC-338714 | G-33 | 45.31 | 42.79 | 1.94 | 3.33 | 1.06 |
| 24 | EC-338717 | G-35 | 204.37 | 24.11 | 4.93 | 5.00 | 1.36 |
| 25 | EC-338725 | G-37 | 55.52 | 32.03 | 1.78 | 5.00 | 0.94 |
| 26 | EC-338735 | G-38 | 85.54 | 43.18 | 3.69 | 3.00 | 0.93 |
| 27 | EC-339057 | G-39 | 70.94 | 22.86 | 1.62 | 4.33 | 1.34 |
| 28 | EC-357839 | G-40 | 43.25 | 74.02 | 3.20 | 4.17 | 1.08 |
| 29 | EC-357845 | G-41 | 108.31 | 40.53 | 4.39 | 4.00 | 1.28 |
| 30 | EC-362940 | G-43 | 79.43 | 28.98 | 2.30 | 3.67 | 0.67 |
| 31 | Punjab Varkha Bahar-1 | G-46 | 38.98 | 72.90 | 2.84 | 4.17 | 1.37 |
| 32 | PNR-1 | G-47 | 44.35 | 61.80 | 2.74 | 6.67 | 0.96 |

| Sl. No | Original code | Genotypes | Fruits per plant | Average fruit weight (g) | Yield per plant (kg/cm ²) | TSS (°Brix) | Firmness (kg/cm ²) |
|--------|-----------------------|-----------|------------------|--------------------------|---------------------------------------|-------------|--------------------------------|
| 33 | S-12 | G-48 | 89.25 | 34.21 | 3.05 | 4.67 | 0.57 |
| 34 | Punjab Varkha Bahar-2 | G-49 | 58.49 | 73.09 | 4.28 | 4.67 | 2.27 |
| 35 | Hisar Lal | G-50 | 57.97 | 51.06 | 2.96 | 4.00 | 2.64 |
| 36 | Punjab Upama | G-51 | 75.66 | 58.12 | 4.40 | 3.73 | 1.97 |
| 37 | Punjab Chhuhar | G-52 | 97.69 | 51.44 | 5.03 | 3.33 | 1.06 |
| 38 | Punjab Kesari | G-53 | 42.35 | 40.40 | 1.71 | 3.70 | 0.76 |
| 39 | Punjab Ratta | G-54 | 42.25 | 79.95 | 3.38 | 3.80 | 2.31 |
| 40 | 52135 | G-57 | 49.32 | 46.29 | 2.28 | 4.13 | 1.17 |
| 41 | Azad-T-1 | G-61 | 62.76 | 60.52 | 3.80 | 5.07 | 1.45 |
| 42 | Kashi Sharad | G-62 | 27.68 | 95.00 | 2.63 | 4.60 | 1.16 |
| 43 | Kashi Hemanth | G-63 | 33.51 | 102.36 | 3.43 | 4.93 | 0.49 |
| 44 | Kashi Amrith | G-64 | 74.33 | 67.28 | 5.00 | 3.00 | 0.66 |
| 45 | Kashi Anupama | G-65 | 71.88 | 77.77 | 5.59 | 3.07 | 1.18 |
| 46 | H-86 | G-66 | 48.85 | 131.94 | 6.45 | 2.93 | 1.13 |
| 47 | Kalyanpur Type | G-68 | 81.32 | 48.87 | 3.97 | 4.03 | 1.14 |
| 48 | A Abha | G-69 | 37.41 | 63.91 | 2.39 | 4.00 | 1.45 |
| 49 | A Saurabh | G-70 | 39.42 | 52.30 | 2.06 | 6.07 | 1.05 |
| 50 | DT-10 | G-72 | 74.08 | 47.80 | 3.54 | 4.50 | 1.02 |
| 51 | H-24 | G-73 | 111.18 | 34.78 | 3.87 | 4.03 | 0.72 |
| 52 | PTR-4 | G-75 | 74.50 | 53.18 | 3.96 | 4.00 | 2.23 |
| 53 | PTR-6 | G-76 | 79.80 | 74.66 | 5.96 | 2.67 | 2.03 |
| 54 | Bhagyashree | G-77 | 58.78 | 90.86 | 5.34 | 3.93 | 1.19 |
| 55 | Dhanashree | G-78 | 85.83 | 52.05 | 4.47 | 3.73 | 1.02 |
| 56 | Pule Raja | G-79 | 65.48 | 76.65 | 5.02 | 4.07 | 1.61 |
| 57 | Utkal Kumari | G-80 | 55.18 | 56.61 | 3.12 | 3.60 | 0.89 |
| 58 | Utkal Deepti | G-81 | 135.55 | 26.13 | 3.54 | 4.60 | 0.99 |
| 59 | Utkal Raja | G-82 | 36.96 | 73.31 | 2.71 | 4.93 | 0.83 |
| 60 | Pragnya | G-83 | 87.68 | 35.72 | 3.13 | 3.60 | 1.34 |
| 61 | Roma | G-84 | 33.00 | 29.46 | 0.39 | 4.93 | 0.96 |
| 62 | Best of all | G-85 | 56.46 | 29.69 | 1.68 | 5.07 | 0.62 |
| 63 | Sioux | G-86 | 24.68 | 102.22 | 2.52 | 4.00 | 1.00 |
| 64 | PKM-1 | G-87 | 11.66 | 43.85 | 0.51 | 4.50 | 2.60 |
| 65 | Akshaya | G-88 | 64.57 | 43.18 | 2.79 | 2.77 | 1.88 |
| 66 | Rajamundri | G-90 | 39.56 | 56.17 | 2.22 | 3.90 | 1.77 |
| 67 | Guntur Local | G-91 | 38.28 | 52.26 | 2.00 | 4.00 | 1.61 |
| 68 | A Alok | G-92 | 36.55 | 74.22 | 2.71 | 3.60 | 1.06 |

| Sl. No | Original code | Genotypes | Fruits per plant | Average fruit weight (g) | Yield per plant (kg/cm ²) | TSS (°Brix) | Firmness (kg/cm ²) |
|--------|----------------|-----------|------------------|--------------------------|---------------------------------------|-------------|--------------------------------|
| 69 | A Vikash | G-93 | 58.08 | 34.46 | 2.00 | 4.90 | 1.62 |
| 70 | A Meghali | G-95 | 75.58 | 60.64 | 4.58 | 4.27 | 0.92 |
| 71 | A Ahuthi | G-96 | 43.00 | 56.02 | 2.41 | 3.27 | 0.83 |
| 72 | Madanapalli | G-98 | 19.69 | 48.19 | 2.50 | 4.27 | 0.62 |
| 73 | Pusa Sheetal | G-99 | 50.34 | 50.68 | 2.55 | 3.87 | 0.78 |
| 74 | Pusa Sadabahar | G-100 | 29.76 | 40.27 | 1.20 | 3.33 | 0.66 |
| 75 | Pusa Rohini | G-101 | 31.30 | 64.53 | 2.02 | 3.60 | 1.49 |
| 76 | Pusa Gourava | G-102 | 67.02 | 57.19 | 3.83 | 3.67 | 1.06 |
| 77 | P-120 | G-103 | 23.99 | 57.70 | 1.38 | 4.33 | 1.19 |
| 78 | Pusa Ruby | G-104 | 102.69 | 36.97 | 3.80 | 4.57 | 0.62 |
| 79 | S-22 | G-105 | 65.40 | 58.04 | 3.80 | 4.00 | 0.46 |
| 80 | Navodaya | G-106 | 48.38 | 31.81 | 1.54 | 4.40 | 0.64 |
| 81 | Selection | G-107 | 83.46 | 35.95 | 3.00 | 4.03 | 0.67 |
| 82 | C-10-2 | G-108 | 81.34 | 27.66 | 2.25 | 4.73 | 0.78 |
| 83 | C-11-2 | G-109 | 85.56 | 26.74 | 2.29 | 5.00 | 3.78 |
| 84 | C-20-1 | G-110 | 73.02 | 63.10 | 4.61 | 5.17 | 1.47 |
| 85 | CO-3 | G-111 | 96.68 | 39.41 | 3.81 | 4.57 | 1.16 |
| 86 | CLN-2026 | G-112 | 123.07 | 27.61 | 3.40 | 5.17 | 2.24 |
| 87 | 41 | G-113 | 32.00 | 58.35 | 0.61 | 3.67 | 1.71 |
| 88 | DVRT-2 | G-114 | 54.37 | 59.06 | 3.21 | 3.10 | 0.56 |
| 89 | EC-13904 | G-116 | 51.12 | 45.58 | 2.33 | 3.80 | 0.67 |
| 90 | C-1-4 | G-119 | 55.49 | 44.46 | 2.47 | 3.40 | 1.29 |
| 91 | C-4-1 | G-120 | 80.72 | 25.67 | 2.07 | 4.40 | 1.85 |
| 92 | EC-381263 | G-121 | 133.39 | 22.64 | 3.02 | 4.93 | 2.22 |
| 93 | EC-501574 | G-123 | 41.29 | 41.85 | 1.73 | 4.40 | 0.89 |
| 94 | EC-501580 | G-124 | 54.13 | 74.64 | 4.04 | 4.77 | 1.71 |
| 95 | EC-501583 | G-126 | 101.61 | 25.70 | 2.61 | 5.43 | 1.18 |
| 96 | EC-538404 | G-129 | 38.43 | 74.65 | 2.87 | 4.43 | 1.10 |
| 97 | EC-538405 | G-130 | 58.16 | 70.93 | 4.13 | 4.63 | 0.80 |
| 98 | EC-620383 | G-133 | 88.54 | 27.92 | 2.47 | 6.00 | 1.10 |
| 99 | EC-620398 | G-134 | 29.61 | 36.53 | 1.08 | 4.33 | 1.04 |
| 100 | EC-620401 | G-135 | 61.04 | 62.49 | 3.81 | 3.93 | 2.57 |
| 101 | EC-620446 | G-136 | 52.03 | 82.28 | 4.28 | 4.13 | 1.13 |
| 102 | EC-620464 | G-137 | 18.82 | 54.46 | 1.03 | 4.93 | 1.00 |
| 103 | EC-620469 | G-138 | 45.78 | 57.08 | 2.61 | 4.47 | 1.88 |
| 104 | EC-620470 | G-139 | 33.00 | 28.38 | 0.57 | 4.97 | 1.24 |

| Sl. No | Original code | Genotypes | Fruits per plant | Average fruit weight (g) | Yield per plant (kg/cm ²) | TSS (°Brix) | Firmness (kg/cm ²) |
|--------|----------------|-----------|------------------|--------------------------|---------------------------------------|-------------|--------------------------------|
| 105 | Monte Fevarate | G-140 | 49.15 | 43.36 | 2.13 | 4.97 | 1.18 |
| 106 | Rio Grande | G-141 | 62.95 | 22.76 | 1.43 | 4.00 | 1.61 |
| 107 | Angoorlata | G-143 | 30.89 | 28.93 | 0.89 | 4.23 | 1.86 |
| 108 | Ageta-2 | G-144 | 28.96 | 57.65 | 1.67 | 4.00 | 1.81 |
| 109 | 85 | G-145 | 45.92 | 42.74 | 1.96 | 4.23 | 0.82 |
| 110 | 4 | G-148 | 74.70 | 40.23 | 3.01 | 5.00 | 1.11 |
| 111 | 80 | G-149 | 76.08 | 45.31 | 3.45 | 4.83 | 1.49 |
| 112 | 200 | G-150 | 70.76 | 52.69 | 3.73 | 3.57 | 1.28 |
| 113 | 8 | G-151 | 125.16 | 30.45 | 3.81 | 3.93 | 1.10 |
| 114 | 1P2 | G-152 | 114.36 | 31.98 | 3.66 | 4.37 | 0.80 |
| 115 | 2P2 | G-153 | 99.43 | 31.63 | 3.15 | 5.03 | 1.03 |
| 116 | 3P2 | G-154 | 154.75 | 36.83 | 5.70 | 5.33 | 0.81 |
| 117 | 4P1 | G-155 | 108.31 | 59.79 | 6.48 | 3.17 | 0.35 |
| 118 | 5T5P6 | G-156 | 121.91 | 38.88 | 4.74 | 5.17 | 2.40 |
| 119 | 6T6P8 | G-157 | 33.34 | 54.68 | 1.82 | 4.33 | 1.64 |
| 120 | 7 | G-158 | 76.46 | 54.13 | 4.14 | 4.43 | 0.71 |
| 121 | 8P3 | G-159 | 60.54 | 48.57 | 2.94 | 3.90 | 1.01 |
| 122 | 9P4 | G-160 | 142.78 | 40.39 | 5.77 | 4.00 | 1.10 |
| 123 | 10P6 | G-161 | 116.16 | 40.53 | 4.71 | 4.00 | 0.83 |
| 124 | 11P4 | G-162 | 81.06 | 38.51 | 3.12 | 4.67 | 0.97 |
| 125 | 12P1 | G-163 | 68.12 | 50.17 | 3.42 | 3.67 | 2.47 |
| 126 | 13P3 | G-164 | 129.52 | 25.76 | 3.34 | 4.00 | 0.95 |
| 127 | 14P6 | G-165 | 146.23 | 30.97 | 4.53 | 4.67 | 1.06 |
| 128 | 15P4 | G-166 | 88.80 | 64.54 | 5.73 | 4.00 | 1.20 |
| 129 | 16P2 | G-167 | 99.61 | 48.77 | 4.86 | 3.67 | 1.58 |
| 130 | 17P5 | G-168 | 81.47 | 39.28 | 3.20 | 6.67 | 1.47 |
| 131 | 18P3 | G-169 | 65.82 | 33.51 | 2.21 | 4.07 | 1.94 |
| 132 | 19P8 | G-170 | 97.58 | 37.35 | 3.64 | 4.00 | 0.97 |
| 133 | 20 | G-171 | 88.40 | 41.97 | 3.71 | 4.67 | 0.57 |
| 134 | 21 | G-172 | 56.01 | 52.59 | 2.95 | 4.37 | 0.59 |
| 135 | 23P4 | G-173 | 96.04 | 43.99 | 4.23 | 3.83 | 0.93 |
| 136 | 25P2 | G-175 | 125.45 | 32.12 | 4.03 | 5.17 | 0.99 |
| 137 | 27P2 | G-177 | 132.23 | 41.51 | 5.49 | 3.33 | 0.60 |
| 138 | 28P2 | G-178 | 173.31 | 22.89 | 3.97 | 5.00 | 2.54 |
| 139 | 29P4 | G-179 | 95.39 | 30.61 | 2.92 | 5.00 | 1.37 |
| 140 | 30P2 | G-180 | 118.69 | 43.27 | 5.14 | 4.90 | 1.01 |

| Sl. No | Original code | Genotypes | Fruits per plant | Average fruit weight (g) | Yield per plant (kg/cm ²) | TSS (°Brix) | Firmness (kg/cm ²) |
|--------|---------------|-----------|------------------|--------------------------|---------------------------------------|-------------|--------------------------------|
| 141 | 33P2 | G-183 | 135.63 | 34.00 | 3.68 | 5.00 | 1.35 |
| 142 | 34P2 | G-184 | 50.43 | 59.62 | 3.01 | 3.67 | 2.76 |
| 143 | 35P2 | G-185 | 82.38 | 43.14 | 3.55 | 5.33 | 0.50 |
| 144 | 36 | G-186 | 42.78 | 51.75 | 2.21 | 4.67 | 0.98 |
| 145 | 37P2 | G-187 | 56.76 | 36.85 | 2.09 | 4.33 | 0.55 |
| 146 | 38P2 | G-188 | 92.06 | 40.03 | 3.69 | 5.00 | 1.55 |
| 147 | 40P4 | G-190 | 56.27 | 33.89 | 1.91 | 4.33 | 1.12 |
| 148 | 43 | G-193 | 48.06 | 31.80 | 1.53 | 4.67 | 1.69 |
| 149 | 44P2 | G-194 | 18.86 | 52.73 | 1.30 | 5.83 | 0.92 |
| 150 | 45 | G-195 | 51.17 | 46.48 | 2.38 | 4.00 | 1.10 |
| 151 | 46P5 | G-196 | 37.26 | 63.94 | 2.38 | 3.90 | 0.37 |
| 152 | 47P2 | G-197 | 38.08 | 64.45 | 2.45 | 2.77 | 0.84 |
| 153 | 48P4 | G-198 | 181.58 | 30.85 | 5.60 | 4.23 | 1.84 |
| 154 | 51 | G-201 | 69.51 | 37.83 | 2.63 | 6.00 | 0.94 |
| 155 | 53 | G-202 | 86.16 | 40.04 | 3.45 | 4.33 | 1.96 |
| 156 | 54P3 | G-203 | 91.70 | 54.25 | 4.97 | 5.23 | 1.13 |
| 157 | 55P2 | G-204 | 56.00 | 54.05 | 3.03 | 5.67 | 1.82 |
| 158 | 56P2 | G-205 | 53.30 | 52.80 | 2.81 | 4.00 | 3.05 |
| 159 | 58 | G-206 | 30.88 | 73.31 | 2.26 | 3.40 | 2.42 |
| 160 | 59 | G-207 | 54.56 | 40.29 | 2.20 | 4.00 | 0.80 |
| 161 | 61P4 | G-208 | 110.96 | 35.00 | 2.53 | 4.50 | 0.92 |
| 162 | 62P4 | G-209 | 49.26 | 35.00 | 1.72 | 5.50 | 1.26 |
| 163 | 63P3 | G-210 | 87.08 | 40.68 | 3.54 | 3.90 | 1.29 |
| 164 | 64 | G-211 | 25.91 | 77.71 | 2.01 | 4.47 | 0.80 |
| 165 | 65P5 | G-212 | 83.94 | 45.52 | 3.82 | 5.00 | 0.65 |
| 166 | 66P1 | G-213 | 72.72 | 37.44 | 2.72 | 3.83 | 1.01 |
| 167 | 70 | G-214 | 62.25 | 27.68 | 1.72 | 4.00 | 0.80 |
| 168 | 71P2 | G-215 | 62.48 | 42.79 | 2.67 | 3.50 | 0.94 |
| 169 | 72P2 | G-216 | 39.35 | 61.34 | 2.41 | 4.00 | 0.59 |
| 170 | 73P2 | G-217 | 120.81 | 20.00 | 1.92 | 4.53 | 0.90 |
| 171 | 74P5 | G-218 | 70.26 | 46.30 | 3.25 | 5.00 | 0.83 |
| 172 | 75P3 | G-219 | 55.23 | 45.31 | 2.50 | 4.00 | 1.54 |
| 173 | 76P1 | G-220 | 31.63 | 28.77 | 0.91 | 4.17 | 0.96 |
| 174 | 77P1 | G-221 | 22.47 | 50.65 | 1.14 | 3.83 | 1.08 |
| 175 | 78P4 | G-222 | 70.41 | 22.97 | 1.62 | 4.50 | 1.30 |
| 176 | 83P1 | G-223 | 51.18 | 47.38 | 2.43 | 5.50 | 0.33 |

| Sl. No | Original code | Genotypes | Fruits per plant | Average fruit weight (g) | Yield per plant (kg/cm ²) | TSS (°Brix) | Firmness (kg/cm ²) |
|--------|------------------|-----------|------------------|--------------------------|---------------------------------------|-------------|--------------------------------|
| 177 | 84P1 | G-224 | 52.77 | 55.51 | 2.93 | 4.33 | 0.58 |
| 178 | 86P2 | G-225 | 75.41 | 45.15 | 3.40 | 4.00 | 1.54 |
| 179 | 67 | G-226 | 63.60 | 48.53 | 3.09 | 4.07 | 2.10 |
| 180 | 68 | G-227 | 40.49 | 59.26 | 2.40 | 4.67 | 0.95 |
| 181 | Nandhi | G-228 | 13.84 | 31.35 | 2.00 | 4.50 | 2.72 |
| 182 | Pant Polyhouse-2 | G-229 | 17.29 | 83.04 | 1.44 | 3.67 | 0.78 |
| 183 | AR-28 | G-230 | 92.10 | 24.25 | 2.23 | 4.17 | 0.80 |
| 184 | Anaga | G-231 | 57.03 | 35.23 | 2.01 | 4.00 | 0.56 |
| 185 | IIHR-2195 | G-233 | 111.66 | 24.92 | 2.78 | 4.00 | 1.53 |
| 186 | AR-21 | G-234 | 71.05 | 47.71 | 3.39 | 4.83 | 1.95 |
| 187 | AR-56 | G-235 | 140.65 | 21.00 | 2.26 | 7.00 | 0.72 |
| 188 | AR-4 | G-237 | 85.45 | 38.21 | 3.27 | 5.33 | 0.92 |
| 189 | IHR-2199 | G-239 | 78.29 | 38.04 | 2.98 | 3.83 | 1.25 |
| 190 | IIHR-2198 | G-240 | 30.09 | 42.63 | 1.28 | 4.00 | 1.45 |
| 191 | IHR-2197 | G-242 | 33.00 | 30.06 | 2.00 | 4.00 | 1.78 |
| 192 | AR-29 | G-244 | 106.14 | 23.47 | 2.49 | 5.67 | 1.14 |
| 193 | IIHR-2196 | G-246 | 102.02 | 24.45 | 2.49 | 4.50 | 1.92 |
| 194 | Utkal Local-2 | G-247 | 31.00 | 62.69 | 2.00 | 4.50 | 1.01 |
| 195 | Pant Polyhouse | G-248 | 21.05 | 73.38 | 1.55 | 4.00 | 2.33 |
| 196 | Pant-3 | G-249 | 124.36 | 30.46 | 3.79 | 4.50 | 0.59 |
| 197 | IIHR-2200 | G-250 | 23.91 | 55.44 | 1.33 | 4.50 | 2.77 |
| 198 | IIHR-2201 | G-251 | 42.89 | 41.17 | 1.77 | 3.83 | 1.53 |
| 199 | H-24-1 | G-252 | 53.62 | 41.20 | 2.21 | 4.17 | 0.71 |
| 200 | Solan-2 | G-254 | 69.17 | 49.50 | 3.42 | 5.17 | 1.07 |