

# Original Research Article

## Weed dynamics of fodder maize as influenced by different herbicides in Central India

### ABSTRACT

A field experiment was conducted at Research Farm, AICRP on Forage Crops, Department of Agronomy, JNKVV, Jabalpur (Madhya Pradesh) during *Kharif* season 2019. The field experiment was laid out in randomized block design (RBD) with ten treatments and replicated thrice. Ten treatments

were atrazine 1000 g/ha, pendimethalin 750g/ha, atrazine 750g/ha + pendimethalin 750 g/ha, 2,4-D 500 g/ha, tembotrione 120 g/ha, topramezone 35 g/ha, tembotrione 120 g/ha + atrazine 250 g/ha, topramezone 35 g/ha + atrazine 250 g/ha, hand weeding twice at 20 and 40 DAS and weedy check.

Among monocots, the experimental field was infested with *Echinochloa colona*, *Commelinacommunis* and *Digitariasanguinalis*. *Phyllanthus niruri* and *Eclipta alba* were dominated among dicots. Among all herbicidal treatment, topramezone 35 g/ha plus atrazine 250 g/ha was found to be significantly superior in reducing total weed density (9.49/m<sup>2</sup>, 11.50/m<sup>2</sup>) and total weed dry weight (6.43 g/m<sup>2</sup>, 7.91 g/m<sup>2</sup>) at 30 and 45 DAS, respectively and recorded the lowest weed index (1.82%). Also, the maximum green fodder yield (472.68 q/ha) was recorded in topramezone 35 g/ha + atrazine 250 g/ha compared to other treatments.

**Keywords:** Herbicidal application, Weed dynamics, Weed control efficiency, Weed index, Yield

### 1. INTRODUCTION

Population growth causes the demand for animal products like milk, eggs, and meat to rise dramatically [1]. The increased demand for animal products has encouraged a growth in the number of cattle within the current farming system while also highlighting the importance of the security of feed and fodder in the nation [2]. Currently, the nation has a net deficit of 64.1% in feeds, 21.9% in dry crop residue, and 61.1% in green fodder [3]. Thus, the country's cattle industry would have a significant problem in meeting the demand for feed and fodder [4].

Maize is the most ideal and appropriate crop for fodder as well as silage production. Maize (*Zea mays* L.) belongs to the *Gramineae* family and is a staple food crops in the universe, second only to wheat and rice [5]. Maize has become an important grain because of its huge production potential and adaptability to a wide range of environments [6]. Maize assumes a special significance on account of

Comment [u1]: Are you only dealing with fodder maize or forage crops?

Formatted: Highlight

Comment [u2]: The author should include the conclusion in the abstract

its utilization as food, feed, fodder and stalk besides several industrial uses. It is one of the most important dual purpose cereal crops all over the world [7]. The production of good quality fodder and forage is of great importance for the development of livestock industry in the country[8].

Fodder maize encounters a serious weed issue because the majority of farmers don't use any weed management techniques, which results in a lower yield of green and dry fodder per unit area [9]. The quantities of growth factors used by weeds are thus unavailable to the crop; the extent of nutrient loss varies from 30-40% of the applied nutrients [10, 11]. Wider row spacing and initial slow crop growth make maize highly sensitive to weed competition up to six weeks growth period, when the maximum yield loss due to weed competition occurs [12].

Hand weeding is very cumbersome, tedious, and costly. The availability of labourers during the critical period of the crop-weed competition is another issue[13]. Chemical weed management by using post-emergence herbicides can lead to the efficient and cost effective control of weeds during critical period of crop weed competition[14], which may not be possible in manual or mechanical weeding due to its high cost of cultivation[15, 16]. Few herbicides like Atrazine, Oxyfluorfen, 2,4-D and Pendimethalin are available for weed control in maize. At present farmers are applying only 2, 4-D at 1.0 kg/ha or Atrazine at 1.0 kg/ha as post-emergence herbicides in maize, but these herbicides control only broad leaf weeds. Control of grasses and sedges remain a problem for the farmers, especially when the too high or too low soil moisture hinders the inter-cultural operation and scarcity of labour during critical stages of weeding [17, 18]. Hence, there is an immense need to find out the best chemical for effective weed management in maize. Hence, this study was undertaken to identify the best chemical weed management practices in fodder maize.

## 2. MATERIALS AND METHODS

The experiment was executed during the *Kharif* season 2019 at Research Farm, AICRP on Forage Crops, Department of Agronomy, JNKVV, Jabalpur (Madhya Pradesh). The experiment was carried out ~~inwith~~ a randomized complete block design with three replications. Ten different treatments for weed management were applied in the fodder maize field viz. T1- tembotrione 120 g a.i/ha, T2- topramezone 35 g a.i/ha, T3- atrazine 1000 g a.i/ha, T4- pendimethalin 750 g a.i/ha, T5- tembotrione 120 ga.i/ha + atrazine 250 g a.i/ha, T6- topramezone 35 g a.i/ha + atrazine 250 g a.i/ha, T7- atrazine 750 g a.i/ha + pendimethalin 750 g a.i/ha, T8- 2,4-D 500 g a.i/ha, T9- hand weeding twice at 20 and 40 DAS and T10- weedy check. African tall variety of fodder maize was sown with row spacing of 50 cm and seed rate of 40 kg/ha. The soil of the experimental field was neutral in reaction (pH 7.21) and

Comment [u3]: Which issue?

Formatted: Highlight

Formatted: Highlight

Comment [u4]: This section lack timing of applications. Were these treatments applied pre or post emergence?

Comment [u5]: Provide a name for this variety. What about plant to plant spacing?

low in organic carbon (0.54%) as well as with medium available nitrogen (231.56 kg/ha), medium available phosphorus (16.59 kg/ha) and medium available potassium (313.66 kg/ha) contents with normal electrical conductivity (0.33). Observations on weed density, weed dry matter, weed control efficiency, weed index and yield of fodder maize were recorded. The calculation of weed control efficiency is based on the reduction of the dry matter yield of the weeds in the treated plots compared with the weed control at 45 DAS, and expressed as a percentage [19].

$$\text{WCE} = \frac{\text{DMC} - \text{DMT}}{\text{DMC}} \times 100$$

Where,  
WCE = Weed control efficiency (%)  
DMC = Dry matter of weeds in weedy check plot  
DMT = Dry matter of weeds in treated plot

Weed Index was calculated using the formula [20]

$$\text{Weed Index} = \frac{\text{Yield from weed free plot} - \text{Yield from treated plot}}{\text{Yield from weed free plot}} \times 100$$

The data obtained on various parameters are tabulated and statistically analyzed. The significance of the difference between a pair of means was tested by the least significant difference (LSD) test at a significance level of 5%.

Comment [u6]: By which software?

Comment [u7]: there are issues with methodology and statistical analyses

### 3. RESULTS AND DISCUSSION

#### 3.1 Dominant weed flora

The dominant weeds associated with crop in the experimental field were mainly comprised of *Echinochloa colona*, *Cyperus rotundus*, *Commelinacommunis*, *Ecliptaalba*, *Digitariasanguinalis*, *Eleusineindica* and *Phyllanthusniruri*. While, other minor weeds were also present. Almost similar weed flora is reported in maize [21] and other crops [22].


#### 3.2 Relative density of weeds

The data for species-wise weeds observed in weedy check plots at the 30 and 45 DAS stages of maize are shown in Table 1 indicated that, monocot weeds are predominated (76.31%) compared to dicot weeds (16.68%). Other species of weeds also marked their presence in less number. At the 30 and 45 DAS stages, *Echinochloa colona* (35.17%), *Commelinacommunis* (12.43%), and *Digitariasanguinalis* (10.79%) were the most prevalent monocot weeds, whereas dicot weeds like

*Phyllanthusniruri* (10.12%) and *Ecliptaalba* (6.55%) were more sparsely distributed in the maize ecosystem. These findings were in conformity to those of [23, 24]

**Table 1. Species wise mean intensity and relative density of weeds inweedy check plots at 30 and 45 DAS**

S.No.	Weeds	Density (no/m <sup>2</sup> )		Mean	Relative density (%)
		30 DAS	45 DAS		
<b>A. Monocots</b>					
1.	<i>Echinochloacolona</i>	99.92	103	101.46	35.17
2.	<i>Digitariasanguinalis</i>	28.42	33.83	31.13	10.79
3.	<i>Eleusineindica</i>	16.57	24.63	20.60	7.14
4.	<i>Cyperusrotundus</i>	31.5	30.67	31.09	10.77
5.	<i>Commelinacommunis</i>	35.5	36.25	35.88	12.43
Sub-total				220.15	76.31
<b>B. Dicots</b>					
	<i>Phyllanthusniruri</i>	24	34.42	29.21	10.12
	<i>Eclipta alba</i>	16.8	21.00	18.90	6.55
Sub-total				48.11	16.68
<b>C. Other weeds</b>		16	24.50	20.25	7.01
<b>Total</b>				<b>288.51</b>	<b>100</b>



**Figure 1. Relative density of weeds at 30 and 45 DAS**

### 3.3 Density and dry weight of weeds:

Total weed density and dry weight at 30 and 45 DAS varied significantly due to different weed control treatments (Table 2). The results clearly show that the weedy check plots had the highest weed density and dry weight of total weeds, including minor weeds, at both stages due to weeds uninterrupted growth in the absence of weed control measures. However, chemical method of weed control resulted in an identical decrease in weed density and dry weight. Pre-emergence

Comment [u8]: Identical to what?

application of atrazine 1000 g/ha (%), pendimethalin 750 g/ha, atrazine 750 g/ha + pendimethalin 750 g/ha (%) and post-emergence application of 2,4-D 500 g/ha(%) slightly decreased the total density and dry weight of monocot and dicot weeds. However, the reduction was more pronounced when tembotrione 120 g/ha and topramezone @ 35 g/ha herbicides were applied. In comparison to other herbicides, the post-emergence applications of topramezone35 g/ha + atrazine 250 g/ha successfully controlled both monocot and dicot weeds and successfully recorded the lowest weed density (9.49, 11.50/m<sup>2</sup>) and dry weight (6.43, 7.91 g/m<sup>2</sup>) at 30 and 45 DAS, respectively. However, due to the removal of short-of weeds, hand weeding performed at 20 and 40 DAS reduced the density including dry weight of weeds to the greatest degree over herbicidal treatments [25, 26].

Comment [u9]: Pre or post?

**Table 2. Weed density (no./m<sup>2</sup>) and weed dry weight (g/m<sup>2</sup>) as influenced by different weed control treatments in fodder maize**

Comment [u10]: Was this data analyzed?

Treatments	Total weed density (no./m <sup>2</sup> )		Total dry weight (g/m <sup>2</sup> )	
	30 DAS	45 DAS	30 DAS	45 DAS
T <sub>1</sub> - Tembotrione 120 g/ha	11.50 (131.78)	13.67 (186.30)	9.10 (82.23)	10.52 (110.26)
T <sub>2</sub> -Topramezone 35 g/ha	11.04 (121.44)	13.20 (173.62)	8.31 (68.65)	9.76 (94.84)
T <sub>3</sub> -Atrazine 1000 g/ha	13.77 (189.25)	15.65 (244.60)	11.14 (123.69)	12.37 (152.62)
T <sub>4</sub> - Pendimethalin 75 g/ha	14.86 (220.34)	16.30 (265.40)	11.79 (138.59)	13.03 (169.28)
T <sub>5</sub> -Tembotrione 120 g/ha+ Atrazine 250 g/ha	10.08 (101.07)	12.24 (149.43)	7.33 (53.27)	8.79 (76.78)
T <sub>6</sub> - Topramezone 35 g/ha+ Atrazine 250 g/ha	9.49 (89.60)	11.50 (131.92)	6.43 (40.90)	7.91 (62.06)
T <sub>7</sub> -Atrazine 750 g/ha+Pendimethalin750g/ha	13.17 (173.05)	14.90 (221.43)	10.44 (108.40)	11.92 (141.49)
T <sub>8</sub> -2,4-D 500 g/ha	12.75 (162.14)	14.29 (203.78)	9.85 (96.60)	11.34 (128.20)
T <sub>9</sub> -Hand weeding	6.78 (45.47)	7.02 (48.80)	2.49 (5.70)	5.28 (27.51)
T <sub>10</sub> -Weedy Check	16.41 (268.70)	17.57 (308.30)	14.16 (199.96)	15.58 (242.27)
SEm±	0.06	0.15	0.06	0.08
CD at 5%	0.19	0.46	0.17	0.22

Transformed values:  $\sqrt{x + 0.5}$   
Original figures are given in parenthesis

### 3.4 Weed control efficiency

Among various weed control treatments, the highest weed control efficiency were recorded under hand weeding treatment and it was closely followed by topramezone 35 g/ha + atrazine 250 g/ha. These results indicate that in addition to post-emergence herbicides, imposition of hand weeding ultimately provided weed free and congenial environment as the outcome of improved weed control

efficiency of fodder maize crop. While, the lowest weed control efficiency was recorded with weedy check treatment. These results are in accordance with the results indicated by [27, 28].

### 3.5 Weed index

Weed index is a measure of reduction in the green fodder yield due to weed competition stress as against weed free treatment. Data showed that maximum yield loss of 31.53% was recorded under weedy check and weeds were not controlled in the entire crop season. Presence of weeds created stressful conditions for nutrients, space, soil moisture and light; thus, suppressed the growth and development of maize crops. The weed index was lowest (1.82%) in plots receiving application of topramezone 35 g/ha + atrazine 250 g/ha followed by tembotrione 120 g/ha + atrazine 250 g/ha (3.87 %) and topramezone 35 g/ha (10.93 %). The lower weed index values under aforesaid treatments are attributed to the reduced competition stress by weed. Therefore, the yield attributes in crop were superior which ultimately resulted into increased green fodder yield. These findings are in line with the results of [29, 30].

**Comment [u11]:** Your data shows the lowest weed index was achieved in plot that were hand weeded

### 3.6 Green fodder yield

Green fodder yield in response to a specific treatment is the outcome of a complex phenomenon that is influenced by both the production methods used and the genetic makeup of the crop plants. Depending on the associated weed species, their density, the length of the crop weed competition, etc., weeds significantly harmed the crop, and their cumulative impact was demonstrated in a lower crop yield. Due to intense competition stress from the start of the critical period of crop growth until the end, the green fodder yield was lowest (343.12 q/ha) in the plots getting no weed control measures (weedy check). All of the treated plots, which either received manual weeding or herbicidal treatments produced greater yields over weedy check plot. The maximum green fodder yield was noted in hand weeding treatment (473.17 q/ha). However, among the herbicidal applications, topramezone 35 g/ha + atrazine 250 g/ha recorded the maximum green fodder yield (472.68 q/ha). Due to the removal of weeds from between and within the rows as well as improved soil aeration, the crop in weed-free plots grew lushly. More space, water, light, and nutrients were thus made available for the better growth and development, leading to superior yield attributes and development, and ultimately the highest yield [31, 32].

**Comment [u12]:** Always indicated or in text table

**Table 3. Weed control efficiency (%), weed index (%) and yield as influenced by different weed control treatments in fodder maize**

Treatments	Weed control efficiency (%) at 45 DAS	Weed index (%)	Green fodder yield (q/ha)
T <sub>1</sub> - Tembotrione 120 g/ha	54.49	11.60	405.37
T <sub>2</sub> - Topramezone 35 g/ha	60.85	10.93	423.42
T <sub>3</sub> - Atrazine 1000 g/ha	37.00	20.38	366.11
T <sub>4</sub> - Pendimethalin 75 g/ha	30.13	22.50	354.79
T <sub>5</sub> - Tembotrione 120 g/ha+ Atrazine 250 g/ha	68.31	3.87	442.66
T <sub>6</sub> - Topramezone 35 g/ha + Atrazine 250 g/ha	74.38	1.82	472.68
T <sub>7</sub> - Atrazine 750 g/ha+Pendimethalin 750 g/ha	41.60	14.92	370.89
T <sub>8</sub> - 2,4-D 500 g/ha	47.08	13.93	372.36
T <sub>9</sub> - Hand weeding	88.64	0.00	473.17
T <sub>10</sub> - Weedy Check	0.00	31.53	343.12
SEm±	-	0.38	5.53
CD at 5%	-	1.14	16.44

#### 4. CONCLUSION

Different weeds were severely infested with fodder maize in the experimental area. One of the possible ways to reduce weed infestation and enhanced fodder productivity is the application of effective herbicides. Therefore, based on the current experiment, it can be concluded that, the application of topramezone 35 g/ha + atrazine 250 g/ha was successfully controls the complex weed flora associated with fodder maize and also gives the higher green fodder yield.

**Comment [u13]:** This conclusion is not supported by data and is contradictory. What about hand weeding?

#### REFERENCES

- Panwar D, Chouhan D, Singh D, Singh RP, Nepalia V. Performance of Fodder Maize (*Zea mays*L.) Under Varying Plant Densities and Fertility Levels. International Journal of Current Microbiology and Applied Sciences. 2020; 11: 255-260.
- Jha AK, Yadav PS, Shrivastava A, Upadhyay AK, Sekhawat LS, Verma B, Sahu MP. Effect of nutrient management practices on productivity of perennial grasses under high moisture condition. AMA, Agricultural Mechanization in Asia, Africa and Latin America. 2023; 54(3): 12283-12288.
- Chaudhary DP, Kumar A, Mandhania SS, Srivastava P, Kumar RS. Maize as Fodder? An alternative approach, Directorate of Maize Research, Pusa Campus, New Delhi, Technical Bulletin. 2012; 04:32.
- Kantwa SR, Agrawal RK, Jha A, Pathan SH, Patil SD, Choudhary M. Effect of different herbicides on weed control efficiency, fodder and seed yields of berseem (*Trifolium alexandrinum* L.) in central India. Range Management and Agroforestry. 2019; 40(2): 323-328.
- Uddin Md. Riadj, Faruq Md. Omar, Azam Md. Golam. Assessing the Effect of Weed Management Practices on Yield of Maize in Chittagong Hill Tracts of Bangladesh. Journal of Agricultural Science & Engineering Innovation (JASEI). 2020;1(1): 32-37.
- Gheith EMS, El-Badry OZ, Lamloom SF, Ali HM, Siddiqui MH, Ghareeb RY, El-Sheikh MH, Jebiril J, Abdelsalam NR and Kandil EE. Maize (*Zea mays* L.) Productivity and Nitrogen Use

- Efficiency in Response to Nitrogen Application Levels and Time. *Front. Plant Sci.* 2022; 13:941343.
7. Chaudhary Dharam, Jat Shankar, Kumar R, Kumar Ashwani, Kumar Balwinder. Fodder Quality of Maize: Its Preservation. 2014;153-159.
  8. Yadav PS, Kewat ML, Jha AK, Hemalatha K, Verma B. Effect of sowing management and herbicides on the weed dynamics of berseem (*Trifolium alexandrinum*). *Pharma Innovation.* 2023; 12(2): 2845-2848.
  9. Sanodiya Pratik, Jha A.K., Shrivastava Arti. Effect of integrated weed management on seed yield of fodder maize. *Indian Journal of Weed Science.* 2013; 45(3): 214-216.
  10. Jha AK, Shrivastava Arti, Raguvanshi NS. Effect of weed control practices on the fodder and seed productivity of Berseem under irrigated condition of Madhya Pradesh. *Range management & Agroforestry.* 2014; 35(1): 61-65.
  11. Mundra SL, Vyas AK, Maliwal PL. Effect of weed and nutrient management on nutrient uptake by maize (*Zea mays*) and weeds. *Indian Journal of Agronomy.* 2002; 47: 378-383.
  12. Nagalakshmi KVV, Chandrasekhar, Subbaiah G. Weed management for efficient use of nitrogen in rabi maize (*Zea mays* L.). *Andhra Agricultural Journal.* 2006; 53(1&2): 14-16.
  13. Sahu, M. P., Kewat, M. L., Jha, A. K., Sondhia, S., Choudhary, V. K., Jain, N., ... & Verma, B. Weed prevalence, root nodulation and chickpea productivity influenced by weed management and crop residue mulch. *AMA, Agricultural Mechanization in Asia, Africa and Latin America.* 2022;53(6):8511-8521.
  14. Verma B, Bhan M, Jha AK, Singh V, Patel R, Sahu MP, Kumar V. Weed management in direct-seeded rice through herbicidal mixtures under diverse agroecosystems. *AMA, Agricultural Mechanization in Asia, Africa and Latin America.* 2022;53(4):7299- 7306.
  15. Sahu, V., Kewat, M. L., Verma, B., Singh, R., Jha, A. K., Sahu, M. P., & Porwal, M. Effect of carfentrazone-ethyl on weed flora, growth and productivity in wheat. *The Pharma Innovation Journal* 2023; 12(3): 3621-3624.
  16. Triveni U, Patro SK, Bharathalakshmi M. Effect of different pre and post emergence herbicides on weed control, productivity and economics of maize. *Indian Journal of Weed Science.* 2017; 49(3): 231–235.
  17. Patel Raghav, Jha AK, Verma Badal, Kumbhare Rahul, Singh Richa. Bio- efficacy of pinoxaden as post-emergence herbicide against weeds in wheat crop. *Pollution research.* 2023;42(1):115-117.
  18. Shiv Swati, Agrawal S.B., Verma Badal, Yadav Pushpendra Singh, Singh Richa, Porwal Muskan, Sisodiya Jirtendra and Patel Raghav. Weed dynamics and productivity of chickpea as affected by weed management practices. *Pollution Research.* 2023; 42 (2):21-24.
  19. Mani VS, Malla ML, Gautam KC, Bhagwandas. Weed killing chemicals in potato cultivation. *Indian Farm.* 1973; 22:17-18.
  20. Gill GS, Kumar V. Weed index a new method for reporting weed control trials. *Indian J Agron.* 1969; 16:96-98.
  21. Sharma P, Duary B, Singh R. Tank mix application of tembotrione and atrazine to reduce weed growth and increase productivity of maize. *Indian Journal of Weed Science.* 2018; 50(3): 305–308.
  22. Verma B, Bhan M, Jha AK, Khatoon S, Raghuwanshi M, Bhayal L, Sahu MP, Patel Rajendra, Singh Vikash. Weeds of direct-seeded rice influenced by herbicide mixture. *Pharma Innovation.* 2022;11(2):1080-1082.
  23. Arvadiya LK, Raj VC, Patel TU, Arvadiya MK. Influences of plant population and weed management on weed flora and productivity of sweet corn (*Zea mays* L.). *Indian Journal of Agronomy.* 2012; 57(2): 162-167.
  24. Kumar B, Prasad S, Mandal D, Kumar R. Influence of Integrated Weed Management Practices on Weed Dynamics, Productivity and Nutrient Uptake of Rabi Maize (*Zea mays* L.). *International Journal of Current Microbiology and Applied Sciences.* 2017; 6(4): 1431-1440.
  25. Swetha K, Madhavi M, Pratibha G, Ramprakash T. Weed management with new generation herbicides in maize. *Indian Journal of Weed Science.* 2015; 47(4): 432–433.
  26. Mandal S, Mandal S, Nath S. Effect of integrated weed management on yield components, yield and economics of baby corn (*Zea mays* L.). *Annals of Agricultural Research.* 2004; 25(2): 242-244.
  27. Madhavi M, Ramprakash T, Srinivas A, Yakadri M. Topramezone (33.6% SC)+atrazine (50%) WP tank mix efficacy on maize. Biennial conference on “Emerging challenge in weed management” Organized by Indian Society of Weed Science. 2014; 15-17, February.

28. Malviya A, Malviya N, Singh B, Singh AK. Integrated weed management in maize (*Zea mays* L.) under rainfed conditions. Indian Journal of Dryland Agriculture Research & Development. 2012; 27(1):70-73.
29. Baldaniya MJ, Patel TU, Zinzala MJ, Gujjar PB, Sahoo S. Weed management in fodder maize (*Zea mays* L.) with newer herbicides. International Journal of Chemical Studies. 2018; 6(5): 2732-2734.
30. Kolage AK, Shinde SH, Bhilare RL. Weed management in kharif maize. Journal of Maharashtra Agricultural Universities. 2004; 29(1):110-111.
31. Subrahmanya DJ, Kumar Rakesh, TyagiNitin, RamHardev, Singh Magan, Meena RK, Tamta A, Pandey AK. Yield of Fodder Maize (*Zea mays*) and its Chemical Composition under Varying Plant Densities and Nutrient Management. Indian Journal of Animal Nutrition. 2017; 34(4): 425-429.
32. Tripathi AK, Tewari AN, Prasad A. Integrated weed management in rainy season maize (*Zea mays* L.) in Central Uttar Pradesh. Indian Journal of Weed Science. 2005; 37(3&4):269-270.

UNDER PEER REVIEW