

Original Research Article

Impact of pre and post-emergence herbicides on weed flora of maize

Comment [A1]: Effect

ABSTRACT

Weeds have been a problem creator in crops since time immemorial competing with them for all the growth the resources. Weeds cause around 28-100% loss in the maize crop which necessitates their control. The usage of herbicides for the control of weeds has been the most prominent method among farmers. However, the continuous usage of similar herbicides can develop herbicide resistance which needs the exploration to newer herbicides. Therefore, a field experiment was carried out to assess the effect of broad-spectrum herbicide, mesotrione on the weed flora and seed yield of maize at JNKVV, Jabalpur (M.P.) in the *kharif* season of 2019. The experiment was carried out in randomised block design consisting of eight treatments of weed control including six herbicidal treatments with hand weeding and weedy check and replicated thrice. Based on the results, it was concluded that the post emergence application of mesotrione 350 g ha⁻¹ at 30 DAS had most effectively controlled the weeds resulting in highest weed control efficiency (69.25%) and lowest weed index (12.62%) than the rest of the treatments. It also resulted in highest grain and stover yield (2.44 and 21.80 t ha⁻¹) in maize. Thus, the application of mesotrione @ 350 g ha⁻¹ can be a promising technology for the control of complex weed flora in maize.

Keywords: *Herbicide resistance, Mesotrione, Stover yield, Weeds, Weed control efficiency, Weed index*

1. INTRODUCTION

Maize (*Zea mays* L.) is a significant and adaptable crop raised in a wide range of environments and geographical regions for human food, feed, and animal fodder [1]. Maize is one of the major cereal crops with wider adaptability to diverse agro-climatic conditions [2]. ~~With~~ **With a production of 945.8 million tonnes and a yield of 5.7 t ha⁻¹, maize is produced across an area of 168 million hectares worldwide.** In India, during 2018–19, it was cultivated over an area of 9.18 million ha, with an annual production of 27.23 million tonnes and average productivity of 2,965 kg/ha [3]. In Madhya Pradesh maize is cultivated in 1.35 million hectares with production of 3.54 million tonnes with a productivity of 2615 kg ha⁻¹ [4].

Comment [A2]: Give Reference

Weeds rank as maize's number one enemy and can be attributed for the low production of the crop under Indian circumstances. Weeds lower the photosynthetic efficiency, dry matter production, and distribution to economically advantageous areas, which in turn diminishes the crop's sink capacity and leads to a low grain yield [5]. Accordingly, depending on the kind of weed species in the standing crop, the amount of the decline in maize grain production has been observed to be in the range of 28 to 100% [6, 7].

Comment [A3]: cause

Comment [A4]:

Comment [A5]: Delete

Many weed management techniques, including mechanical, cultural, chemical, and biological ones, are available to reduce weed losses [8, 9]. The most effective and extensively used method of weed management is manual weeding, although it is labor-intensive, time-consuming, and expensive owing to high pay rates, which reduce the profitability of the cultivation [10]. Timely weeding is most important to minimize yield losses and therefore under such circumstances the only effective tool is left to control weeds through the use of chemicals [11].

Management of weeds through the use of chemicals has also been found as effective as realized under manual eradication in various crops with over and above benefits in saving extra costs involved in use of labour on manual eradication of weed [12]. Therefore, keeping the weeds below threshold level, herbicides provide the low-cost and effective tool through which excessive weed population can be controlled before crop-weed competition [13].

The use of herbicides is the best approach for quickly and cheaply resolving the various weed issues in maize fields, and as a result, it has surpassed conventional methods in importance [14]. Pre-emergence herbicides have been recommended as the best choice since they can suppress weeds during the early phases of crop growth and also provide an environment free from weed competition to guarantee improved crop establishment [15, 16]. This is especially true for maize. However, weeds may develop a resistance problem if a single herbicide or herbicides with the same mechanism of action are used continuously. As a result, new herbicides are required to manage the mixed weed flora in maize.

Mesotrione is a herbicide of broad-spectrum, selective stem and leaf before bud and after seedling mainly used to effectively control broad-leaved weeds and some gramineous weeds by inhibiting the catalytic factors of plant photosynthetic process [17, 18]. Mesotrione is widely used in weed control of maize and winter wheat due to its high activity, low residue, strong compatibility and safety to the environment and

Comment [A6]: Not clear

subsequent crops [19, 20]. In order to assess the possibility of mesotrione application and understand the related mechanism, we investigated the weed control in this current study.

2. MATERIALS AND METHODS

An experimental trial was conducted at Research Farm, AICRP on Forage Crops, Department of Agronomy, JNKVV, Jabalpur (Madhya Pradesh) during *kharif* season of the year 2019. The soil of the research field was neutral in reaction (pH 7.21), and medium in organic carbon (0.58%) as well as with medium available nitrogen (285.56 kg ha⁻¹), available phosphorus (16.59 kg ha⁻¹) and available potassium (262.66 kg ha⁻¹) contents with normal electrical conductivity (0.33 dS/cm). The experiment consisted of eight treatments viz., pre-emergence application of atrazine 1000 g ha⁻¹, pendimethalin @750 g ha⁻¹ and post emergence application of mesotrione 250 g ha⁻¹, mesotrione 300 g ha⁻¹, mesotrione 350 g ha⁻¹ and tembotrione 286 g ha⁻¹ and hand weeding twice at 20 and 40 DAS and weedy check laid out in randomized block design with three replications. Sowing of maize cultivar "African tall" at the rate of 20 kg/ha was carried out at a spacing of 60 cm. The crop was cultivated as per recommended practices. Weed parameters such as relative density of weeds, weed density and dry weight, weed index, weed control efficiency and grain and stover yields of maize were recorded in the present investigation. The data were statistically analysed and then the results were interpreted.

Comment [A7]: Delete

Comment [A8]: Add rain fall data and temperatures

Comment [A9]: Was this pre ? Indicate

Comment [A10]: Pre or Post

Comment [A11]: 60 cm x ?

Comment [A12]: What are they? Did you apply fertilisers etc?

Comment [A13]: Mention how each was carried out

Comment [A14]: How? Using which methods?

3. RESULTS AND DISCUSSION

3.1 Weed flora

Different weeds severely infested maize in weedy check plot of experimental site. The relative density of predominant monocot weeds was *Echinochloa colona* (31.48%), *Commelina communis* (11.48%), *Digitaria sanguinalis* (11.37%), *Cyperus rotundus* (10.31%) and *Eleusine indica* (7.79%) and relative density of broad-leaved weeds was *Phyllanthus niruri* (9.78%) and *Eclipta alba* (9.36%) and relative density of other minor weeds was 8.42%. Similar weed flora also observed by [21, 22].

Comment [A15]: Is the Data in any Table? Indicate Table no. or Fig no.

Comment [A16]: Did the researchers carry out asimilar study to get similar results?




Figure 1. Relative density of weeds in weedy check at 30 and 60 DAS

3.2 Weeds density and dry weight

The weed density and dry weight was significantly affected by different weed control treatments at 30 and 60 DAS in comparison to weedy check plot (Table 1-4). It is evident from the data that the density and dry weight of all the weeds was maximum under weedy check plot due to uninterrupted growth of weeds as no weed control measures were adopted in weedy check plots. However, considerable reduction in density and dry weight of weeds were observed when weeds were controlled by chemical method of weed control [23, 24]. Among herbicides, post emergence application of mesotrione 350 g/ha controlled all the weed species successfully and recorded a lower weed population fb mesotrione 300 g/ha and tembotrione 286 g ha⁻¹. Similar findings have been reported by [25, 26]. Among the weed control treatments, hand weeding done at 20 and 40 DAS reduced the density including dry weight of weeds to maximum extent over herbicidal treatments due to elimination of most of weeds. Similar view was also endorsed by [27, 28].

Comment [A17]: No LSD figures to show significant differences in Table

Comment [A18]: Pre or Post?

Comment [A19]: Wrong

Comment [A20]:

Comment [A21]: Not clear

Comment [A22]: What did they report?

Table 1. Effect of different weed control treatments on weed density (no./m²) at 30 and 60 DAS (grassy weeds and sedges)

Treatments	<i>Echinochloa colona</i>		<i>Digitaria sanguinalis</i>		<i>Eleusine indica</i>		<i>Cyperus rotundus</i>	
	30 DAS	60 DAS	30 DAS	60 DAS	30 DAS	60 DAS	30 DAS	60 DAS
T ₁ : Mesotrione 250 g ha ⁻¹	7.28 (52.50)	7.80 (60.42)	4.65 (21.17)	5.49 (29.67)	2.94 (8.17)	4.14 (16.67)	4.32 (18.17)	4.87 (23.25)
T ₂ : Mesotrione 300 g ha ⁻¹	6.67 (44.00)	7.57 (56.75)	4.31 (18.08)	5.09 (25.42)	2.36 (5.08)	3.59 (12.42)	4.19 (17.08)	4.67 (21.33)
T ₃ : Mesotrione 350 g ha ⁻¹	6.18 (37.67)	7.34 (53.42)	4.04 (15.83)	4.89 (23.42)	2.18 (4.25)	3.30 (10.42)	3.77 (13.75)	4.32 (18.42)

Comment [A23]: Insert Type of weeds on top

T ₄ : Tembotrione 286 g ha ⁻¹	6.92 (47.33)	7.61 (57.42)	4.40 (18.83)	5.28 (27.33)	2.41 (5.33)	3.85 (14.33)	4.26 (17.67)	4.86 (23.17)
T ₅ : Atrazine 1000 g ha ⁻¹	7.36 (53.67)	8.00 (63.58)	4.70 (21.58)	5.54 (30.17)	3.01 (8.58)	4.20 (17.17)	4.44 (19.25)	5.02 (24.67)
T ₆ : Pendimethalin 750 g ha ⁻¹	7.51 (55.97)	8.03 (63.92)	4.87 (23.25)	5.69 (31.83)	3.17 (9.58)	4.40 (18.83)	4.52 (19.92)	5.19 (26.42)
T ₇ : Hand weeding	3.50 (11.75)	4.35 (18.50)	3.28 (10.33)	4.01 (15.58)	1.50 (1.75)	2.25 (4.58)	2.80 (7.33)	3.29 (10.33)
T ₈ : Weedy Check	9.61 (91.92)	10.09 (101.42)	5.44 (29.17)	6.42 (40.67)	4.32 (18.17)	5.49 (29.67)	5.45 (29.17)	5.89 (34.17)
SEm±	0.27	0.33	0.21	0.26	0.17	0.21	0.19	0.25
CD at 5%	0.81	1.01	0.65	0.80	0.51	0.65	0.58	0.75

Table 2. Effect of different weed control treatments on weed density (no./m²) at 30 and 60 DAS (broad-leaved & other weeds)

Treatments	<i>Commelina communis</i>		<i>Phyllanthus niruri</i>		<i>Eclipta alba</i>		Other weeds	
	30 DAS	60 DAS	30 DAS	30 DAS	30 DAS	60 DAS	30 DAS	60 DAS
T ₁ : Mesotrione 250 g ha ⁻¹	4.20 (17.17)	5.00 (24.33)	2.98 (8.42)	2.98 (8.42)	2.98 (8.42)	4.02 (15.75)	2.59 (6.22)	3.49 (11.75)
T ₂ : Mesotrione 300 g ha ⁻¹	4.02 (15.75)	4.65 (20.42)	2.52 (5.92)	2.52 (5.92)	2.52 (5.92)	3.90 (14.75)	2.04 (3.72)	3.35 (10.75)
T ₃ : Mesotrione 350 g ha ⁻¹	3.59 (12.42)	4.27 (17.75)	2.22 (4.42)	2.22 (4.42)	2.22 (4.42)	3.61 (12.58)	1.79 (2.70)	3.00 (8.58)
T ₄ : Tembotrione 286 g ha ⁻¹	4.08 (16.17)	4.43 (20.58)	2.83 (7.50)	2.83 (7.50)	2.83 (7.50)	3.98 (15.33)	2.40 (5.30)	3.44 (11.33)
T ₅ : Atrazine 1000 g ha ⁻¹	4.37 (18.58)	5.09 (24.83)	3.00 (8.50)	3.00 (8.50)	3.00 (8.50)	4.04 (15.83)	2.61 (6.30)	3.50 (11.83)
T ₆ : Pendimethalin 750 g ha ⁻¹	4.55 (20.25)	5.06 (25.83)	3.05 (8.83)	3.05 (8.83)	3.05 (8.83)	4.15 (16.75)	2.67 (6.63)	3.60 (12.75)
T ₇ : Hand weeding	3.04 (8.75)	3.57 (11.92)	1.67 (2.30)	1.67 (2.30)	1.67 (2.30)	2.08 (3.85)	1.65 (2.22)	2.04 (3.67)
T ₈ : Weedy Check	5.38 (28.50)	6.58 (42.03)	4.86 (23.17)	4.86 (23.17)	4.86 (23.17)	5.90 (34.33)	4.54 (20.13)	5.66 (31.55)
SEm±	0.16	0.20	0.14	0.14	0.14	0.17	0.12	0.14
CD at 5%	0.47	0.61	0.42	0.42	0.42	0.52	0.38	0.44

Table 3. Effect of different treatments on dry weight of weeds (g/m²) at 30 and 60 DAS (grassy weeds and sedges)

Treatments	<i>Echinochloa colona</i>		<i>Digitaria sanguinalis</i>		<i>Eleusine indica</i>		<i>Cyperus rotundus</i>	
	30 DAS	60 DAS	30 DAS	60 DAS	30 DAS	60 DAS	30 DAS	60 DAS
T ₁ : Mesotrione 250 g ha ⁻¹	5.07 (25.20)	5.41 (28.80)	4.14 (16.60)	4.39 (18.75)	3.18 (9.60)	3.53 (11.95)	3.61 (12.53)	3.76 (13.67)

T ₂ : Mesotrione 300 g ha ⁻¹	3.78 (13.82)	3.96 (15.21)	4.01 (15.59)	4.24 (17.44)	3.02 (8.59)	3.34 (10.64)	2.96 (8.25)	3.11 (9.17)
T ₃ : Mesotrione 350 g ha ⁻¹	2.74 (7.00)	3.59 (12.42)	3.93 (14.92)	4.19 (17.08)	2.95 (8.20)	3.28 (10.28)	2.63 (6.43)	3.06 (8.83)
T ₄ : Tembotrione 286 g ha ⁻¹	3.90 (14.73)	4.43 (19.08)	4.08 (16.15)	4.32 (18.16)	3.11 (9.15)	3.44 (11.36)	3.57 (12.24)	3.56 (12.17)
T ₅ : Atrazine 1000 g ha ⁻¹	5.19 (26.43)	5.44 (29.05)	4.94 (23.93)	5.02 (24.70)	3.97 (15.30)	4.18 (16.93)	3.62 (12.63)	3.81 (14.00)
T ₆ : Pendimethalin 750 g ha ⁻¹	5.38 (28.45)	5.86 (33.87)	4.95 (24.02)	5.11 (25.63)	4.16 (16.83)	4.42 (19.00)	3.91 (14.77)	4.45 (19.27)
T ₇ : Hand weeding	1.49 (1.73)	3.13 (9.28)	1.33 (1.27)	1.94 (3.26)	1.21 (0.97)	1.26 (1.10)	0.97 (0.45)	1.75 (2.57)
T ₈ : Weedy Check	8.12 (65.38)	9.22 (84.50)	5.17 (26.23)	5.40 (28.63)	4.43 (19.14)	4.78 (22.35)	4.98 (24.30)	5.54 (30.20)
SEM±	0.31	0.33	0.20	0.31	0.26	0.15	0.13	0.16
CD at 5%	0.93	1.00	0.60	0.94	0.78	0.45	0.40	0.48

Table 4. Effect of different treatments on dry weight of weeds (g/m²) at 30 and 60 DAS (broad-leaved & other weeds)

Treatments	<i>Commelina communis</i>		<i>Phyllanthus niruri</i>		<i>Eclipta alba</i>		Other weeds	
	30 DAS	60 DAS	30 DAS	60 DAS	30 DAS	60 DAS	30 DAS	60 DAS
T ₁ : Mesotrione 250 g ha ⁻¹	4.43 (19.14)	5.06 (25.13)	4.15 (16.72)	4.73 (21.84)	3.54 (12.05)	4.16 (16.83)	3.25 (10.03)	3.74 (13.47)
T ₂ : Mesotrione 300 g ha ⁻¹	3.39 (11.00)	4.17 (16.93)	3.21 (9.83)	3.60 (12.47)	3.35 (10.74)	3.59 (12.41)	2.58 (6.17)	2.83 (7.50)
T ₃ : Mesotrione 350 g ha ⁻¹	2.86 (7.66)	3.57 (12.25)	2.74 (7.02)	3.34 (10.66)	2.98 (8.40)	3.30 (10.38)	2.28 (4.70)	2.47 (5.58)
T ₄ : Tembotrione 286 g ha ⁻¹	4.18 (16.93)	4.44 (19.25)	3.92 (14.85)	4.16 (16.83)	3.35 (10.74)	3.91 (14.77)	3.11 (9.18)	3.59 (12.36)
T ₅ : Atrazine 1000 g ha ⁻¹	5.06 (25.13)	5.24 (27.00)	4.22 (17.33)	5.09 (25.38)	3.73 (13.40)	4.20 (17.16)	3.36 (10.80)	3.78 (13.77)
T ₆ : Pendimethalin 750 g ha ⁻¹	5.24 (27.00)	5.87 (34.00)	4.42 (19.06)	5.21 (26.60)	3.83 (14.19)	4.22 (17.33)	3.44 (11.32)	3.82 (14.10)
T ₇ : Hand weeding	0.99 (0.48)	3.37 (10.84)	0.97 (0.43)	3.34 (10.65)	1.17 (0.87)	1.59 (2.03)	1.49 (1.73)	1.94 (3.26)
T ₈ : Weedy Check	5.83 (33.50)	6.51 (41.91)	5.24 (27.00)	5.75 (32.53)	4.68 (21.40)	5.28 (27.33)	3.89 (14.64)	4.20 (17.11)
SEM±	0.14	0.18	0.16	0.15	0.13	0.09	0.11	0.11
CD at 5%	0.41	0.55	0.49	0.45	0.40	0.26	0.34	0.34

3.3 Weed control efficiency

Weed control efficiency (WCE) of various treatments was determined based on relative weed biomass recorded under weedy check and other treatments at 30 and 60 DAS. The weed control efficiency significantly varied due to different weed control treatments (Table 5). The weed control efficiency (WCE) had significant inverse relationship with dry matter production by weeds. The weed control efficiency was maximum with hand weeding twice (96.58% and 84.89%) at 30 and 60 DAS, respectively. It was due to

Comment [A24]: No LSD values and superscripts to show similarity or differences

Comment [A25]: As in A 23

the production of minimum dry matter of all the associated weeds. All herbicidal treatments reduced weed growth, but the post emergence application of mesotrione 350 g ha⁻¹ had greatest amount of suppression of weeds, *fb* mesotrione 300 g ha⁻¹ *fb* tembotrione 286 g ha⁻¹ resulting in highest weed control efficiency. The results are in conformation with [29, 30].

3.4 Grain and stover yields

Weed control treatments also significantly influenced the maize grain yield and stover yield (Table 5). The uncontrolled weed competition during the crop duration in weedy check resulted in the lowest grain yield and stover yield of 1.39 and 16.25 t ha⁻¹ respectively. It was due to severe weed and crop competition from germination to harvest, leading to poor growth parameters, yield attributing characteristics and finally the grain yield [31, 32]. Except, application of pendimethalin 750 g ha⁻¹, the rest of herbicide treatments (atrazine 1000 g ha⁻¹, mesotrione 250 g ha⁻¹, mesotrione 300 g ha⁻¹, mesotrione 350 g ha⁻¹ and tembotrione 286 g ha⁻¹) produced significantly higher grain yield and stover yield compared with untreated weedy check. However, higher grain and stover yield were recorded with the application of mesotrione 350 g ha⁻¹ (2.44 and 21.80 t ha⁻¹). Among all the weed control treatments, highest grain and stover yield were observed under hand weeding treatment. The crop under weed free plots attained lush growth due to elimination of weeds from inter and intra row spaces besides better aeration due to manipulation of surface soil and thus, more space, water, light and nutrients were available for the better growth and development, which resulted into superior yield attributes and development, and consequently the highest yield. The research findings are in conformity with [33, 34].

Comment [A26]: How?

Comment [A27]: How?

Table 5. Effect of different weed control treatments on weed control efficiency, grain yield and stover yield

Treatments	Weed control efficiency (%)		Grain yield (t/ha)	Stover yield (t/ha)
	30 DAS	60 DAS		
T ₁ : Mesotrione 250 g ha ⁻¹	47.37	47.13	1.83	19.70
T ₂ : Mesotrione 300 g ha ⁻¹	63.73	64.24	2.18	20.95
T ₃ : Mesotrione 350 g ha ⁻¹	72.23	69.25	2.44	21.80
T ₄ : Tembotrione 286 g ha ⁻¹	55.10	56.43	2.07	20.61
T ₅ : Atrazine 1000 g ha ⁻¹	37.40	40.96	1.76	18.40
T ₆ : Pendimethalin 750 g ha ⁻¹	32.80	33.30	1.63	17.12
T ₇ : Hand weeding	96.58	84.89	2.80	22.52

T ₈ : Weedy Check	0.00	0.00	1.39	16.25
SEm±	-	-	0.04	0.90
CD at 5%	-	-	0.13	2.60

Comment [A28]: No statistical analysis and ISD for comparison of means

3.5 Weed index

Weed index measures the reduction in crop yield due to weed competition as against two hand weeding treatment and is expressed in percentage (**Figure 2**). Weed index was significantly affected by different weed control treatments. The data revealed that maximum reduction in yield (50.22%) occurred in weedy check pots where weeds were not controlled throughout the crop season. This might be due to the competition of weeds with the crop for the useful resources which may have resulted in suppressed the growth and development of the crop. The post emergence application of mesotrione 350 g ha⁻¹ resulted in the minimum weed index (12.62%) among the weed control treatments due to the effective and timely control of weeds in the crop. [35, 36] also reported a similar findings.




Figure 2. Effect of weed control treatments on weed index in maize

4. CONCLUSION

From the one-year experimental findings, it can be concluded that the application of mesotrione 350 g ha⁻¹ as post emergence application effectively controlled the diverse weed flora of maize with highest weed control efficiency and found to be most promising weed control treatment for obtaining highest grain and stover yield.

Comment [A29]: The conclusion should give a summary of ALL the research results
This should have been more than one statement

REFERENCES

1. Arvadiya LK, Raj VC, Patel TU, Arvadiya MK and Patel AM. Effect of plant population and weed management practices on productivity of sweet corn. *Indian Journal of Weed Science*. 2012; 44(3): 167–171.
2. Kumar A, Behera UK, Dhar S, Shukla L, Bhatiya A, Meena MC, Gupta G and Singh RK. Effect of tillage, crop residues and phosphorus management practices on the productivity and profitability of maize cultivation under Inceptisols of IGPs. *Indian Journal of Agricultural Sciences*. 2018; 85(2): 182–188.
3. DOAC&FW. 2019. *Agriculture Statistics at a Glance*. Ministry of Agriculture and Farmers Welfare, Government of India, New Delhi, p. 78.
4. Anonymous. 2019. *Annual Report 2021-2022*, Department of Agriculture and Cooperation, Ministry of Agriculture, Government of Madhya Pradesh.
5. Verma B, Bhan M, Jha AK, Khatoon S, Raghuvanshi M, Bhayal L, Sahu MP, Patel Rajendra, Singh Vikash. Weeds of direct-seeded rice influenced by herbicide mixture. *Pharma Innovation*. 2022;11(2):1080-1082.
6. Patel VJ, Upadhyay PN, Patel JB and Meisuriya MI. Effect of herbicide mixture on weeds in Kharif maize (*Zea mays* L.) under middle Gujarat conditions. *Indian Journal of Weed Science*. 2006; 38(1 and 2): 54-57.
7. Patel Raghav, Jha AK, Verma Badal, Kumbhare Rahul, Singh Richa. Bio- efficacy of pinoxaden as post-emergence herbicide against weeds in wheat crop. *Pollution research*. 2023;42(1):115-117.
8. Sahu, M. P., Kewat, M. L., Jha, A. K., Sondhia, S., Choudhary, V. K., Jain, N., ... & Verma, B. Weed prevalence, root nodulation and chickpea productivity influenced by weed management and crop residue mulch. *AMA, Agricultural Mechanization in Asia, Africa and Latin America*. 2022;53(6):8511-8521.
9. Shiv Swati, Agrawal S.B., Verma Badal, Yadav Pushpendra Singh, Singh Richa, Porwal Muskan, Sisodiya Jirtendra and Patel Raghav. Weed dynamics and productivity of chickpea as affected by weed management practices. *Pollution Research*. 2023; 42 (2):21-24.
10. Yadav PS, Kewat ML, Jha AK, Hemalatha K, Verma B. Effect of sowing management and herbicides on the weed dynamics of berseem (*Trifolium alexandrinum*). *Pharma Innovation*. 2023; 12(2): 2845-2848.
11. Tanisha Nirala, A.K. Jha, Badal Verma, Pushpendra Singh Yadav, Mahendra Anjna and Lakhan Bhalse. Bio efficacy of Pinoxaden on Weed Flora and Yield of Wheat (*Triticum aestivum* L.). *Biological Forum – An International Journal*. 2022; 14(4): 558-561.
12. Verma B, Bhan M, Jha AK, Singh V, Patel R, Sahu MP, Kumar V. Weed management in direct-seeded rice through herbicidal mixtures under diverse agroecosystems. *AMA, Agricultural Mechanization in Asia, Africa and Latin America*. 2022;53(4):7299- 7306.
13. Sahu, V., Kewat, M. L., Verma, B., Singh, R., Jha, A. K., Sahu, M. P., & Porwal, M. Effect of carfentrazone-ethyl on weed flora, growth and productivity in wheat. *The Pharma Innovation Journal* 2023; 12(3): 3621-3624.

14. Chikoye D, Akobundu IO, Isichei AO and Ekeleme F. Cover crops reduce weed seedbanks in maize-cassava systems in Southwestern Nigeria. *Weed Science*. 2004; 51(5): 774-780.
15. Jha AK, Yadav PS, Shrivastava A, Upadhyay AK, Sekhawat LS, Verma B, Sahu MP. Effect of nutrient management practices on productivity of perennial grasses under high moisture condition. *AMA, Agricultural Mechanization in Asia, Africa and Latin America*. 2023; 54(3): 12283-12288.
16. Sunitha N, Reddy PM and Sadhineni M. Effect of cultural manipulation and weed management practices on weed dynamics and performance of sweet corn (*Zea mays* L.). *Indian Journal of Weed Science*. 2010; 47(3&4): 184-188.
17. Armel GR, Wilson HP, Richardson RJ and Hines TE. Mesotrione combinations for postemergence control of Horsenettle (*Solanum carolinense*) in corn (*Zea mays*). *Weed Technology*. 2003; 17(1): 65-72.
18. Zhang J, Zheng Li, Jäck O, Yana D, Zhang Z, Gerhards R and Ni H. Efficacy of four post-emergence herbicides applied at reduced doses on weeds in summer maize (*Zea mays* L.) fields in North China Plain. *Crop Protection*. 2013; 52: 26-32.
19. Yang HH, Wang LQ, Xie ZJ, Yu Q, Liu G, Tan HR. The tyrosine degradation gene hpdD is transcriptionally activated by HpdA and repressed by HpdR in *Streptomyces coelicolor*, while hpdA is negatively autoregulated and repressed by HpdR. *Molecular Microbiology*. 2007; 65(4): 1064-1077.
20. Guo WL, Feng L, Cui Y, Wu DD, Zhang TJ, Zhang C, et al. Weed control effect of 30% nicosulfuro+atrazine+mesotrione oil dispersion and the safety to sweet corn in field. *Journal of Maize Sciences*. 2020; 28(4): 178-183. <https://doi.org/10.13597/j.cnki.maize.science.20200426> (in Chinese).
21. Jha AK, Shrivastva Arti, Raguvanshi NS. Effect of weed control practices on the fodder and seed productivity of Berseem under irrigated condition of Madhya Pradesh. *Range management & Agroforestry*. 2014; 35(1): 61-65.
22. Kantwa SR, Agrawal RK, Jha A, Pathan SH, Patil SD, Choudhary M. Effect of different herbicides on weed control efficiency, fodder and seed yields of berseem (*Trifolium alexandrinum* L.) in central India. *Range Management and Agroforestry*. 2019; 40(2): 323-328.
23. Kumar B, Prasad S, Mandal D and Kumar R. Influence of integrated weed management practices on weed dynamics, productivity and nutrient uptake of rabi maize (*Zea mays* L.). *International Journal of Current Microbiology and Applied Sciences*. 2017; 6(4): 1431-1440.
24. Verma S, Verma DK, Giri SP, Singh K, Singh RB and Panday A. Response of chemical weed management in maize and cow intercropping system grown for quality fodder in Eastern Uttar Pradesh. *Indian Journal of Forestry and Crop Improvement*. 2015; 6: 105-119.
25. Idziak R, Skrzypczak W, Waligora H, and Woznica Z. The effect of mesotrione applied with adjuvants on weed control efficacy and forage sorghum tolerance. *Turkish Journal of Agriculture and Forestry*. 2013; 37(3): 265-270.
26. Soltani N, Kaastra AC, Swanton CJ, and Sikkema PH. Efficacy of topramezone and mesotrione for the control of annual grasses. *International Research Journal of Agricultural Science and Soil Science*. 2012; 2(1): 46-50.
27. Elmore MT, Brosnan JT, Kopsell DA and Breeden GK. Nitrogen-enhanced efficacy of mesotrione and topramezone for smooth crabgrass control. *Weed Science*. 2012; 60(3): 480-485.
28. Sharma P, Duary B and Singh R. Tank mix application of tembotrione and atrazine to reduce weed growth and increase productivity of maize. *Indian Journal of Weed Science*. 2018; 50(3): 305-308.
29. Kaur M, Singh K, and Singh H. Varying weed management treatments impact on weeds and fodder yield relationship in fodder maize. *Indian Journal of Weed Science*. 2023; 55(1): 103-106.
30. Ahmed MAA and Susheela R. Weed management studies in Kharif maize. *Journal of Research ANGRAU*. 2012; 40(3): 121- 123.
31. Walia US, Singh S and Singh B. Integrated control of hardy weeds in maize (*Zea mays* L.). *Indian Journal of Weed Science*. 2007; 39(1&2): 17-20.
32. Swetha K, Madhavi M, Pratibha G and Ramprakash T. Weed management with new generation herbicides in maize. *Indian Journal of Weed Science*. 2015; 47(4): 432-433.
33. Sanodiya P, Jha AK, and Shrivastava A. Effect of integrated weed management on seed yield of fodder maize. *Indian Journal of Weed Science*. 2013; 45(3): 214-216.

34. Barod NK, Dhar S and Kumar A. Effect of nutrient sources and weed control methods on yield and economics of baby corn (*Zea mays* L.). Indian Journal of Agronomy. 2012; 57(1): 96-99.
35. Chhokar RS, Sharma RK, Gill SC and Singh RK. Mesotrione and Atrazine combination to control diverse weed flora in maize. Indian Journal of Weed Science. 2019; 51(2): 145–150.
36. Dharam Bir Yadhav, Ashok Yadhav, Punia SS, Anil Duhan. Temobotrione for post-emergence control of complex weed flora in maize. Indian Journal of Weed Science. 2018; 50(2): 133-136.

UNDER PEER REVIEW