

PROXIMATE AND ANTINUTRIENT COMPOSITION OF  
*Hura crepitans* (SANDBOX TREE) SEEDS

ABSTRACT

Analysis of proximate and antinutrient composition of *Hura crepitans* (Sandbox tree) seeds obtained from campus of Akwa Ibom State Polytechnic Ikot Osurua, Ikot Ekpene were carried out using standard analytical procedures. The result of the proximate composition analysis showed that *H.crepitans* seeds contain moisture content ( $13.875 \pm 0.176\%$ ), crude fat ( $36.750 \pm 1.060\%$ ), crude fibre ( $0.5 \pm 0.000\%$ ), crude protein ( $36.625 \pm 0.884\%$ ), ash content ( $0.25 \pm 0.000\%$ ) and carbohydrate ( $25.875 \pm 0.200\%$ ). The result of the antinutrient analysis showed that *H.crepitans* seeds contain phytic acid (phytate) ( $0.615 \pm 0.002\text{mg}/100\text{g}$ ), hydrogen cyanide ( $3.289 \pm 0.003\text{mg}/100\text{g}$ ), oxalate ( $1.386 \pm 0.001\text{mg}/100\text{g}$ ) and tannins ( $8.130 \pm 0.003\text{mg}/100\text{g}$ ). Tannins had the highest concentration while phytate had the lowest concentration of all the antinutrients analyzed for. The antinutrients can be removed during processing. The result of the proximate composition reveals high crude fat and protein contents of the seed of *H.crepitans*, showing that the seed is nutritional. Findings from this study therefore suggest intensified research on the seeds of *H.crepitans* so as to harness its potentials in the production of animal feed.

Keywords: Antinutrient, proximate composition, Sandbox tree, seeds.

## Introduction

*Hura crepitans*, commonly referred to as sandbox tree (Fig.1) is a tropical plant of Euphorbiaceae family (Abdulkadir *et al.*, 2013). The tree is native to Tropical America (Owojuyigbe *et al.*, 2020), and is traditionally grown as shade tree (Ola and Jekayinfa, 2015), and also has been found to be useful in many ethnomedicinal applications such as its use in the treatment of leprosy and as antimicrobial among others (Owojuyigbe *et al.*, 2020). Sandbox tree can grow above 40 metres (Azubuiké *et al.*, 2016).



Fig.1: Sandbox Tree

It has many dark, pointed spines and smooth brown bark. Because of these spines, it is normally called “Monkey no-climb” (Oyeleke *et al.*, 2012).

The fruits of this tree resemble small pumpkin pods with 16 seeds enclosed and because the fruits explode on maturation, it is usually called dynamite tree (Okolie *et al.*, 2012). The seeds are small and brown in colour with an average diameter of about 1-5cm (Mohammed, 2004).

*H. crepitans* seed has been found to contain glucosamine, lectin that has mitogenic and haemagglutinating properties (Owojuyigbe *et al.*, 2020). The seeds while still green have strong purgative effect. The juice secreted by sandbox tree had been used to poison darts (Azubuikwe *et al.*, 2016).

Fowomola and Akindahunsi (2007) observed that the seeds of *H. crepitans* cause burning sensation to the throat and induces vomiting, but noted that it is a good protein source for animal feeds when properly treated.

Although sandbox trees are being used as shade in most part of the world (Idowu *et al.*, 2012), the tree is one of the underutilized trees in Nigeria (Otoikhian *et al.*, 2016). Hence, this study was carried out to determine the proximate and antinutritional composition of the seeds of *H. crepitans*.

## **Materials and Methods**

### ***Sample Collection and Preparation***

The seeds of *Hura crepitans* (Sandbox tree seeds) were collected from matured dry pods (fruits) of *Hura crepitans* tree around the campus of Akwa Ibom State Polytechnic, Ikot Osurua, in Ikot Ekpene Local Government Area of Akwa Ibom State.

The seeds were decorticated and pounded into powdered form using mortar and pestle, and the analysis was carried out in the Chemistry laboratory of Akwa Ibom State Polytechnic, Ikot Osurua.

### ***Proximate Analysis***

The recommended methods of AOAC (2000) were used for the determination of moisture, ash, crude fat, crude fibre and crude protein contents while carbohydrate

was calculated by difference by subtracting total sum of crude protein, crude fibre, crude fat, moisture and ash content from 100% dry weight sample.

### ***Antinutrient Analysis:***

The method described by Ola and Oboh (2000) was used for the determination of Phytate, while the methods described by AOAC (1990), Krishna and Ranjhan (1980) and Allen *et al.* (1974) were used to determine the levels of hydrocyanic acid, oxalate and tannins respectively.

### ***Results***

The result of the proximate composition of *Hura crepitans* seeds is shown in Table 1 while that of antinutrient analysis of *Hura crepitans* seeds is shown in Table 2.

Table 1: Proximate Composition of *Hura crepitans* Seeds.

Parameter	Percentage (%)
Moisture	13.875 ± 0.176
Crude fat	36.750 ± 1.060
Crude fibre	0.5 ± 0.000
Crude protein	36.625 ± 0.884
Ash	0.25 ± 0.000
Carbohydrate	25.875 ± 0.200

Mean of 2 Determinations ± S.D.

Table 2: Antinutrient Composition of *Hura crepitans* Seeds

Antinutrient	Composition (mg/100g)
Phytic acid	0.615 ± 0.002
Hydrogen cyanide	3.289 ± 0.003
Oxalate	1.386 ± 0.001
Tannins	8.130 ± 0.003

Mean of 2 Determinations ± S.D.

## ***Discussion***

The result of the proximate composition of *H. crepitans* seeds as shown in Table 1, revealed the moisture content to be  $13.875 \pm 0.176\%$ , which is higher than 3.10% reported by Abdulkadir *et al.* (2013), and that of Ige *et al.* (2019) which varied from 5.123 - 8.23%. The value was also higher than 7.89% reported by Nasir and Usman (2019) for the seeds of *Vigna membranacea*. This high moisture content shows that the seeds of *H. crepitans* will not have a longer shelf-life. High moisture content has been implicated in food spoilage (Okezie *et al.*, 2017).

The crude fat was found to be  $36.750 \pm 1.060\%$  which was lower than 53.61% reported by Okolie *et al.* (2012). This value was within the range reported by Auta and Anwa (2007) for oil seeds from 18% in soya beans to 43% in groundnut oil. As such, *H. crepitans* seed can be seen as an oil seed.

Crude fibre content of the sandbox seeds was found to be  $0.5 \pm 0.000\%$ . This value is lower than 1.45% reported by Okolie *et al.* (2012). The crude protein content of the sample was found to be  $36.625 \pm 0.884\%$ . The value was higher than 22.36% reported by Okolie *et al.* (2012), but lower than  $25.16 \pm 0.22\%$  as reported by Fowomola and Akindahunsi (2007) and 10.50% reported by Akpabio *et al.*, (2012) for *Cassia hirsute* seeds. Oyeleke *et al.* (2012) reported  $25.76 \pm 0.04\%$  crude protein in *H. crepitans* seed and noted that high crude protein content of the seed suggests it to be a good protein source. Nwokenkwo *et al.* (2020) also suggested that the seeds of *H. crepitans* should be used in areas of food industries where protein (amino acids) is critically needed since it contains significant amount of essential amino acids.

Esonu *et al.* (2014) reported that sandbox seed meal could enhance the performance of broiler finishers of 10% dietary levels as the seeds contain amino acids (the building units for body protein) at levels comparable to soya bean and groundnut seeds.

The ash content was  $0.25 \pm 0.000\%$  which is lower than  $3.54 \pm 0.02\%$  obtained by Oyeleke *et al.* (2012) and 7.70% reported by Akpabio *et al.*, (2012) for *Cassia hirsute* seeds. This value was very small when compared to that of *Vigna membranacea* (25.67%) (Nasir and Usman, 2019). The carbohydrate content of the seed of *H. crepitans* was  $25.875 \pm 0.200\%$  which is higher than  $0.69 \pm 0.2\%$  reported by Abdulkadir *et al.* (2013).

### ***Antinutrients***

Antinutrients are substances from plants which can affect the availability of nutrients by interfering with metabolic processes (Okezie *et al.*, 2017). Nutritional deficiencies, headaches, rashes and nausea are some of the symptoms of large amount of antinutrients in the body (Essack *et al.*, 2017). Levels of antinutrient can be reduced by traditional processing methods such as cooking (Sha'a *et al.*, 2019). The result of the antinutrient composition of *H. crepitans* seeds are presented in Table 2. From the result, phytic acid was found to be  $0.615 \pm 0.002$  mg/100g, this value is lower than  $20.28 \pm 0.90$  mg/100g obtained by Hassan *et al.* (2018). Phytic acid can bind calcium, zinc, magnesium, manganese and iron to form indigestible complexes and thereby reduce the bioavailability of the elements for absorption (Umaru *et al.*, 2007). Hydrogen cyanide was found to be  $3.289 \pm 0.003$  mg/100g. This value is higher than 0.18 mg/100g of hydrocyanic acid reported by Hassan *et al.* (2018). Oxalate was found to be  $1.386 \pm 0.001$  mg/100g. This value is higher than 0.017 mg/100g obtained by Hassan *et al.* (2018). Regular consumption food with excess amount of oxalic acid may results in nutritional deficiencies and severe irritation of the living of the gut (Gemedé and Ratta, 2014).

Tannins was found to be  $8.130 \pm 0.003$  mg/100g which is higher than 0.43mg/100g reported by Hassan *et al.* (2018). Tannins decreased protein digestibility in humans and animals (Gemedé and Ratta, 2014). Certain studies have indicated tannins for healing of wounds (Ukpong *et al.*, 2014; Ambreen and Mirza, 2020; Eneh and Ukpong, 2021), treatment of urinary tract infection, diarrhoea and dysentery (Fahey, 2005; Akinpelu and Onakoya, 2006; Godfrey, 2015), and in the production and storage of some foods in order to increase the products' shelf life because of the anti-microbial activity of tannins (Bajaj, 1999).

### ***Conclusion***

The results of this work show that sandbox tree seeds possess all the analysed antinutrients which can even be removed during processing. The proximate composition results reveal high crude fat and protein contents of the seeds of *H. crepitans*, showing that the seed is nutritional. Findings from this study therefore suggest intensified research on the seeds of *H. crepitans* so as to harness its

potentials in the production of animal feed.

## REFERENCES

- Abdulkadir, M. N., Amoo, I. A. and Adesina, A. O. (2013). Chemical composition of *Hura crepitans* seeds and antimicrobial activities of its oil. *International Journal of Sciences and Research*, 2(3):440-445.
- Akinpelu, D. A. and Onakoya, T. M. (2006). Antimicrobial Activities of Medicinal Plants used in Folklore Remedies in South-western. *African Journal of Biotechnology*, 5, 1078-1081.
- Akpabio, U.D., Wilson, L. A., Akpakpan, A. E. and Obot, I. B. (2012). Phytochemical screening and proximate composition of *cassia hirsute* seeds. *Elixir Appl. Chem.* 47 8704-8707
- Allen, E. S., Grimshaw, H. M., Parkinson, J. A. and Quarmby, C. (1974). Chemical Analysis of Ecological Materials Blackwell Scientific Publication, London. Pp. 52-62.
- Ambreen, M. and Mirza, S. A. (2020). Evaluation of Anti-inflammatory and Wound Healing Potential of Tannins Isolated from Leaf callus Cultures of *Achyranthes aspera* and *Ocimum basilicum*. *Pak. J. Pharm. Sci.* 33(1(supplementary)), 361-369.
- Association of Official Analytical Chemist (AOAC) (1990). Official Methods of analysis, 14<sup>th</sup> Edition. *Association of Analytical Chemists Washington D. C.* pp. 1112-1114.
- Association of Official Analytical Chemist (AOAC) (2000). Official methods of analysis of the AOAC 17<sup>th</sup> Horwitz W. ed. AOAC. *International Published. Maryland USA.*
- Auta, J. and Anwa, P. (2007). Preliminary Studies on *Albizia lebbek* Seeds - Proximate Analysis Phytochemical Screening. *Medwell Research Journal of Biological Sciences*, 2(1), 33-35.
- Azubuikwe, A. E., Iheanyichukwu, E., Stanley, O. and Ogwo, J. I. (2016). Comparative Antihypertensive Properties of Aqueous Extracts of Leaves, Stem Bark and Roots of *Hura crepitans* (L) in Adrenaline induced Hypertensive Albino Rats. *International Journal of Technical Research and Applications* 4(1):185-193.
- Bajaj, Y. S. (1999). Medicinal and Aromatic Plants V. Biotechnology in Agriculture and Forestry Berlin, Heidelberg. Springer, pp. 24.
- Eneh, G.D.O. and Ukpong, E. G. (2021). Comparative Study of Phytochemical Analysis of Two Generic Species of Cocoyam (*Xanthosoma sagittifolium* and *Colocasia esculenta*). In: Albayrak, H. (ed.) *Full Text Book, Cukurova 7th International Scientific Research Conference, 7th - 8th September, 2021.* Adana: IKSAD Global Publications p.1182

- Esonu, B. O., Ozeudu, E., Emenalom, O. O., Nnaji, C., and Onyeikegbulem, I. K. (2014). Nutritional value of sandbox (*Hura crepitans*) seed meal for Broiler Finisher Birds. *Journal of Natural Sciences Research*. Pp. 4(23):95-99.
- Essack, H., Odhar, B., and Mellen, J. J. (2017). Screening of Traditional South African Leafy Vegetables for Selected Anti-nutrient Factors before and after Processing. *Food Sci. Technol.* 3,462-1.
- Fahey, J. W. (2005). *Moringa oleifera*: A Review of the Medicinal Evidence for its Nutritional, Therapeutic and Prophylactic Properties. *Tree of Life Journal*, 15, 1-5.
- Fowomola, M. A., and Akindahunsi, A. A. (2007). Nutritional Quality of Sandbox Tree (*Hura crepitans* Linn.). *Journal of Medicinal Food* 10(1), 159-164.
- Genede, H. F. and Ratta, N. (2014). Antinutritional factors in plant foods: Potential Health Benefits and Adverse Effects. *International Journal of Nutrition and Food Science*, 3(4), 284-289.
- Godfrey, N. E. (2015). Proximate and Phytochemical Composition of the Pulp of *Tetrapleura tetraptera* Fruits Consumed in Abakaliki, Nigeria. *International Journal of Engineering Research & Technology*, 4(6), 1286-1294.
- Hassan, L. G., Sokoto, A. M., Ngaski, M. A. and Anka, S. A. (2018). Nutritional and Antinutritional analysis of *Hura crepitans* seeds cultivated in Sokoto North L.G.A., North-North-Western Nigeria. *Bayero Journal of Pure and Applied Sciences* 11(1), 126-130.
- Iidowu, D. O., Abegunrin, T. P., Ola, F. A., Adediran, A. A. and Olaniran, J. A. (2012). Measurement of some Engineering Properties of Sandbox seeds (*Hura crepitans*). *Agriculture and Biology Journal of North America*, 3(8), 318-325.
- Ige, M. M., Gbadamosi, S. O. and Solam, O. I. (2019). Proximate composition and functional properties of sandbox seeds as influenced by processing methods. *The Journal of Science*, 21(1), 129-144.
- Krishna, G. and Ranjhan, S. K. (1980). Laboratory manual for nutritional research. Vikas Publishing House PVT Ltd, Gaziabad India. Pp. 76-89.
- Mohammed, A. A. (2004). Chemical composition and characteristic of pumpkin (*curcubita maxima*) seed kernels, Res. Bult No. 129, 5-8.
- Nasir, A. and Usman, M. I. (2019). Proximate analysis and Elemental composition of

Vigna membranacea seeds. Food Science and Quality Management, 91, 65-68.

- Nwokenkwo, E. C., Nwosu, J. N., Onuegbu, N. C., Olawuni, I. A. and Ofoedum, A. F. (2020). Evaluation of the Antinutrients, Amino Acid Profile and Physicochemical Properties of *Hura crepitans* seeds. *Archives of Current Research International*, 2(5), 1-17.
- Okezie, E., Ubogu, A. E., Odungide, A. A. and Atani, C. S. (2017). Proximate, Antinutritional and Mineral Estimation of some selected consumed Green Vegetables in Afaha Eket, Akwa Ibom State, Nigeria. *American Journal of Food Science and Technology*, 5(5), 182-191.
- Okolie, P. N., Uaboi-Egbenni, P. O. and Ajekwene, A. E. (2012). Extraction and Quality Evaluation of Sandbox Tree Seed (*Hura crepitans*) oil. *World Journal of Agricultural Sciences* 8(4), 359-365.
- Ola, F. A. and Jekayinfa, S. O. (2015). Pyrolysis of Sandbox (*Hura crepitans*) shell: Effect of Pyrolysis parameters on biochar yield. *Res. Agr. Eng.* 61, 170-176.
- Ola, F. L., and Oboh, G. (2000). Food value of two Nigerian Edible Mushrooms (Termitomyces Stratus and termitomyces robustus). *The J. Technol. Sci.* 4, 1-3.
- Otoikhian, S. K., Aluyor, E. O. and Audu, T. O. K. (2016). Mechanical Extraction and Fuel Properties Evaluation of *Hura crepitans* seed oil. *Chemical Technology: An Indian Journal*, 11(6), 1-11.
- Owojuyigbe, O. S., Firempong, C. K., Larbie, C., Komlaga, G. and Emikpe, B. O. (2020). Hepatoprotective potential of *Hura crepitans* L. A Review of Ethnomedical, Phytochemical and Pharmacological Studies. *Journal of Complementary and Alternative Medical Research*, 8(2), 1-10.
- Oyeleke, G. O., Olayiwola, O. A. and Latona, D. F. (2012). Chemical Examination of Sandbox (*Hura crepitans*) seed: Proximate, Elemental and Fatty Acid Profile. *IOSR Journal of Applied Chemistry (IOSRJAC)*, 1(2), 10-13.
- Sha'a, K. K., Clarkson, G. P. and Artimas, S. P. (2019). Phytochemical Analysis, Proximate Composition and Antinutritional Factors of Corchorus oliterius plant. *International Journal of Biological and Chemical Sciences*, 13(4), 2147- 2157.
- Ukpong, I. J., Abasiokong, B. O. and Etuk, B. A. (2014). Phytochemical Screening and Mineral Elements Composition of *Xanthosoma sagittifolium* Inflorescence. *Asian Journal of Plant Science and Research*, 4(6), 32-35.
- Umaru, H. A., Adamu, R., Dahiru, D., and Nadro, M. (2007). Levels of Antinutritional factors in some wild Edible, Fruits of Northern Nigeria. *African Journal of Biotechnology* 6(16), 1935-1938.