

## Genetic Variability of Floral and Agronomic Characteristics that Influence Outcrossing Rate Percentage of Cytoplasmic Male Sterile Rice

### ABSTRACT

Developing high-yielding hybrid rice varieties is much needed to ensure food security to meet the requirement of the increasing population across the world. Cytoplasmic male genetic sterility system is a valuable technology for the exploitation of heterosis and production of high-yielding hybrid rice with better grain quality. Large-scale commercial exploitation of hybrid vigor in rice largely depends on the availability of good CMS (Cytoplasmic Male Sterile) lines. A total of eighteen traits of five CMS lines of rice were evaluated to study the variability and correlation between floral and other yield-contributing traits with outcrossing rate using Randomized Complete Block Design with three replications. High significant variation was observed among the genotypes for all the studied traits. High GCV (Genotypic Coefficient of Variance) and PCV (phenotypic Coefficient of Variance) coupled with high heritability and high genetic advance in percent of mean were observed for number of panicles per plant, filled grain, flag leaf breadth, stigma length and breadth, filament length and outcrossing rate which indicated the broad genetic base, less environmental influence, and these traits are under the control of additive gene action. character association and path analysis indicated that number of grains per panicle, filled grains per panicle, stigma length, and breadth possessed both positive significant correlation and positive direct effects on outcrossing rate therefore, selection for the traits mentioned above will certainly bring in the improvement of the outcrossing percentage of the lines, which in turn enhances the seed quantity of the hybrid.

**Keywords:** CMS, Outcrossing, Rice, Variability, Heritability

## 1. INTRODUCTION

"Rice is one of the most ancient grain crops which is used as a staple food in most of the countries for containing high amount of carbohydrates, low fat, and rich amounts of proteins, vitamins, and minerals" [1]. Daily we get 1329 kcal energy from rice which is about 60% of our total energy requirement. Its digestibility is also high than others cereal crop (99.70%) [2]. "Rice is a major source of dietary carbohydrates for more than half of the world's population" [3]. "In Bangladesh, about 75% (11.2 million ha) of total cropped land is covered by rice and more than 60% of man are engaged with it. Rice alone contributes around 10% of the GDP in this country" [4]. "Thus the single crop rice has multiple effect day to day daily life and the economy" [5]. "A modest estimate suggests that the demand for rice in Bangladesh will increase by over 80% in the next 20 years to feed the growing population" [6].

"It is predicted that the population in the world will increase continually to 8 billion in 2030. According to a document from FAOSTAT 2019" [7], "the annual loss of land for other uses is estimated to be about 10 to 35 million m<sup>2</sup>/year, half of the coming from the cropland. Facing the challenges of population growth and cropland reduction, it is obvious that the only way to solve the problem is to improve the yield of cereal crops. Fortunately, the importance of rice to feed the population is recognized by the United Nations. The year 2004 was declared the international year of rice, which reflects the importance of rice as a primary food source. It attracts a growing number of scientists around the world attempting to increase grain yield to keep pace with the demand of the burgeoning world population" [7]

Large-scale commercial exploitation of hybrid vigor in rice largely depends on the availability of good CMS lines and restorer lines. In recent times, cytoplasmic male sterility (CMS) technology has emerged as a promising approach for the dramatic enhancement of productivity. However, there is a difference between male sterility and cytoplasmic male sterility. Cytoplasmic male sterility is total or partial male sterility in plants as the result of specific nuclear and mitochondrial interactions whereas Male sterility is the failure of plants to produce functional anthers, pollen, or male gametes. CMS of rice with boro II cytoplasm is caused by a cytotoxic peptide and is restored by two related PPR motif genes via distinct modes of mRNA silencing. In the CMS system generally, three lines are essential that is A, B, and R lines. A line (CMS) is the cytoplasmic genetic male sterile line which is characterized by the absence of pollen grain, abortive anther, and stable sterility. The economic feasibility of hybrid rice seed production largely depends on the availability of good CMS lines with desirable traits, the knowledge of the desirable traits of CMS lines is quite essential in enhancing the seed yield of the potential rice hybrids. If parental lines of hybrids are locally adapted and adequate knowledge of agronomical and floral morphology of parental lines were gathered, it could lead self-sufficiently in F<sub>1</sub> seed production. So, the knowledge of agronomic and floral traits of A line is very much important. There is ample scope to study these characteristics influencing outcrossing rate of component lines of hybrid rice in the local environment. It would help in designing row ratio estimation for CMS multiplication and F<sub>1</sub> seed production and ultimately helps in building cost-effective seed production package development. The promising CMS lines are now being studied for their combining ability and use in breeding programs. Hence, the present study is undertaken to evaluate the locally

adaptable CMS lines for their traits to identify promising parental lines for developing heterotic hybrid rice. Therefore, this study was conducted to identify the most important characteristics of the breeding program by exploiting the genetic variation, heritability, and path analysis of outcrossing and related attributes of CMS lines of rice.

## 2. MATERIALS AND METHODS

The present investigation was carried out during *Aman*, 20018-19 at the experimental field of Bangabandhu Sheikh Mujibur Rahman Agricultural University (BSMRAU), Gazipur located at center of Madhupur Tract (24°29 N latitude and 90°26 E longitude) having an altitude of 8.2m from the sea level. The climate of the experimental site is subtropical in nature characterized by heavy rainfall during the months from June to September and scanty in winter with a gradual fall of temperature from the month of September. The experimental materials were collected from the Department of Genetics and Plant Breeding of Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur. The experimental material comprised five Cytoplasmic Male Sterile (A) lines those are IR 58A, IR 62A, IR 68888A, Gan 46A, and BRRI 1A. The experiment was conducted in a completely randomized block design with three replications. Thirty days old seedlings were transplanted in each row by adopting inter and intra-row spacing of 20 cm x 15cm. Fertilizer application and other cultural management practices were performed as per the recommendation of Aman Season. Standard crop management practice was done as and when necessary.

Five plants of each genotype in each replication were selected randomly to record different parameters. The parameters were collected for the characters *i. e* Days to 50% flowering, days to maturity, plant height (cm), panicle length (cm), number of tillers per hill, number of panicles per plant, number of grains per panicle, 1000 grain weight (gm), filled grain, flag leaf length (cm), flag leaf width (cm), anther length ( $\mu\text{m}$ ), anther breadth ( $\mu\text{m}$ ), stigma length ( $\mu\text{m}$ ), stigma breadth ( $\mu\text{m}$ ), filament length ( $\mu\text{m}$ ), pollen sterility (%), and outcrossing rate (%). Most of the traits were measured in the field except anther length, anther breadth, stigma length, stigma breadth, filament length, and pollen sterility which were measured in laboratory. Anther length, anther breadth, stigma length, stigma breadth, and filament length were measured with a microscope using an ocular micrometer (level 1) before anthesis.

For statistical analysis, analysis of variance, estimation of mean, range, standard error, coefficient of variation, Estimation of phenotypic and genotypic correlations, and Path coefficient analysis were considered. To test the significance of treatments critical difference was calculated [8]. Phenotypic and genotypic coefficients of variation (PCV and GCV) were analyzed according to [9] Heritability in broad sense and genetic advance was calculated according to methods specified by [10] and categorized according to [11]. Correlation coefficients were calculated for all quantitative character combinations at the phenotypic, genotypic, and environmental level by the formula set by [12]. Path coefficient analysis proposed by [13] and explained by [14] was used to analyze the direct and indirect impact of various traits to outcrossing.

### **3. RESULTS AND DISCUSSION**

#### **3.1 Analysis of variance**

The analysis of variance studied for 18 characters was furnished in Table 1a, 1b. The analysis of variance revealed significant differences for all the characters studied except panicle length and pollen sterility, which indicated that a wide range of variability present among the genotype. Previous studies also found significant variation in these traits [15].

Table 1a: Analysis of variance for different parameters of CMS (A) lines

Source of variation	d.f	50% F	DMT	PH (cm)	PL (cm)	NT/H	NP/P	NG/P	TGW (gm)	Filled grain
Replication	2	0.0	0.000	0.800	0.285	4.067	3.267	5.400	2.319	0.467
Genotype	4	33.0**	287.4**	146.267**	5.965NS	20.50*	49.833**	123.933**	30.47**	47.767**
Error	8	0.0	0.000	2.717	1.81	1.900	1.433	1.983	0.806	1.467

50% F = Days to 50% flowering, DMT = Days to maturity, PH = Plant height, PL = Panicle length, NT/H = Number of tiller per hill NP/P = Number of panicle per tiller, NG/P = Number of grain per panicle, TGW = Thousand grain weight

Table 1b: Analysis of variance for different parameters of CMS (A) lines

Source of variation	d.f	FLL (cm)	FLW (cm)	AL ( $\mu\text{m}$ )	AB ( $\mu\text{m}$ )	SL ( $\mu\text{m}$ )	SB ( $\mu\text{m}$ )	FL ( $\mu\text{m}$ )	PS (%)	OCR (%)
Replication	2	0.129	0.109	.9080	1.817	17.317	0.267	7.371	0.001	0.467
Genotype	4	62.16**	0.311*	298.25*	188.558**	13356.29**	1079.64**	41759.02**	1.5NS	21.5**
Error	8	0.308	0.027	25.350	2.983	15.192	6.142	16.614	0.001	0.800

FLL = Flag leaf length, FLW = flag leaf width, TW = Test weight, AL = Anther length, AB = Anther breadth, SL = Stigma length, SB = Stigma breadth, FL = Filament length, PS = Pollen sterility, OCR = Outcrossing rate

\* = Significant at 5% level, \*\* = Significant 1% level, d.f = Degrees of freedom

### 3.2 Genetic parameters for different characters of tested entries

The estimates of mean, range, phenotypic coefficients of variation (PCV), genotypic coefficients of variation (GCV), heritability ( $h^2$ ) and genetic advance as percent of mean (GAM) are presented in Table 2.

Table 2: Mean, range, genetic variability, heritability (broad sense), and genetic advance as percent of mean for outcrossing and its component traits for five CMS lines of rice

Sl. No	Character	Mean	Range	Coefficient of variation		Heritability (%)	GAM (%)
				PCV(%)	GCV(%)		
1	50% flowering	66.66	63-70	5.08	5.04	98.55	10.32
2	Days to maturity	82.93	73-96	11.98	11.97	99.89	24.65
3	Plant height (cm)	70.20	61.66-77.67	10.12	9.85	94.62	19.74
4	Panicle length (cm)	22.29	20.48-24.30	7.95	5.22	43.26	7.08
5	Number of tiller/hill	19.33	15.67-22.33	14.72	12.87	76.54	23.21
6	Number of panicle/plant	11.66	8.33-17.33	35.92	34.42	91.84	67.96
7	Number of grain/panicle	137.2	146.3-131.7	4.75	4.64	95.34	9.34
8	Thousand weight (gm)	17.97	13.47-22.30	18.19	17.49	92.46	34.65
9	Filled grain	17.86	14-22.33	52.25	49.93	91.32	98.30
10	Flag leaf length (cm)	24.47	20.52-32.51	18.48	18.35	98.52	37.52
11	Flag leaf breadth(cm)	1.24	0.83-1.53	27.97	24.68	77.80	44.84
12	Anther length( $\mu$ m)	395.86	384-407.33	2.71	2.40	78.46	4.39
13	Anther breadth( $\mu$ m)	74.36	66-83.16	10.82	10.57	95.39	21.27
14	Stigma length( $\mu$ m)	257.83	190.6-365.5	25.90	25.86	99.65	53.18
15	Stigma breadth ( $\mu$ m)	69.13	47.33-92.33	27.59	27.36	98.31	55.88
16	Filament length ( $\mu$ m)	518.69	413.1-667.5	22.75	22.74	99.88	46.81
17	Pollen sterility (%)	97.73	96-99	1.15	1.05	84.21	1.99
18	Outcrossing rate %	19.43	17.96-21.27	39.14	37.37	85.03	32.24

PCV = Phenotypic coefficient of variation, GCV = Genotypic coefficient of variation,

GAM = Genetic advance as percent of mean

### 3.3 Coefficient of variation (PCV, GCV)

Higher phenotypic co-efficient of variation, compared to genotypic co-efficient of variation was recorded for all the quantitative traits studied in the present investigation, indicating the influence of environment. Similar findings were reported earlier by Mamta Singh et al. (2007) [16]. However, high (>20%) phenotypic co-efficient of variation for number of panicles per plant, filled grains/panicle, flag leaf breadth, stigma length, stigma breadth, filament length, and outcrossing rate in the present research was noted to be close and essentially related with high genotypic co-efficient of variation,

indicating the negligible influence of environment and existence of high genetic variability for these traits. Hence, selection on the basis of the phenotype of these genotypes can be effective. Similar results were described earlier by [17] and Hossain [18]. However, moderate (10-20%) genotypic and phenotypic coefficients of variation were recorded in the present study for days to maturity, plant height, thousand grain weight, flag leaf length, and anther breadth. These results are similar with the findings of Bornane [19] for thousand grain weight. In contrast, low (<10%) estimates of genotypic and phenotypic coefficients of variation were observed in the present study for days to 50% flowering, number of grain per panicle, anther length, and pollen sterility indicating low variability for these characters in the present experimental material and therefore, there is little scope for improvement of these traits. Similar findings were reported earlier by Satish [20] for plant height.

### **3.4 Heritability (broad sense) and Genetic Advance as Percent Mean (GAM)**

Heritability estimates in broad sense ( $h^2_b$ ) were relatively higher (>60%) for almost all the traits studied. [5] stated a similar result in hybrid rice. High heritability evaluations have been found to be helpful in making a selection of superior genotypes on the basis of phenotypic performance.

High heritability (>60%) coupled with high genetic advance as percent mean (>20%) was recorded for days to maturity, number of tillers per hill, number of panicles per plant, thousand grain weight, filled grains/panicle, flag leaf length, flag leaf breadth, anther breadth, stigma length and breadth, filament length, and outcrossing rate indicating that the high heritability observed is due to additive gene effects and selection may be effective for these characters. Similar findings were observed by [22] for thousand grain weight, and [23] for outcrossing rate. On the contrary, high heritability coupled with moderate genetic advance in percent of mean (10-20%) was observed for days to 50% flowering, and plant height indicated the role of both additive and non-additive gene effects for controlling these characteristics. The results are similar with the reports of [24]. Further, information on genetic variation along with heritability and genetic advance assessments has been informed to give a better idea about the efficiency of selection.

In the present study, high GCV and PCV coupled with high heritability and high genetic advance in percent of mean were observed for number of panicles per plant, filled grain, flag leaf breadth, stigma length and breadth, filament length, and outcrossing rate indicating the preponderance of additive gene action and therefore, there is scope for improvement of the trait through selection. Similar results were given earlier by [25] and [26].

### **3.5 Mean Performance**

BRR1 1A recorded for maximum days to 50% flowering (70) and minimum in Ghan 46 A (63) ; Ghan 46A recorded for maximum days to maturity (90) followed by BRR1 1A (89); highest plant height observed in IR 58A (75.67 cm) and minimum in BRR1 1A (61.67 cm) ; panicle length found maximum in Ghan 46 A (24.30 cm) followed by IR 62A (23.21 cm); BRR1 1A revealed maximum number of tiller per hill (22.33) and panicle per plant (17.33) followed by Ghan 46 A (21.33) and (14.67) respectively; number of grain per panicle found highest in IR 58A (146.3) but number of filled grain observed highest in IR 62A (22.33); thousand grain weight found maximum in Ghan 46A (22.30gm) followed by IR 58 A (18.56 gm); highest flag leaf length found in IR 58A (32.52 cm) followed by Ghan 46A (24.10 cm) and

highest flag leaf width observed in Ghan 46A (1.533 cm) followed by IR 58A (1.467); Ghan 46A recorded for maximum anther length (444.0 um) followed by IR 58 A (407.3); maximum anther breadth recorded in IR 58A (93.17 um) followed by Ghan 46 A (92.50 um); BRR1 1A recorded for maximum stigma length (365.5um) and breadth (82.33um) followed by Ghan 46A (191.2 and 80.83) respectively; filament length observed highest in IR 62A (713.2 um) followed by Ghan 46 A (644.7 um); outcrossing rate found maximum in IR 58A (21.17) and minimum in BRR1 1A (17.96). Results revealed that out crossing rate varied significantly among the CMS lines (Table 3b) suggested outcrossing rate appeared as less flexible over the genetic background of the CMS lines. [27] reported a significant difference in outcrossing potential among the CMS lines.

UNDER PEER REVIEW

Table 3a: Mean performance of 5 CMS lines for 18 characters in rice (*Oryza sativa* L.)

CMS line	50% F	DMT	PH (cm)	PL (cm)	NT/H	NP/T	NG/P	TGW	Filled grain
Ghan46A	63 d	96 a	71.67 b	24.30 a	21.33 a	14.67 b	135.7 c	22.30 a	15.000 b
BRR1 1A	70 a	89 b	61.67 c	20.48 b	22.33 a	17.33 a	131.7 d	13.47 c	16.000 b
IR 68A	65 c	75 d	77.67 a	22.18 ab	18.67 b	9.000 c	131.7 d	17.00 b	14.000 b
IR 62A	64 d	73 e	64.33 c	23.21 a	18.67 b	9.000 c	141.0 b	18.55 b	22.33 a
IR 58A	69 b	79 c	75.67 a	22.29 ab	15.67 c	8.333 c	146.3 a	18.56 b	22.00 a
CV	.612	.3813	2.34	5.98	7.12	10.26	1.02	4.99	15.39
S.E	.235	.182	.95	.77	.79	.69	.81	.51	.69

50% F = Days to 50% flowering, DMT = Days to maturity, PH = Plant height, PL = Panicle length, NT/H = Number of tiller per hill NP/P = Number of panicle per tiller, NG/P = Number of grain per panicle, TW = Thousand grain weight

Table 3b: Mean performance of 5 CMS lines for 18 characters in rice (*Oryza sativa* L.)

CMS Line	FLL(cm)	FLW(cm)	AL (µm)	AB (µm)	SL (µm)	SB (µm)	FL (µm)	PS	OCR
Ghan46A	24.10 b	1.533 a	444.0 a	92.50 a	191.2 b	80.83 b	644.7 b	100.00 a	20.92 b
BRR1 1A	20.52 c	0.8333 b	405.3 c	66.00 c	365.5 a	82.33 a	443.3 c	97.00 b	21.27 a
IR 68A	23.27 b	1.433 a	418.0 b	91.83 b	197.2 c	62.50 d	354.7 d	95.00 c	18.77 c
IR 62A	23.32 b	0.9667 b	392.5 b	81.33 b	194.7 d	72.67 c	713.2 a	97.00 b	17.96 c
IR 58A	32.52 a	1.467 a	407.3 a	93.17 a	190.7 e	47.33 e	547.6 a	95.00 c	18.22 c
CV	2.24	13.18	1.2	2.32	2.25	3.5	.78	0.75	5.40
S.E	.32	.094	2.88	.99	2.25	1.43	2.35	1.34	0.60

FLL = Flag leaf length, FLW = flag leaf width, AL = Anther length, AB = Anther breadth, SL = Stigma length, SB= Stigma breadth, FL = Filament length, PF = Pollen sterility, OCR= Out crossing rate




Fig 1: Different floral traits of CMS line; a) White color anther with filament, b) Sterile pollen, c) Black color stigma, d) White color stigma

### Correlation and Coefficient

The phenotypic and genotypic correlation coefficients between outcrossing and other characters and **inter-relations** among them were estimated and presented in Table 4a, 4b. In

general, the results revealed that the genotypic correlations were higher than the phenotypic correlations for almost all the characters under study. The outcrossing ability of a CMS line is very important for seed set. It is influenced by several inter dependable traits. So, selection for outcrossing rate might not be effective unless influential traits would not be taken into consideration. Character association studies revealed a positive and significant association of outcrossing rate with number of grains per panicle, filled grain per panicle, stigma length and breadth both at genotypic and phenotypic levels, indicating that these traits were influenced by outcrossing rate. panicle length (PL) is an important yield-related trait as it strongly affects yield components which showed positive correlation with outcrossing rate, thousand grain weight. Similar results are reported [29]. While this trait showed a significant negative association anther length and anther breadth. Similar results are reported [30]. Plant height exposed positive association with panicle length, number of grain per panicle, thousand grain weight. This trait showed a negative association with number of tillers, number of panicles per plant and outcrossing rate at both genotypic and phenotypic levels respectively. Related findings were carried out by [31].

Table 4a : Phenotypic (above diagonal) and genotypic (below diagonal) correlations for floral and agronomical characters in CMS lines rice (*Oryza sativa* L.)

Character	Days to 50% flowering	Days to maturity	Plant height	Panicle length	Number of tiller /hill	Number of panicle /plant	Number of grain /panicle	thousa nd grain weight	Filled grain	Out crossin g rate
<b>Days to 50% flowering</b>	-	.0371	-.0933	-.6588*	-.0863	.2228	-.0376	-.7312**	.0658	-.0156
<b>Days to maturity</b>	.0399	-	-.2003	.0994	-.6272	-.8153*	.3091**	.2270	.4832*	.1230
<b>Plant height</b>	-.1127	-.2005	-	.2576	-.4999	-.5363*	.1504	.3984	-.1938	-.4505
<b>Panicle length</b>	-1.068*	.1622	.3605	-	-.0256	-.1834	.1834	.6848**	.0328	.1897
<b>Number of tillers/hills</b>	-.0945	-.7199*	-.6594	-.2280	-	.8747**	-.7453**	-.1514	-.645*	-.2723
<b>Number of panicles/tillers</b>	.2280	-.8541**	-.609*	-.3615	.9324**	-	-.5731*	-.2540	-.533*	-.1642
<b>Number of grain/panicle</b>	-.0311	.3158	.1657	.4106	-.7868**	-.5854*	-	.3559	.872**	.775**
<b>Thousand grain weight</b>	-.7415	.2314**	.4306	1.22**	-.2328	-.2606	.4006	-	.0745	.1805
<b>Filled grain</b>	.0693	.5059	-.1822	.1481	-.6449*	-.5302*	.9119**	.1041	-	.877**
<b>Outcrossing rate</b>	-.0164	.1235	-.4604	.2063	-.3131	-.1629	.8078**	.1957	.914**	-

**Table 4b- Phenotypic (above diagonal) and genotypic (below diagonal) correlations for floral and agronomical characters in CMS lines rice (*Oryza sativa* L.)**

Character	Flag leaf length	Flag leaf width	Anther length	Anther breadth	Stigma length	Stigma breadth	Filament length	Outcrossing rate
Flag leaf length	-	.5171	-.5427*	.7546**	.7442**	.7276**	.7021**	-.4878
Flag leaf width	.6162	-	.3300	.7109*	-.5248	-.5829*	.5654*	-.1041
Anther length	-.6554*	.3015	-	-.1080	.5553*	.8490**	.8491**	-.6112
Anther breadth	.7753**	.9009*	-.1103	-	.5362*	-.4139	-.3894	-.1087
Stigma length	.7486**	-.6163	.6340*	.5426*	-	-.851**	.5631*	.7637*
Stigma breadth	-.7459**	-.6413*	.9319**	-.4159	-.8615**	-	.8869**	.6905**
Filament length	.7086**	.6302*	.9572**	-.4051	.5659*	.8961**	-	-.4072
Outcrossing rate	-.5643	-.1894	-.7562	-.0692	.9453**	.8454**	-.5107	-

### 3.6 Path Coefficient Analysis

“As simple correlation does not provide the true contribution of the characters towards the yield, these genotypic correlations were partitioned into direct and indirect effects through path coefficient analysis” [32]. “Path coefficient analysis allows the separation of the direct effect and their indirect effects through other attributes by apportioning the correlations for a better interpretation of the cause-and-effect relationship. Path coefficient analysis was carried out using genotypic correlation, using outcrossing rate as a dependent variable. The results of path analysis revealed that panicle length, number of tillers per hill, number of grains per panicle, filled grain per panicle, thousand grain weight, flag leaf width, stigma breadth, and filament length exhibited positive direct effects with outcrossing rate” [32].

Critical analysis of results obtained from character association and path analysis indicated that number of grains per panicle, filled grain per panicle, stigma length, and breadth possessed both positive significant correlation and positive direct effects on the outcrossing rate. The results clearly indicated the true association of these traits with the outcrossing of these lines. Hence, that could be used in a selective breeding programs.

Table 5a: Direct and indirect effects (phenotypic) of agronomic and floral characters on yield in 5 CMS (A) lines of rice (*Oryza sativa*)

Character	Days to 50% flowering	Days to maturity	Plant height	Panicle length	Number of tiller /hill	Number of panicle /tiller	Number of grain /panicle	Thousand grain weight	Filled grain
Days to 50% flowering	<b>.2618</b>	-.0062	.0451	-.1396	-.0273	.0100	.0143	-.2082	.0337
Days to maturity	.0104	<b>-.1544</b>	.0803	.0212	.2083	.0373	-.1454	.0650	-.2462
Plant height	-.0295	.0310	<b>-.4004</b>	.0471	-.1908	-.0266	.0763	.1209	-.0887
Panicle length	-.2796	-.0251	-.1443	<b>.1308</b>	-.0660	-.0158	.1890	.3452	.0721
Number of tiller/hill	-.0247	-.1112	.2604	-.0298	<b>.2893</b>	.0407	-.3623	.0654	-.3139
Number of panicle/tiller	.0597	.1319	.2439	-.0473	.2697	<b>-.0437</b>	-.2695	-.0732	-.2581
Number of grain/panicle	.0081	.0488	-.0664	.0537	-.2276	.0256	<b>.4604</b>	.1125	.4439
Thousand grain weight	-.1941	-.0357	-.1724	.1607	-.0673	-.0114	.1845	<b>.2808</b>	.0507
Filled grain	.0181	.0781	.0729	.0194	-.1866	-.0232	.4199	.0292	<b>.4867</b>
Outcrossing rate	<b>-.0164</b>	<b>-.1235</b>	<b>-.4607</b>	<b>.2063</b>	<b>.3131</b>	<b>.1629</b>	<b>.8078</b>	<b>.1957</b>	<b>.9146</b>

RESIDUE= 0.0234

Table 5b: Direct and indirect effects (phenotypic) of floral and agronomical characters on yield in 5 CMS lines of rice

Character	Flag leaf length	Flag leaf width	Anther length	Anther breadth	Stigma length	Stigma breadth	Filament length
Flag leaf length	<b>-.1463</b>	.1541	-.2015	-.1733	.1054	-1.1093	.8067
Flag leaf width	-.0902	<b>.2501</b>	-.0927	-.1427	.1224	-.9538	.7174
Anther length	-.0959	.0754	<b>-.3074</b>	-.1468	.0150	-1.3861	1.0896
Anther breadth	-.1134	.2253	-.0339	<b>-.1256</b>	.1359	-.6186	.4612
Stigma length	.1095	-.1541	.1949	.2315	<b>-.0737</b>	1.2814	-.6442
Stigma breadth	.1091	-.1604	.2865	.1994	-.0565	<b>1.4873</b>	-1.0201
Filament length	-.1037	.1576	-.2943	-.1310	.0551	-1.3328	<b>1.1384</b>
Outcrossing rate	<b>-.5643</b>	<b>-.1894</b>	<b>-.7562</b>	<b>-.0691</b>	<b>.9452</b>	<b>.8454</b>	<b>-.5106</b>

RESIDUE=.3914

#### 4. Conclusion

The germplasm utilized in this study possessed useful variability and genetic parameter estimates for the studied traits that could be exploited for the rice genetic improvement program. The maximum outcrossing rate was found in IR 58A and the minimum in BRR1 1A. Correlation and path analysis revealed that number of grains per panicle, filled grain per panicle, stigma length, and breadth had a positive significant correlation along with the direct effect on outcrossing. Thus, it can be determined that selection for the aforementioned traits will undoubtedly increase the outcrossing rate of the lines, which in turn increases the amount of hybrid seed produced, making hybrid seed production more cost-effective and profitable for the seed industry and ultimately for the farming community.

**Conflict of interest:** The authors declare that they have no conflict of interest.

#### References

- [1] Hossain, M. S., Ivy, N. A., Raihan, M. S., Kayesh, E., & Maniruzzaman, S. (2020). Genetic Variability, Correlation and Path Analysis of Floral, Yield and its Component Traits of Maintainer Lines of Rice (*Oryza sativa* L.). *Bangladesh Rice Journal*, 24(1), 1-9.
- [2] Shozib, Habibul & Jahan, Saima & Sultan, Zakir & Alam, Samsul & Amin, Rifat & Hasan, Mahedi & Siddiquee, Muhammad. (2018). Nutritional Properties of Some BRR1 HYV Rice in Bangladesh. *Journal of Vitamins and Minerals*. 07. 10.4172/2376-1318.1000174.
- [3] Shivani, Dustakar & Jasmine, C & P., Senguttuvel & Naik, Saida. (2022). Genetic variability and association studies in maintainer and restorer lines of rice [*Oryza sativa* (L.)]. 11. 569-576.
- [4] Yearbook of Agricultural Statistics-2021 July 2020 Bangladesh Bureau of Statistics (BBS) Statistics and Informatics Division (SID) Ministry of Planning Government of the People's Republic of Bangladesh [www.bbs.gov.bd](http://www.bbs.gov.bd).
- [5] Faysal ASM, Ali L, Azam MG, Sarker U, Ercisli S, Golokhvast KS, Marc RA. Genetic Variability, Character Association, and Path Coefficient Analysis in Transplant Aman Rice Genotypes. *Plants* (Basel). 2022 Nov 2;11(21):2952. doi: 10.3390/plants11212952. PMID: 36365406; PMCID: PMC9655179.
- [6] Deepthi, K. P., Mohan, Y. C., Hemalatha, V., Yamini, K. N., & Singh, T. V. J. (2022). Genetic variability and character association studies for yield and yield related, floral and quality traits in maintainer lines of rice (*Oryza sativa* L.). *The Pharma Innovation Journal*, 11(2), 191-197.
- [7] FAOSTAT. 2017. World's crop production. Food and agricultural organization.
- [8] Fisher, R. A., & Yates, F. (1963). Statistical tables for biological, agricultural and medical research. *Statistical tables for biological, agricultural and medical research.*, (6th ed).
- [9] Burton, G.W. (1952, August). Qualitative inheritance in grasses. Vol. 1. In *Proceedings of the 6th*

*International Grassland Congress, Pennsylvania State College* (pp. 17-23).

- [10] Allard, R W. 1960. Principles of Plant Breeding. John Willy and Sons, Inc., New York
- [11] Johnson, H W, H F Robinson and R E Comstock. 1955. Estimation of genetic and environmental variability in soybean. *Agronomy Journal*, 47:314-318.
- [12] Miller, P.A., Williams, J. C., Robinson, H.F., & Comstock, R.E. (1958). Estimate of genotypic and environmental variances and co-variance in upland cotton and the implication selection. *Agronomy Journal*, 50, 126-131.
- [13] Wright, S. 1921. Correlation and causation. *Journal of Agricultural Research*, 20: 257-287.
- [14] Awad-Allah, M., Shafie, W. W., Alsubeie, M. S., Alatawi, A., Safhi, F. A., ALshamrani, S. M., ... & Masrahi, A. S. (2022). Utilization of Genetic Resources, Genetic Diversity and Genetic Variability for Selecting New Restorer Lines of Rice (*Oryza sativa* L.). *Genes*, 13(12), 2227.
- [15] S. Li, D. Yang, and Y. Zhu, Characterization and Use of Male Sterility in Hybrid Rice Breeding. *J. Integr. Plant Biol.*, 49, 791 (2007).
- [16] Niharika, T., Srikanth, T., Madhukar, P., & Yellagoud, T. (2022). Assessment of Genetic Variability and Correlation for Floral, Yield and Yield Contributing Traits in Rice (*Oryza sativa* L.). *International Journal of Environment and Climate Change*, 12(10), 322-331.
- [17] Singh, A. K., Singh, A. K., Tarkeshwar, A. G., & Maurya, S. P. (2022) Assessment of Genetic Variability and Heritability in CMS Lines and their F1's of Rice (*Oryza sativa* L.).
- [18] Hossain, Md & Maniruzzaman, Sheikh & Ivy, Nat & Sharif Raihan, Mohammad & Akter, Anowara & Hossain, Md. (2022). Assessment of Genetic Diversity, Correlation and Principal Component Analysis of Restorer Lines of Hybrid Rice. *International Journal Of Advance Research And Innovative Ideas In Education*. Volume 8. 1359-1374.
- [19] Majid, A., Waza, S. A., Shikari, A. B., Sofi, N. R., & Parray, G. A. (2022). Marker-based analysis of CMS lines for grain and kernel traits in rice hybrids. *Bangladesh Journal of Botany*, 51(2), 343-351.
- [20] Kulsum, U., Sarker, U., & Rasul, M. (2022). Genetic variability, heritability and interrelationship in salt-tolerant lines of T. Aman rice. *Genetika*, 54(2), 761-776.
- [21] Cochran, W. G. (1950). *Experimental design*. John Wiley & Sons. Developing Pusa 5A, a stable indica CMS line with high outcrossing potential.
- [22] Adilakshmi, D and M Girijarani. 2012. Variability, character association and path analysis in rice

- under submergence. *Crop Research*, 44(1 & 2): 146-151.
- [23] Madhavalatha, L, M ReddiSekhar, Y Suneetha and T Srinivas. 2005. Genetic variability, correlation and path analysis for yield and quality traits in rice (*Oryza sativa L.*). *Research on Crops*, 6 (3): 527-534.
- [24] Seyoum, M, S Alamerew, and K Bantte. 2012. Genetic variability, heritability, correlation coefficient and path analysis for yield and yield related traits in upland rice (*Oryza sativa L.*). *Journal of plant sciences*, 7(1):13-22.
- [25] Aswin, M., Umarani, R., Rajeswari, S., & Jeyaprakash, P. (2021). Genetic variability and association analysis in rice (*Oryza sativa L.*) germplasm based on biometrical and floral traits. *Electronic Journal of Plant Breeding*, 12(3), 912-917.
- [26] Hossain, M. S., Ali, M. A., Khan, M. A. I., & Maniruzzaman, S. (2020). Prevalence and Transmission of *Fusarium moniliforme*: A Seed Borne Pathogen of Rice. *Bangladesh Rice Journal*, 24(1), 11-19.
- [27] Hashim S, Ding P, Ismail MF, Ramli A. Floral traits and flowering behaviors of Malaysian rice cytoplasmic male sterile and maintainer lines and its relationship with out-crossing performance. *Aust. J. Crop Sci.* 2021;15:180–186
- [28] R. El-Namaky et al., *Putting plant genetic diversity and variability at work for breeding: Hybrid rice suitability in West Africa. Diversity*, 9, 27 (2017).
- [29] Anis, G. B. (2019). Research Article Assessment of Genetic Variability and Identification of Fertility Restoration Genes Rf3, Rf4 of WA-CMS in RILs Population of Rice.
- [30] Hasan, J. M., Kulsum, U. M., Majumder, R. R., & Sarker, U. (2020). Genotypic variability for grain quality attributes in restorer lines of hybrid rice. *Genetika*, 52(3), 973-989.
- [31] Gramaje, L. V., Caguiat, J. D., Enriquez, J. O. S., dela Cruz, Q. D., Millas, R. A., Carampatana, J. E., & Tabanao, D. A. A. (2020). Heterosis and combining ability analysis in CMS hybrid rice. *Euphytica*, 216, 1-22.
- [32] Maniruzzaman, S., Rahman, M. A., Hasan, M., Rasul, M. G., Molla, A. H., Khatun, H., & Akter, S. (2022). Genetic Mapping to Detect Stringent QTLs Using 1k-RiCASNP Genotyping Platform from the New Landrace Associated with Salt Tolerance at the Seedling Stage in Rice. *Plants*, 11(11), 1409.