

Original Research Article

Remote sensing indexes assessment for Drought monitoring using sentinel satellite imagery: case study from Natanz County, Iran

Abstract

Water demand is increasing worldwide, mainly owing to the growing population and industrial development [1]. Excessive population growth [2], accompanied by rapid industrialization and urbanization [3] In recent years, the problem of climate change and decrease in rainfall has become a big challenge in Iran, so it's monitoring and control is very vital and necessary. In addition to the classical methods, nowadays with the progress of the space industry in the world, the use of remote sensing techniques has greatly helped to detect and monitor drought. The science of remote sensing and the use of satellite images is considered a useful and optimal tool for drought monitoring. In this project, we investigated the occurrence of drought using satellite images and remote sensing techniques. In this way, we have calculated the drought-related remote sensing indicators, including NDVI, VCI, TCI, and VHI, in the six-year period between 2015 and 2021 for Natanz city located in Isfahan province. In order to calculate the indicators, Sentinel satellite data was used, and to calculate the Standard Rainfall Index (SPI), data from the synoptic station of Natanz city was used. According to the general results obtained from the indicators, currently the drought situation in the southwestern region of this city has been increasing compared to the last five years, and in other regions we are witnessing the growth and occurrence of drought sporadically. The results of this research show that the best index for calculating and monitoring drought is the VCI index because it has a higher correlation with meteorological data and the SPI index.

Key words: drought index, satellite imagery, remote sensing, sentinel satellite;

Comment [Ma1]: Abstract should tell us about the aim of the study Statement of the Problem, Methodology employed, Results, Conclusion and Contribution to knowledge. Author should try and re-write the abstract to clearly reflect all the characteristics as listed.

Comment [Ma2]: Reference not required at this stage in Abstract.

Comment [Ma3]: Arrange in alphabetical order

Introduction

Drought is a complex, globally occurring phenomenon that affects humans and nature alike [1]. Due to climate change and decrease in precipitation, drought has become a big challenge in the world, especially in dry and semi-dry areas like Iran; Thus, monitoring and management of this is an important issue. In front of traditional methods that depend on meteorological station observations and focused on more investigation of the meteorological drought, by using remote sensing techniques and satellite imagery as a useful method for temporal and spatial agricultural monitoring are of interest for researchers. Subsurface water storage situation in Iran is not affirmative. In recent decades, the value of subsurface water storage in Iran equal to 131 billion m^3 according to statistical tests reports, but in recent years it decreased to 111 billion m^3 , also global warming and defect of precipitation caused this disaster. According to some researches, the average of precipitation was 275 mm, have mentioned that represent the most part of Iran is dry and semi-dry region [2]. So because of these problems and climate change of Iran, drought is an inevitable issue in this country, therefore monitoring and investigation of this phenomenon is necessary. In this paper we evaluated remote sensing drought indicators by using satellite imagery. Several studies had been done in this field; Sepulcre-Canto et al [3], Applied a combination of SPI, soil moisture anomaly, and the fraction of absorbed photosynthetically active radiation

, While Trnka et al [4] used a combination of soil moisture data and EVI in the Czech Republic and Slovakia. Both approaches were shown to be promising. Finally, approaches also exist within this topic to determine areas and time periods where water is the primary limiting factor for plant growth. In

Europe, a correlation analysis between LST and NDVI has revealed that energy was a limiting factor mainly in northern Europe at high altitude, and in spring while water limited plant growth mainly southern Europe in summer. Despite the large numbers of existing studies on agricultural and vegetation-based drought in central Europe using remote sensing, uncertainties and larger knowledge gaps still exist [5]. Rezaei et al evaluated performance of temperature conditional index (TCI) and vegetation conditional index (VCI) using MODIS satellite images for estimating drought in Urmia lake; they concluded that MODIS satellite images have fine accuracy for estimating drought. They recommended remote sensing techniques except of meteorological methods due to better temporal resolution, less expenditure, more surface coverage and more resampling points [6]. Rezaei-Banafshe et al for drought analysis at east Azarbaijan province by using MODIS satellite images from 2011 to 2017 computed the VCI (vegetation conditional index) they understood maximum drought depended on 2011, 2012, consequently, and minimum values depend on 2013. For validation VCI results in 2 synoptic station data used and compare with SPI values for three months in 2011, 2012, 2013. This results represented according to SPI values confirm VCI values [7]. In 2006 Kogan et al evaluated Mongolia's drought by

Comment [Ma4]: In recent decades, and not in recent decade value.....

Comment [Ma5]: Check if indicated or indicates

Comment [Ma6]: Check if the two words are together or not e.g. photosynthetically or photo synthetically

using AVHRR-NOAA satellite data; They found out that VHI had the best results on the other hand in Afghanistan, Pakistan and two country in west side of India Then kaba and his coworkers estimated VHI, VCI, TCI and NDVI for drought monitoring. Their results represented that due to lack of ground observations, remote sensing have a very important role, also they recommended merging AVHRR, MODIS daily data method in 2006 [8]. Dilayda Soyulu Pekpostalci et, al demonstrated in their research that the emerging RS technology and satellite-driven indicators were rarely used in the country. The review showed that there is room for more research on agricultural and hydrological drought monitoring, forecasting, and pattern detection in Turkey [9]. Ziyu Wanget, al. used soil moisture was added to calculate a new drought index based on TVPDI in southeastern Tibet, named the Temperature–Vegetation–Soil–Moisture–Precipitation–Drought Index (TVMPDI). Then, the TVMPDI was validated by using the Standardized Precipitation Evapotranspiration Index (SPEI) and other remote sensing drought indices, including the Vegetation Health Index (VHI) and Scale Drought Conditions Index (SDCI), during the growing seasons of 2003–2018. [10] despite of the important role of Natanz county there isn't any new study about drought monitoring by using remote sensing technique, in this study investigation of drought monitoring have been done by using several indexes to choose the best one for future studies by considering meteorological data of case study.

2. Materials and methods

In this study main flow-chart is shown in figure 1, including of collecting precipitation stations data, validation and SPI computation; on the other side satellite data acquisition, LST and NDVI computation, coding and VCI, TCI and VHI estimation. The final part of this flow-chart is computation of correlation factor between satellite-based indicates and ground-based indicates, best indicate selection and the last step is drought monitoring.

Comment [Ma7]: Methodology not clearly explain enough. Mythology should state clearly the research design, sources of data: either primary or secondary methods of data sources. It should also include research population, sample size and frame, sample techniques and process of data collection.




Figure 1: Work flowchart

Comment [Ma8]: Flow chart of what? Figure reference must be properly done and should be properly refer to within the text contexts

2.1 study area

The case-study of this research is central part of Natanz county between 33° 25' and 33° 40' N and 51° 25' and 52° E. This region have mountains and deserts that a mountainous area with a maximum height of 2242 meters above sea level and a desert area with a height of 272 meters above sea level, it has 2 types of climate: cold, desert and temperate. From the north to Aran and

Bidgol deserts, from the south to Mimeh is limited to Ardestan from the east and Kashan from the west. This county is located 752 km north of Isfahan, 72 km southeast of Kashan and 254 km from Tehran. Water of this area is provided from seasonal rivers and mostly from springs and aqueducts .average rainfall is 772 mm annually and there is no rainfall in the months of July, August and September. This issue indicates that the growing season of the plant without rain and the water from the winter rain is waste water, unavailable and unusable. Maximum degree of the absolute temperature is 4.22 degrees Celsius and the minimum absolute temperature is -4.72. Maximum

humidity is 60% and minimum Average humidity is 34%. The dry season in this region is relatively long and starts from May to the end of October. February is the coldest month and the driest month July is the hottest month of the year. The rainiest month of the year is March, which is also the wettest month of the year [9]




Figure 2. case study location [19]

Comment [Ma9]: Refer to Figure 2 by explanation in the test above (Study Area)

2.2 Data

Sentinel data used for this research and compute NDVI and LST raster data then calculate drought indicates. These data was downloaded from USGS data center. Chosen period for all sentinel satellite product was from 2015 to 2021. The spatial resolution of sentinel 3 satellite images is 300 m and temporal resolution is 10 days [10]. On the other hand collecting meteorological data from ground synoptic station including pressure, temperature, humidity and rainfall that archived to the governmental organization which provided valid statistical parameters using high precision instruments [11]. by using previous parameters, Standard Precipitation Index (SPI) was calculated into monthly average. All of data in this research processed by ENVI 5.6 (The Environment for Visualizing Images) for computing VCI, TCI and VHI Indexes. In this research classification upon to maximum likelihood algorithm is used.

2.3 calculation and assessment of drought indices

VCI, TCI and VHI the SPI is the geo-referenced indicate which is computed upon from ground station data.

SPI is one of the basic indices, for its calculation, it is necessary to have the long-term average and standard deviation of rainfall values for the studied periods. Lack of precipitation in short-term time scales mainly affects soil moisture and long-term often affects underground water, river flow and water resource reserves [12]. This index can be calculated through equation 1:

$$SPI = \frac{P_i - \bar{P}}{SD} (1)$$

In this regard; SPI is the standard index of precipitation, P_i is the precipitation of the desired year in millimeters, \bar{p} is the long-term average precipitation of the station, and SD is the standard deviation of precipitation. Figure (3) shows the classification of the drought situation based on the SPI index. [13] The LST is drive from below formula:

$$LST = \frac{T_b}{1 + (L\lambda * T_b + P) * Ln\epsilon} \quad (2)$$

In this formula $L\lambda$ scattered wave length radiance, T_b brightness temperature radiation power and P value is calculated from this formula:

$$P = h * c * (1.438 * 10^2 m.) \quad (3)$$

That h is Planc factor ($6.626 * 10^{-34}$ J.S), σ EstephanBoltezman factor ($1.38 * 10^{-23}$ /K), c is speed of light ($2.998 * 10^8$ m/s).

NDVI that is Normalized Difference Vegetation Index is known as one of the most famous, simplest and most practical plant indices. The normalized difference vegetation index is expressed as follows:

$$NDVI = \frac{NIR + RED}{NIR - RED} \quad (4)$$

That NIR is near infrared band and RED is the red band values. This index has values between +1 and -1. The usual range of changes in green vegetation is equal to 0.2 to 0.8 (Tucker, 1997). The value of NDVI index is usually between 0.05 and 0.1 for sparse vegetation areas, and between 0.1 and 0.6 for normal and semi-dense vegetation areas. And for very dense plant areas, it is between 0.6 and 0.7. [14]

VCI is the Vegetation Condition Index, The amount of vegetation on the surface of the earth is basically determined by the main of the environmental Physical components such as climate and soil are controlled [15]. Purpose of VCI is identifying the effects of ecosystem change on vegetation in the region.

$$VCI = \frac{NDVI - NDVI_{MIN}}{NDVI_{MAX} - NDVI_{MIN}} \quad [16] \quad (5)$$

Table (1) VCI value range

Class	Very High	High	Almost High	Mean	Normal
VCI	<10	<20	<30	<40	40>

TCI is Temperature Condition Index, that is assumed the phenomenon of drought causes a decrease in soil moisture and creates the heat stress will be on the earth's surface. The range of changes of this index is between 0 1. [17]

$$TCI = \frac{LST_{MAX} - LST}{LST_{MAX} - LST_{MIN}} \quad (6)$$

That LST_{min} and LST_{max} are the minimum and maximum temperature values in the time period. [18]

VHI index is a combined from VCI and TCI indices, the purpose of this index is, employing vegetation conditions and ground surface temperature in total index. [19]

$$VHI = (0.5 * VCI) + (0.5 * TCI) / 7$$

3. Result analysis

In this paper our purpose is assessment the drought indicates using by remote sensing satellite imagery during 2015 to 2021 for central part of Natanzcountyand recognize the most appropriate indicate.

3.1 Processing results




Figure3.Monthly average of synoptic station data during 2015 – 2021 (mm)

Comment [Ma10]: Refer to Figure 3 by explanation in the test above (Study Area)

As mentioned in the definition of the SPI index, the input values of this index are the only factor of rainfall, and the purpose of creating the SPI index is to digitalize the analysis of the factor of rainfall, increasing positive values show a wet year indicate and decreasing negative values show more drought .As can be seen in the diagram, from 2015 to the end of 2019, despite the decrease, the values are positive, but their decrease in the period of 2019 to the end of the research period has even approached the number -2, which indicates a very severe drought situation.

Comment [Ma11]: Can you put this above figure 3 and not below figure 3




Figure.4 the normalized difference vegetation index (NDVI) during 2015 – 2021

According to the above figure, the extent of areas with weak vegetation is increasing in the northwestern areas of the city, also in the central areas of the city and in the southwestern part, the growth of areas with weak vegetation can be seen, and also in the southwestern region in 2015, 2016 and 2017, the vegetation pattern is same and generally no vegetation was seen, but in 2018, 2019 and 2020, the vegetation slightly improved and grew towards the areas with weak coverage and higher NDVI values.

But in 2021, it can be seen that the loss of weak vegetation and NDVI is less than the previous years.

Comment [Ma12]: Mention the figure in full e.g figure 4




Figure .5 Temperature Condition Index during (TCI) 2015 – 2021

According to Figure 5, for study area using the TCI index, toward 2015 in 2016 and 2017, an increase in severe drought areas is shown, especially in the southwestern, but in 2018, in the central and western, the drought situation improved and even some part of the region appeared without drought, while in 2019 there was a sharp decrease in drought-free areas and an increase in mild drought areas, in 2020 and 2021 toward 2019, by checking the index in the central areas according to the classification, the situation can be seen to improve.




Figure 6.vegetation health index (VHI) during 2015-2021

According to the survey based on the VHI index, in 2016 and 2017 compared to 2015, an increase in the extent of areas with very severe drought can be seen in the southwest region, and in 2019, compared to 2018, there has been an improvement in the areas without drought and severe drought, but in the southwest region, severe drought has increased, a strong trend and by 2020 is shown, the drought situation in this area is getting worse. It can be seen in 2021 increase and decrease of areas with severe drought in the south and southeast regions, therefore, same as the previous indicators, the conclusion needs to match the maps produced with the numerical values obtained from the formulas and the standard deviation.




Figure 7.vegetation condition index (VCI) during 2015-2021

In 2016, compared to 2015, can be seen an improvement in drought situation, in 2017 southwest of this area drought is increased toward 2016.although In 2018, 2019, 2020 same pattern is shown but in 2021 drought situation has become much worse. According to SPI graph, the decreasing trend can be seen Better in the VCI index obtained maps.

3.2 Numerical review tables of the above indicators

Table 2: TCI values

	2015	2016	2017	2018	2019	2020	2021
Mean	0.95	0.903	0.39	0.877	0.73	1.05	0.8
Standard deviation	0.60	0.59	0.62	0.39	0.691	0.84	0.695

Table 3: VCI index

	2015	2016	2017	2018	2019	2020	2021
Mean	39.178	31.52	48.48	29.98	33.102	28.55	21.3
Standard deviation	0.137	0.2575	0.388	0.28	0.2234	0.24	0.212

Table 4: VHI index

	2015	2016	2017	2018	2019	2020	2021
Mean	20.05	16.25	20.44	14.43	17.08	14.82	11.08
Standard deviation	0.33	0.32	0.34	0.32	0.33	0.44	0.37

According to the correlation between the standard deviation values of all three indicators, the correctness of the results is validated by acceptable values that is depended to the accuracy of the Sentinel satellite images.

4 .conclusion

In this paper assessment of drought indicates using sentinel satellite high resolution images has been done during 2015 to 2021 at Natanzcountyby remote sensing methods that is appropriate technique for drought monitoring in new studies. At first step the NDVI has been estimated and vegetation maps is extracted the results show that the studied area have poor land vegetation. According to purpose of this research the land surface temperature (LST) has been calculated and three indicates VCI, TCI, VHI which are related to both of these arrival data (LST and NDVI) has choose. All results demonstrated increase of drought .According to this indicates drought classified to very high, high, middle, light and without drought classes.The Vegetation Conditional Index (VCI) has the highest correlation with the results of the ground depended data methods that is, obtained by using of meteorological parameters and the Standard Precipitation Index (SPI). The annual rainfall in

2021 is very low. On the other hand, the obtained drought situation from this index in 2021 has also been associated with an increase in the amount of drought in the southwest and northwest regions.Also, in the examination of the numerical values according to the definition of the drought index, when the number obtained from the formula is between 10 and 40, the drought conditions are increasing with the increasing trend of this value, which is exactly the same problem that can be seen in the linear representation of the index, so in this research, the best index for The assessment of the degree of drought is VCI. Among the calculated indices, VHI has the lowest correlation with the

Comment [Ma13]: Remove

Comment [Ma14]: Change to Has

Comment [Ma15]: Demonstrated an increase

Comment [Ma16]: According to the purpose

Comment [Ma17]: Ground-depended

Comment [Ma18]: Remove that is

Comment [Ma19]: Remove of

Comment [Ma20]: drought situation obtained

SPI index and, naturally, it has provided the weakest result. It should be noted that the NDVI and TCI indices show a relatively similar performance both in the generated maps and in the numerical values of the table and have a relatively good correlation with the SPI index. But the range of their changes does not show the changes made in the region as well as VCI. By use of this indicates, gives an ability to drought management, also use of this indicates due to high spatial and temporal resolution can assist to making decision in several agricultural purpose.

Comment [Ma21]: decisions

Comment [Ma22]: purposes

References

- [1] Sterling, S.M.; Ducharme, A.; Polcher, J. The impact of global land-cover change on the terrestrial water cycle. *Nat. Clim. Chang.* 2013, 3, 385–390. [CrossRef]
- [2] Rockström, J.; Lannerstad, M.; Falkenmark, M. Assessing the water challenge of a new green revolution in developing countries. *Proc. Natl. Acad. Sci. USA* 2007, 104, 6253–6260. [CrossRef] [PubMed]
- [3] Henderson. Urbanization in developing countries. *World Bank Res. Obs.* 2002, 17, 89–112. [CrossRef]
- [4] Heim, R.R. A Review of Twentieth-Century Drought Indices Used in the United States. *Bull. Am. Meteor. Soc.* 2002, 83, 1149–1166
- [5] EngdaworkAsfaw, K., Suryabagavan, V. & Argaw, M. (2018). Soil salinity modeling and mapping using remote sensing and GIS: The case of Wonji sugar cane irrigation farm, Ethiopia. *Journal of the Saudi Society of Agricultural Sciences*, 17 (3), 250-258.
- [6] Sepulcre-Canto, G.; Horion, S.; Singleton, A.; Carrao, H.; Vogt, J. Development of a Combined Drought Indicator to detect agricultural drought in Europe. *Nat. Hazards Earth Syst. Sci.* 2012, 12, 3519–3531.
- [7] Trnka, M.; Hlavinka, P.; Možný, M.; Semerádová, D.; Štěpánek, P.; Balek, J.; Bartošová, L.; Zahradníček, P.; Bláhová, M.; Skalák, P.; et al. Czech Drought Monitor System for monitoring and forecasting agricultural drought and drought impacts. *Int. J. Climatol.* 2020, 40, 5941–5958
- [8] Karnieli, A.; Ohana-Levi, N.; Silver, M.; Paz-Kagan, T.; Panov, N.; Varghese, D.; Chrysoulakis, N.; Provenzale, A. Spatial and Seasonal Patterns in Vegetation Growth Limiting Factors over Europe. *Remote Sens.* 2019, 11, 2406

[9] rezaeimoghadam,m ; valizade.k ;Modis data performance analysis in drought estimation ,journal of geography and environmental stability 1991 ,4,97-54, 5.

[10]Ziyu Wang 1, Zegen Wang 2,*, JunnanXiong 1,3,WenHe 4, Zhiwei Yong 2 and Xin Wang 1,Responses of the Remote Sensing Drought Index with Soil Information to Meteorological and Agricultural Droughts in Southeastern Tibet Article in Remote Sensing · December 2022 DOI: 10.3390/rs14236125

[11] Rezaie,b ; rezaie.m ; freydonpoor.a agricultural drought analysis in eastern azarbayjan province with remote sensing approach and vegetation condition index ,1991 water and soil journal,52 ,1,119-149 ,1.

[12] Mckee, T.B.; Doesken, N.J.; Kleist, J. The Relationship of Drought Frequency and Duration to Time Scales. In Proceedings of the 8th Conference on Applied Climatology, Anaheim, CA, USA, 17–22 January 1993; Volume 17, pp. 179–183

[13] Bhuiyan, C., Singh, R.P., Kogan, F.N., 2006. Monitoring drought dynamics in the Aravalli region (India) using different indices based on ground and remote sensing data. International Journal of Applied Earth Observation and Geoinformation 289-302

[14] DilaydaSoyluPekpostalci 1 , Rifat Tur 2, Ali DanandehMehr 3,4,* , Mohammad Amin Vazifekhah Ghaffari5,Dominika D abrowska 6 and VahidNourani 7,8,Drought Monitoring and Forecasting acrossTurkey: A Contemporary Review,Article in Sustainability · March 2023, DOI:10.3390/su15076080

[15] <http://zistnatanz.blogfa.com>

[16] Sentinel User Handbook, ESA Standard Document Date 24/07/2015 Issue 1 Rev 2

[17]<https://www.irimo.ir>

Comment [Ma23]: This is not a good way of writing references

[18] Afshar, M.H.; Al-Yaari, A.; Yilmaz, M.T. Comparative Evaluation of Microwave LBand VOD and Optical NDVI for Agriculture Drought Detection over Central Europe. *Remote Sens.* 2021, *13*, 1251.

[19] Bachmair, S.; Tanguy, M.; Hannaford, J.; Stahl, K. How well do meteorological indicators represent agricultural and forest drought across Europe? Environ. Res. Lett. 2018, 13, 34042.

[20]www.wikipedia.org

Comment [Ma24]: Please use APA Format of Presenting references

UNDER PEER REVIEW