

Original Research Article

Frequency of *H. pylori* infection among Patients with Gastrointestinal Symptoms Attending Somali Sudanese Specialized Hospital (SSSH), Mogadishu, Somalia

ABSTRACT

Background: *Helicobacter pylori* is a human-specific pathogen, which leads to gastric pathologies including gastric cancer. Fifty percent of the world's population from 62 countries is infected with *H. pylori* infection, making it the most widespread infection in humans. The aim of the study was to determine the frequency of *H. pylori* infection among patients with gastrointestinal symptoms.

Materials and methods: This study was descriptive cross-sectional Hospital-based study conducted at Somali Sudanese Specialized Hospital (SSSH), Mogadishu, Somalia during the period of December 2022 to March 2023. A total of 1009 subjects were included in the study. Fresh stool samples were collected for antigen detection. The test was done in the ICT of *H. pylori* stool antigen detection (Intec, China; catalog number ITP10003-DS50). The data was gathered using pre-designed structural questionnaire and the SPSS 26.0 statistical software (SPSS Inc., USA) was used for statistical analysis.

Results: The age of the study population ranges from 5 months to 62 years with a mean age of 34 ± 27.5 years. 57.4% of them were female patients, where 42.6% were male patients. The study showed no significant correlation between gender and frequency of *H. pylori* infection with P-value of 0.137. According to the frequency of *H. pylori* infection among the patients with gastrointestinal symptoms; 44.8% of them had *H. pylori* infection. Regarding to frequency of *H. pylori* infection in the recent months, the most frequent period of prevalence was in February (56.5%), followed by January (51.1%), followed by December (43.8%) where the least frequent was in March (26.9%) which showed significant correlation between prevalence period and *H. pylori* infection frequency with P-value of 0.000.

Conclusion: This study concluded a high frequency of *H. pylori* infection among patients with gastrointestinal symptoms.

Keywords: *H. pylori* infection, ICT, Stool antigen test, gastrointestinal system, Somalia

INTRODUCTION

“*Helicobacter pylori* is a spiral-shaped, flagellated, Gram-negative pathogen that only affects humans. It colonises the human stomach and causes gastric issues. It is a bacterium that is extremely rare and is thought to be carcinogenic. 90% of cases of gastric cancer are still linked to *H. pylori*, which continues to be a serious health concern for people. *H. pylori* was identified in 1982 by Barry Marshall and Robin Warren as the primary cause of chronic gastritis and peptic ulcer disease” [1,2].

“It is estimated that 4.4 billion cases of *H. pylori* infection occurred worldwide in 2015, with significant regional and national differences in the disease's prevalence. *H. pylori* infection affects 50% of the world's population, spread across 62 different countries. In Africa and developing countries, it is estimated that the highest prevalence of *H.pylori* infection to be around 80%” [3,4]

A study done by Alebie [5] in Jigjiga city, Ethiopia Jigjiga, Somali Regional State of Ethiopia in 2016 about the prevalence of *H.pylori* infection in Jigjiga University reported high prevalence of *H.pylori* infection among gastritis students.

There is lack of data about the prevalence of *H.pylori* infection in Somalia and getting basic data about this disease is important for Health institutions to reduce the burden of the disease. Regarding to this, the present study was designed to determine the frequency of *H.pylori* infection among patients complaining from gastrointestinal symptoms in Somalia.

MATERIALS AND METHODS

This study was descriptive cross-sectional Hospital-based study conducted at Somali Sudanese Specialized Hospital (SSSH), Mogadishu, Somalia during the period of December 2022 to March 2023. a total of 1009 subjects were included in this study, Patients with Gastrointestinal Symptoms Attending Somali Sudanese Specialized hospital (SSSH) during the aforementioned period were included in the study, while patients who had no symptoms of *H. pylori* infection and any patient who refused to give consent were excluded. Fresh stool samples were collected into spoon-cover and outer-labeled stool container for antigen detection. Using a wood stick, a small portion of the stool sample was transferred into buffer, incubated for 2 minutes and then

two to three drops of the mixture were poured in the hole of the ICT of *H. pylori* stool antigen detection (Intec, China; catalog number ITP10003-DS50). The color migrated from the well containing the tested sample in the ICT device. The presence of two bands (test band (T) and control band (C)) within the result window, no matter which band appeared first, indicated a positive result. The data was gathered using per-designed structural questionnaire and the SPSS 26.0 statistical software (SPSS Inc., USA) was used for statistical analysis. Finally, the study was licensed by the ethical committee of Somali Sudanese Specialized Hospital (SSSH).

RESULTS

A Total of 1009 samples were collected from the patients with gastrointestinal symptoms, Their age ranged from 5 months up to 62 years with a mean age of 34 ± 27.5 years. 57.4% of the study population were female patients, where 42.6% were male patients. The study showed no significant correlation between demographic data (Age and Gender) and frequency of *H.pylori* infection with P-value of 0.50 and 0.137 respectively. Regarding to the frequency of *H.pylori* infection among these patients with gastrointestinal symptoms attending Somali Sudanese Specialized Hospital; 44.8% of them had *H.pylori* infection. In the last three months, the most frequent period of *H.pylori* infection was in February (56.5%), followed by January (51.1%), followed by December (43.8%) where the least frequent was in March (26.9%) which showed significant correlation between prevalence period and *H.pylori* infection frequency with P-value of 0.000. The results were shown in tables (1,2,3,4,5,6).

Table 1: Distribution of gender in the study population

| Gender | Frequency | Percent | Valid Percent | Cumulative Percent |
|--------|-----------|---------|---------------|--------------------|
| Female | 579 | 57.4 | 57.4 | 57.4 |
| Male | 430 | 42.6 | 42.6 | 100.0 |
| Total | 1009 | 100.0 | 100.0 | |

Table 2: Distribution of *H.pylori* infection in the study population

| H.pylori infection | Frequency | Percent |
|---------------------------|------------------|----------------|
| Positive | 452 | 44.8 |
| Negative | 557 | 55.2 |
| Total | 1009 | 100.0 |

Table 3: Frequency of H.pylori infection in the recent months

| Duration period | Frequency | Percent |
|------------------------|------------------|----------------|
| March 2023 | 261 | 25.9 |
| February 2023 | 232 | 23.0 |
| January 2023 | 235 | 23.3 |
| December 2022 | 281 | 27.8 |
| Total | 1009 | 100.0 |

Table 4: Correlation between age and frequency of H.pylori infection

| H.pylori infection | N | Mean Age \pm Std. Deviation | P-value |
|---------------------------|----------|---|----------------|
| Positive | 452 | 31.49 \pm 17.13 | 0.50 |
| Negative | 557 | 33.73 \pm 19.26 | |

Table 5: Correlation between gender and frequency of H.pylori infection

| | | | H.pylori infection | | Total | P=value |
|--------|--------|-----------------|--------------------|----------|--------|---------|
| | | | Positive | Negative | | |
| Gender | Female | Count | 271 | 308 | 579 | 0.137 |
| | | % within Gender | 46.8% | 53.2% | 100.0% | |
| | Male | Count | 181 | 249 | 430 | |
| | | % within Gender | 42.1% | 57.9% | 100.0% | |
| Total | | Count | 452 | 557 | 1009 | |
| | | % within Gender | 44.8% | 55.2% | 100.0% | |

Table 6: Correlation between Duration period and frequency of H.pylori infection

| | | | H.pylori infection | | Total | P-value |
|-----------------|---------------|--------------------------|--------------------|----------|--------|---------|
| | | | Positive | Negative | | |
| Duration period | March 2023 | Count | 78 | 183 | 261 | 0.000 |
| | | % within Duration period | 29.9% | 70.1% | 100.0% | |
| | February 2023 | Count | 131 | 101 | 232 | |
| | | % within Duration period | 56.5% | 43.5% | 100.0% | |
| | January 2023 | Count | 120 | 115 | 235 | |
| | | % within Duration period | 51.1% | 48.9% | 100.0% | |
| | December 2022 | Count | 123 | 158 | 281 | |
| | | % within Duration period | 43.8% | 56.2% | 100.0% | |
| Total | | Count | 452 | 557 | 1009 | |
| | | % within Duration period | 44.8% | 55.2% | 100.0% | |

DISCUSSION

A human-specific pathogen called *Helicobacter pylori* causes diseases of the stomach, including gastric cancer. 62 countries and 50% of the global population are home to *H. pylori* infections, making it the human infection with the greatest spread. The present study was descriptive cross-sectional hospital-based study conducted at Somali Sudanese specialized hospital, Mogadishu, Somalia, for determination of *H. pylori* infection frequency among patients with gastrointestinal symptoms.

The results of this study revealed that the mean age of the study population was 34 ± 27.5 years (ranged from 5 months up to 62 years). 57.4% of the study population were female patients, where 42.6% were male patients. There was no statistically significant correlation between demographic data (age and gender) and frequency of *H. pylori* infection (P-value of 0.5 and 0.137 respectively). these findings disagree with a study done by W. Hong¹, et al (2019) who reported that *H. pylori* infection frequency is related to age, gender.⁽⁶⁾

Regarding to the frequency of *H. pylori* infection among the patients with gastrointestinal symptoms attending Somali Sudanese Specialized Hospital; 44.8% of them had *H. pylori* infection, The prevalence of *H. pylori* infection in this study was lower when compared with results reported by Maria P. C., et al (2019), and Bakka et al (2002) which show high positive percentage of *H. pylori* infection in Latin America and the Caribbean populations (adults 69.26%) and Libya (94%), that might be the reason of demographic differentiation as well as characteristics of the populations.⁽⁷⁻⁸⁾ But the finding of this study was similar with a study done by Mohammed A.I.A., et al (2020) in Western Sudan which found that *H. pylori* positive was 45.8%,⁽⁹⁾ also the frequency of *H. pylori* infection in this study was similar with results reported by Shuai Ren in mainland china (2022) which found that 44.2% (95% CI: 43.0–45.5%) of the population had *H. pylori* infection.⁽¹⁰⁾

CONCLUSION

This study concluded high frequency of *H.pylori* infection among patients with gastrointestinal symptoms attending Somali Sudanese Specialized Hospital, Mogadishu, Somalia.

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

Consent

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

REFERENCES

- 1-Meliş, L.E.; Mărginean, C.O.; Mărginean, C.D.; Mărginean, M.O., “The Relationship between Toll-like Receptors and *Helicobacter Pylori*-Related Gastropathies: Still a Controversial Topic”, *J. Immunol. Res*, 2019; 8197048.
- 2-Li RJ, Dai YY, Qin C, Huang GR, Qin YC, Huang YY, Huang ZS, Luo XK, Huang YQ, “Application of traditional Chinese medicine in treatment of *Helicobacter pylori* infection”, *World J Clin Cases*, 2021; 16;9(35):10781-10791.
- 3-Boustany A, Onwuzo S, Almomani A, Asaad I, “Epidemiology and risk of colorectal cancer in patients with a history of *Helicobacter pylori* infection: a population-based study”, *Ann Gastroenterol*, 2023; 36(2):203-207. doi: 10.20524/aog.2023.0783. Epub 2023, Feb 3.
- 4-Hooi JKY, Lai WY, Ng WK, Suen MMY, Underwood FE, Tanyingoh D, Malfertheimer P, Graham DY, Wong VWS, Wu JCY, Chan FKL, Sung JJY, Kaplan GG, Ng SC, “Global prevalence of *Helicobacter pylori* infection: systematic review and meta-analysis. *Gastroenterology*”, 2017; 153(2):420–429. <https://doi.org/10.1053/j.gastro.2017.04.022>.
- 5-Alebie G, Kaba D., “Prevalence of *helicobacter pylori* infection and associated factors among gastritis students in Jigjiga University, jigjiga, somali regional state of Ethiopia”, *J Bacteriol Mycol Open Access*. 2016; 3(3):234-239. DOI: 10.15406/jbmoa.2016.03.00060.

6-W. Hong, H.-L. Tang, X.-L. Dong, S.-K. Hu, Y. Yan, "Prevalence of *Helicobacter pylori* infection in a third-tier Chinese city: relationship with gender, age, birth-year and survey years", *Microb Health Dis*, 2019; 1: e150.

7-Maria Paula Curado, Max Moura de Oliveira, Marcela de Araújo Fagundes, "Prevalence of *Helicobacter pylori* infection in Latin America and the Caribbean populations: A systematic review and meta-analysis", *Cancer Epidemiology*, 2019; 60, 141-148.

8-Bakka AS, El-Gariani AB, Abou Ghrara FM, Salih BA, "Frequency of *Helicobacter pylori* infection in dyspeptic patients in Libya", *Saudi Med J*, 2002; 23(10): 1261-1265.

9-Ahmed M. A. I, Altayeb M. A. A, Abdelrahman N. K, Abdurrahman N. A. M and Osman E. I., "Prevalence of *Helicobacter pylori* infection among Adults in Elfashir, North Darfur, Western Sudan", *International Journal of Biomedical and Advance Research*, 2020; 11(03): e5342.

10-Shuai Ren, Pengpeng Cai, Yaqian Liu, Tianpei Wang, Yan Zhang, Qian Li, Yuanliang Gu, Liqin Wei, Caiwang Yan, Guangfu Jin, "Prevalence of *Helicobacter pylori* infection in China: A systematic review and meta-analysis", *Journal of Gastroenterology and hepatology*, 2022; 37(3); 464-470.