

## Original Research Article

### MANAGEMENT OF WHITEFLY *BEMISIA TABACI* (GENN.) IN POTATO

#### **Abstract**

A field experiment was conducted at Potato Research Station, S. D. Agricultural University, Deesa during 2018-19 and 2019-20 for evaluation of efficacy of insecticides against whitefly in potato. The seven different treatments viz. T<sub>1</sub>: Control, T<sub>2</sub>: Seed treatment with imidacloprid (200 SL) @ 0.04% followed by foliar sprays of imidacloprid @60 gm a.i./ha at 85% emergence + second spray with thiamethoxam 25WG @100g a.i./ha, T<sub>3</sub>: Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence, T<sub>4</sub>: Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence followed by second spray of diafenthiuron after 10 days, T<sub>5</sub>: Foliar spray of castor oil @0.05% at 85% emergence, T<sub>6</sub>: Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05%, T<sub>7</sub>: Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05% by second spray with diafenthiuron after 10 days. Based on pooled data of two years, The significantly highest percent mortality (*i.e.* 68.57 %) after 2<sup>nd</sup> spray was recorded in T<sub>7</sub> *i.e.* foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05% by second spray with diafenthiuron after 10 days which was at par with T<sub>4</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence followed by second spray of diafenthiuron after 10 days). The tuber yield and benefit cost ratio were also recorded with T<sub>7</sub>.

**Keywords:** Potato, whitefly, management, difenthiuron, sucking pest

#### **Introduction:**

Potato (*Solanum tuberosum* Linn.) is an important vegetable crop cultivated throughout the world. It is a perennial herb belonging to family *Solanaceae*, a native of Andean Plateau of South America. The power of potato is known for sustaining millions of lives by providing nutritious food in the time of war and hunger and also used as a staple food in several countries of the world. Globally, it is ranked as the fourth most important food crop after maize, wheat and rice with approximately having double calories per hectare. It is maximum uses for making potato flour, potato chips, French fries, frozen potato, potato starch and tapioca of potato. Each 100 g serving of fresh potato, provides protein (1.6 g), carbohydrate (22.6 g), minerals (0.6 g), crude fibre (0.4 g), fat (0.10 g), vitamin C (25 mg) (Saini and Umrav, 2008). Potato contains significant levels of phenolic compounds and vitamin C as potent antioxidants (Brown, 2005), which inactivate reactive oxygen species, reduce oxidative damage, lead to improved immune functions and reduce risk of cardiovascular diseases, cancer, cataract, diabetes and ageing (Kour *et al.*, 2004). Recognizing the major role of potato as a staple food and providing food security, the year of

2008 was declared as the International Year of the Potato by the United Nations (Shailbala and Pathak, 2008). Hence, potatoes are called “King of Vegetables”.

This crop is attacked by several pests in the field and in store. According to Simpson (1977), potato is attacked by more than 100 arthropod pests among them 80 have been reported from India (Anonymous, 1971; Saxena and Rizvi, 1974; Rataul and Misra, 1979; Saxena and Misra, 1983). Among various insect-pests; aphids, jassids, cutworms, white grubs, epilachna beetles, defoliating caterpillars and potato tuber moths are serious and responsible for considerable economic losses, while the rest are sporadic and of minor importance. Among the sucking pests the whitefly, *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae), is a worldwide polyphagous insect pest that has wreaked havoc on agricultural productivity, particularly in some plant families such as Solanaceae, Cucurbitaceae, and Fabaceae (Oliveira *et. al.*, 2001 and Cruz-Estrada *et. al.*, 2013). The potato whitefly, *Bemisia tabaci* (Gennadius) are the most destructive ([Roditakis \*et al.\*, 2005](#)) pests of numerous crops ([Perring, 2001](#)). *Bemisia tabaci* biotype A (Gennadius) has the ability to transmit 111 economically important viruses ([Jones, 2003](#)). By keeping in mind notorious devastating sap-sucking insect pest this experiment was conducted for evaluation of various systemic insecticide against potato whitefly.

### **Materials and methods:**

The field experiment was conducted at Potato Research Station, S. D. Agricultural University, Deesa to study the management of whitefly in potato for two consecutive years Rabi 2018-19 and 2019-20. The experiment was laid out in a randomized block design with seven treatments and five replications with 3.0 m × 2.0 m plot size. The planting was done at row spacing of 50 cm and plant to plant spacing of 20 cm. The recommended dosage of fertilizers 275 kg N, 138 kg P and 275 kg K per hectare was applied. Insecticides were sprayed using high volume knapsack sprayer with 500 L ha<sup>-1</sup> solution. Seven different treatments *viz.*, T<sub>1</sub>: Control, T<sub>2</sub>: Seed treatment with imidacloprid (200 SL) @ 0.04% followed by foliar sprays of imidacloprid @60 gm a.i./ha at 85% emergence + second spray with thiamethoxam 25WG @100g a.i./ha, T<sub>3</sub>: Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence, T<sub>4</sub>: Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence followed by second spray of diafenthiuron after 10 days, T<sub>5</sub>: Foliar spray of castor oil @0.05% at 85% emergence, T<sub>6</sub>: Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05%, T<sub>7</sub>: Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05% by second spray with diafenthiuron after 10 days. Observations were taken on whitefly numbers before the foliar spray and on per cent reduction after 24, 48 and 72 hours of the spray on 5 tagged plants (3 leaves lower, middle and upper) as per method described by Butter and Vir (1990). The per cent reduction was work out by using standard formula.

### **Results and Discussion**

It was observed that there were non-significant differences in whitefly counts in all the treatment before first spray (Table 1). The significantly highest percent reduction (*i.e.*

56.35 %) after 1<sup>st</sup> spray during *rabi* 2018-19 was recorded in T<sub>7</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05% by second spray with diafenthiuron after 10 days) which was at par with T<sub>6</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05%) recorded 54.21% reduction, T<sub>3</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence) recorded 48.62% reduction and T<sub>4</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence followed by second spray of diafenthiuron after 10 days) recorded 46.79% reduction. The similar trend was also observed in pooled data while in *rabi* 2019-20, significantly higher reduction per cent were recorded in all the treatments except T<sub>5</sub> (Foliar spray of castor oil @0.05% at 85% emergence) over untreated control (Table: 1).

The significantly highest percent reduction (*i.e.* 62.70 %) after 2<sup>nd</sup> spray during *rabi* 2018-19 was recorded in T<sub>7</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05% by second spray with diafenthiuron after 10 days) which was at par with T<sub>4</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence followed by second spray of diafenthiuron after 10 days) and T<sub>2</sub> (Seed treatment with imidacloprid (200 SL) @ 0.04% followed by foliar sprays of imidacloprid @60 gm a.i./ha at 85% emergence+ second spray with thiamethoxam 25WG @100g a.i./ha). The next best treatment in sequence was T<sub>6</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05%) and T<sub>3</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence). In *rabi* 2019-20 significantly highest percent mortality (*i.e.* 74.44 %) after 2<sup>nd</sup> spray was recorded in T<sub>7</sub> which was at par with T<sub>4</sub> (69.68%). The next best treatment in sequence was T<sub>2</sub> (62.09 per cent mortality). The similar trend was also observed in pooled data. The zero per cent mortality was recorded in T<sub>5</sub> in both years as well as in pooled data. (Table: 2).

The economics of treatments shows that higher BC ratio (2.26) was recorded in T<sub>7</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence mixed with castor oil @0.05% by second spray with diafenthiuron after 10 days) which was followed by T<sub>4</sub> (Foliar spray of diafenthiuron 50WP 350 g a.i. at 85% emergence followed by second spray of diafenthiuron after 10 days). (Table : 3)

Present results are in agreement with findings of Yadav *et. al.*, (2022) they reported that two sprays of diafenthiuron 50WP 350 g a. i./ ha mixed with castor oil @ 0.05% at 10 days interval gave maximum mortality (82.0%), maximum tuber yield (24.7 t/ ha) with incremental cost benefit ratio (ICBR 1:2.1). Tomar *et. al.*, (2017) reported the highest reduction of potato whitefly was recorded in Difenthiuron 50 SC @ 350 g ai/ha (84.13%) in comparison to rest of insecticide. Meenu and Dahiya (2017) observed that diafenthiuron 50 WP @ 500 g/ ha showed better performance against *B. tabaci*. Bharpoda *et al.* (2014) found that diafenthiuron 50 WP (at 0.05%) was the most effective compared to imidacloprid 17.8 SL (at 0.008%) against *B. tabaci*. Rajasekhar *et al.* (2018) observed that diafenthiuron 50% WP @ 1.25 g/ l was found effective in cotton. Praveen *et al.* (2019) found that diafenthiuron 50%WP @ 1 g/ l and triazophos 40 % EC @ 2 ml/ l were found effective against whitefly in sunflower. Thumar *et. al.*, (2018) diafenthiuron 25% + pyriproxyfen 5% SE @ 1000 and 1250 ml/ha were found relatively more effective against sucking insect pests

viz., aphids, leafhoppers, whiteflies and thrips in Bt cotton. Two doses of diafenthiuron 25% + pyriproxyfen 5% SE (1250 and 1000 ml/ha) also gave highest seed cotton yield. Furthermore, previous reports demonstrated extreme effectiveness of diafenthiuron against whitefly (Naveed *et al.*, 2008; Reghuraman and Gupta., 2005; Shah *et al.*, 2007; Barrania and telab., 2014) on different crops.

**Table 1: Per cent reduction of potato whitefly after first spray (2018-19, 2019-20 & pooled).**

Treatment	1 <sup>st</sup> Spray					
	Nos before spray			Per cent reduction after spray		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T <sub>1</sub>	**24.73 *(17.60)	22.64 (15.00)	23.74 (16.30)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
T <sub>2</sub>	25.80 (19.00)	25.06 (18.00)	25.44 (18.50)	40.49 (42.40)	46.60 (52.80)	43.60 (47.60)
T <sub>3</sub>	25.02 (18.00)	23.32 (15.80)	24.20 (16.90)	44.19 (48.62)	50.47 (59.44)	47.30 (54.03)
T <sub>4</sub>	25.20 (18.20)	25.48 (18.60)	25.36 (18.40)	43.13 (46.79)	51.36 (60.93)	47.20 (53.86)
T <sub>5</sub>	27.34 (21.20)	27.34 (21.20)	27.34 (21.20)	17.03 (9.60)	19.35 (11.57)	18.42 (10.59)
T <sub>6</sub>	26.76 (20.40)	26.53 (20.00)	26.67 (20.20)	47.41 (54.21)	52.45 (62.71)	49.87 (58.46)
T <sub>7</sub>	26.53 (20.00)	26.76 (20.40)	26.66 (20.20)	48.65 (56.35)	51.75 (61.55)	50.15 (58.95)
<b>SEd</b>	1.51	1.55	1.31	3.06	3.00	2.07
<b>CD (0.05)</b>	NS	3.22	NS	6.35	6.23	4.29
<b>CV (%)</b>	9.20	9.70	8.11	14.04	12.22	8.91

\*\*Data are transformed values \*Data in the parenthesis are original values,

**Table 2: Per cent reduction of potato whitefly after second spray (2018-19, 2019-20 & pooled).**

Treatment	2 <sup>nd</sup> Spray					
	Nos before spray			Per cent reduction after spray		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled

T <sub>1</sub>	**0.00 *(0.00)	30.89 (26.40)	21.28 (13.20)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
T <sub>2</sub>	19.29 (11.00)	19.33 (11.00)	19.33 (11.00)	45.16 (50.28)	52.07 (62.09)	48.57 (56.18)
T <sub>3</sub>	17.17 (8.80)	16.93 (8.60)	17.07 (8.70)	24.94 (17.91)	15.97 (9.47)	21.61 (13.69)
T <sub>4</sub>	17.98 (9.60)	17.61 (9.20)	17.83 (9.40)	50.10 (58.69)	56.68 (69.68)	53.26 (64.18)
T <sub>5</sub>	26.75 (20.40)	27.39 (21.20)	27.09 (20.80)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
T <sub>6</sub>	16.92 (8.60)	18.00 (9.60)	17.53 (9.10)	27.17 (21.52)	20.43 (12.28)	24.13 (16.90)
T <sub>7</sub>	15.96 (7.60)	18.52 (10.20)	17.31 (8.90)	52.91 (62.70)	60.43 (74.44)	55.99 (68.57)
<b>SEd</b>	1.32	1.17	0.96	3.77	3.28	1.86
<b>CD (0.05)</b>	2.74	2.43	2.0	7.83	6.80	3.86
<b>CV (%)</b>	12.82	8.72	7.71	20.83	17.64	10.10

\*\*Data are transformed values \*Data in the parenthesis are original values,

**Table 3 : Economics of the treatments (pooled data)**

<b>Treatme</b>	<b>Yield (t/ha) pooled</b>	<b>Cost of cultivation (Rs./ha)</b>	<b>Produce (Rs./ha)</b>	<b>Sale Price (Rs./t)</b>	<b>Net returns (Rs./ha)</b>	<b>B:C ratio</b>
<b>T<sub>1</sub></b>	40.08	133900	360750	9000	226850	1.69
<b>T<sub>2</sub></b>	42.13	135440	379125	9000	243685	1.80
<b>T<sub>3</sub></b>	41.85	136700	376650	9000	239950	1.76
<b>T<sub>4</sub></b>	48.38	139500	435375	9000	295875	2.12
<b>T<sub>5</sub></b>	41.70	133950	375338	9000	241388	1.80
<b>T<sub>6</sub></b>	45.83	136750	412500	9000	275750	2.02
<b>T<sub>7</sub></b>	50.62	139550	455580	9000	316030	2.26

**Conclusion:**

It is concluded that the foliar spray of alone diafenthiuron 50WP @ 350 g a.i/ha or in combination with castor Oil @ 0.05% at 85% crop emergence and followed by at 10 days interval for the management of potato whiteflies & for higher net returns.

**References:**

- Anonymous. Pests of potato. Potato in India, CPRI Tech. 1971; Bull. No. 1: 25-26.  
 Barrania AA and Telab HKA. Relative efficacy of some newer insecticides on the mortality of jassid and whitefly in brinjal. Int. J. of Res. In. Biol. Sci. 2014;4(3):89-93.

- Bharpoda T M, Patel NB, Thumar RK, Bhatt NA, Ghetiya LV, Patel HC. Evaluation of insecticides against sucking insect pests infesting Bt cotton BG- II. The Bioscan. 2014; 9 (3): 977-980.
- Brown CR. Antioxidants in potato. American Journal of Potato Research. 2005; 82, 163-172.
- Butter NS, Vir BK. Sampling of whitefly, *Bemisia tabaci* Genn. In cotton. PAU Agricultural Research Journal. 1990; 27(4): 615-619.
- Cruz-Estrada A, Gamboa-Angulo M, Borges-Argáez R, Ruiz-Sánchez E. Insecticidal effects of plant extracts on immature whitefly *Bemisia tabaci* Genn.(Hemiptera: Aleyroideae). Electron. J. Biotechnol. 2013; 16, 6.
- Huded PB, Ghante, VN, Kulkarni V. Chemical Management of Whiteflies in Sunflower. International Journal of Current Microbiology and Applied Sciences. 2019; 8 (4): 2885-2890.
- Jones DR. Plant viruses transmitted by whiteflies. Eur. J. Plant Pathol. 2003; 109:95-219.
- Kaur C, George B, Deepa N, Singh B, & Kapoor HC. Antioxidant status of fresh and processed tomato-A review. Journal of Food Science and Technology, 2004; 41(5),479-486.
- Meenu and Dahiya KK. Efficacy of some insecticides against whitefly (*Bemisia tabaci*). International Journal of Applied and Natural Science, 2017;6(2):1-4.
- Naveed M, Saleem MA, Salam A. and Ali S. The effects of different groups of insecticides in cotton fields on *Bemisia tabaci*, Genn (Homoptera: Aleyrodidae), predators and parasitoids: Implication in whitefly management. Phytoparasitica. 2008;36: 377-387.
- Oliveira MRV, Henneberry TJ, Anderson P. History current status, and collaborative research projects for *Bemisia tabaci*. Crop Prot. 2001; 20,709–723.
- Perring TM. The *Bemisia tabaci* complex. Crop Prot. 2001;20:725-737.
- Raghuraman M, and Gupta GP, Field evaluation of neonicotinoids against whitefly *Bemisia tabaci* Gennadius in cotton.J. Entomol. 2005; 67(1): 29-33.
- Rajasekhar N, N. V. V. S. Durga Prasad, D.V. Sai Ram Kumar, Adinarayana, M. 2018. Incidence and management of cotton whitefly *Bemisia tabaci* under high density planting system (HDPS). International Journal of Current Microbiology and Applied Sciences 7(3): 2074-2079.
- Rataul HS And Misra SS. Potato pests and their control Pesticides;1979;13(7)27-38.
- Roditakis E, Roditakis NE, and Tsagkarakou A. Insecticide resistance in *Bemisia tabaci* (Homoptera: Aleyrodidae) populations from Crete. Pest Manage. Sci. 2005;61: 577-582.
- Saini GS and Umrav VK. Morden oleri and floriculture production. Rama publication house 2008;82-83.
- Saxena AP and Misra SS. Pests of potato crop in high altitude and their control. Proc. Wkshp. High Alt. Ent. & Wild). Ecol. Zoo/. Surv. India. 1983; 165-180.
- Saxena AP And Rizvi SMA. Insect-pests problems of potato in India. Journal of the Indian Potato Association. 1974;1:45-50.
- Shah MJ, Ahmad A, Hussain M, Malik MY and Ahmad B. Efficiency of different insecticides against sucking insect-pest complex and effect on the growth and yield of mungbean (*Vigna radiate* L.). Pak. Entomol. 2007;29(2):83-85.

- Shailbala & Pathak C. Harnessing the potential of potato to meet increasing food demand. *Kurukshetra*. 2008;56(3), 45-48.
- SIMPSON GW. Potato insects and their control. In O. Smith, Potatoes, Production, Storing and Processing. pp 505-605, AVI Publishing Co. Westport, Connecticut, U.S.A. 1977.
- Thumar RK, Borad PK, Pathan NP, Bharpoda TM, Saiyad, MM and Chaudhary HK. Bio-efficacy of diafenthiuron 25% + pyriproxyfen 5% SE against sucking insect pests of Bt Cotton, *Journal of Entomology and Zoology studies*. 2018; 6(5): 1024-1029.
- Tomar swati, Sharma Seema and Malik Kamlesh "Diafenthiuron (a novel insecticide) for management of whitefly, *Bemisia tabaci*.", *Indian Journal of Scientific Research*. 2017;16(1):120-122.
- Yadav DL, Patidar BK, and Nagar BL. Efficacy of Insecticides against Whitefly *Bemisia tabaci* (Genn.) in Potato. *Indian Journal of Entomology*. 2022;84(4), 900–902.