

Short Research Article

**COMMUNITY COMPOSITION AND DYNAMICS OF BUTTERFLIES IN DIFFERENT
MICROHABITATS OF CENTRAL ARAVALLI HILL REGIONS OF AJMER
DISTRICT, RAJASTHAN, INDIA**

ABSTRACT

Aims: Butterflies are the symbol and target species for conservation in many parts of the world, and are key indicators of an environment's ecological status. The abundance, brief generation time, quick movement, and sensitivity to climatic changes of the butterfly fauna make it a significant predictor. It is crucial for effective and suitable butterfly protection to conduct research on biodiversity, ecology, and habitat suitability.

Place and Duration of Study: The study was conducted in Ajmer City, Rajasthan, India. The four sites for butterfly collection were Open Land, Scrub Land, Aravalli Hills, and MDS University Campus to reflect the variety of environments in Ajmer. Scrub land makes up 1.5 km², open ground 1.3 km², the MDS University campus 0.5 km², and the Aravalli hills 2 km² of the 5.3 km² research regions.

Methodology: For butterfly studies, the Pollard walk technique was used, with 20 fixed transects placed stratified and randomly across four environments. Adult butterfly individuals were noted while walking at a slow, steady speed within a assumptive 5 m radius and Alpha and Beta diversity analysis was performed using PAST 4.06 and Microsoft Excel 2010

Results: During the study, 54 butterfly species from five groups were identified. The most diverse families were Nymphalidae and Pieridae, then Lycaenidae, Hesperidae, Papilionidae.

Conclusion: It is essential to track changes in the climate and human impacts because they are sensitive to changes in their surroundings. By conducting additional research into the study of novel and inventive butterfly diversity, the previously mentioned list needs to be updated. It is impossible to totally eradicate the detrimental effects of industrialization and sustainable development. Because they perform functions like pollinating various plant species, nutrient depletion, pest control, and decomposition, insects are crucial to a society's structure. However, by planting appropriate trees, plants, and other vegetation that will support the organisms' continued health, we can at least try to lessen them. At a minimum, this attempt will prevent the common species from facing extinction.

Keywords: Butterfly, Community Composition, Microhabitat, Rajasthan, Conservation.

1. INTRODUCTION

Ecosystems and habitats are always transforming, and some of these alterations are fueled by human influences. As a result, continuous ecological monitoring is required to assess the status of the ecosystem and biodiversity (Lindenmayer and Likens, 2018). We study biological

indicators to better understand biodiversity, and they help us understand the importance of ecosystems and their management (Maleque *et al.*, 2009). The human development process is one of several elements proven to have an impact on biodiversity in many regions. Construction operations such as road construction, power transmission line construction, energy development, and waterway excavation and impoundment for development have been shown to have an impact on land cover and on variety of species including butterflies (Gasparatos *et al.*, 2017).

Butterflies are a flagship and target species for conservation in many parts of the world, particularly for invertebrates (New, 2011). Butterflies are simple to examine because of their fairly large size and distinctiveness, as well as their well-known taxonomy (Brown, 1991), and are a key component of biodiversity in natural environments due to their strong complicated linkages in the food web (Bonebrake *et al.*, 2010). The adult forms pollinate, which is essential for the ecological functioning (Tiple *et al.*, 2006), and larval forms serve as primary consumers and prey to species at higher trophic levels, butterflies provide dual functions as pollinators and energy transferors. Butterflies are key indicators of an environment's ecological status and biotope quality (Launer and Murphy, 1994; Sharma *et al.*, 2020), as prospective pollinators of their nectar plants and indicators of the health and quality of their host plants and the ecosystem as a whole. The butterfly fauna is an important indicator because it is abundant, its generation is short, its migration is fast, and it is very sensitive to environmental changes (Lee and Kwon, 2014).

Many species are habitat specific and are extremely vulnerable to habitat degradation, which includes changes in microclimate, vegetation structure, and the co-occurrence of plant types on a local scale (Bonebrake *et al.*, 2010; Sharma *et al.*, 2020). Seasonal factors, in addition to habitat types, play an important role in defining the spatiotemporal patterns of butterfly species richness and diversity (Kunte, 1997; Sengupta *et al.*, 2014; Sharmila *et al.*, 2020). Temperature and rainfall patterns impact the similarity and variety of local butterfly populations (Grøtan *et al.*, 2012, 2014)

Butterflies are extremely sensitive to fractional fluctuation in climatic conditions and disturbance in their habitats due to their short life spans, host plant specialization, and limited dispersion capabilities (Kocher and Williams, 2000). The undisturbed natural flora and seasonal flowering plantation provide promising habitat for butterfly populations as there are no development operations or contamination from industrial hazardous waste (Tiple, 2012).

In this context, studies on diversity, ecology, and habitat appropriateness are critical for successful and proper butterfly conservation (Sharma *et al.*, 2020). Studies are vital for understanding butterfly diversity, ecology, and numerous functions in an ecosystem, as well as for investigating the influence of disturbance and land use changes on them (Bhardwaj *et al.*, 2012). Furthermore, monitoring and quantifying butterflies across seasons and habitats has significant implications for systematic conservation planning. All of these characteristics make them a viable model for ecological and conservation research, emphasizing the importance of establishing good conservation approaches (Ehrlich and Hanski, 2004). As an outcome of our current work, an inventory of butterfly faunal diversity and abundance is developed.

2. MATERIAL AND METHODS

2.1 Study Area

All of the investigation locations were in the Central Aravalli Hill Ranges of Ajmer City, Rajasthan, India. Ajmer has semi-arid climate with bushy and thorny vegetation (Singh *et al.*, 2022). The average annual rainfall in Ajmer is 525 mm, with 90% of it falling between June and September, while the average yearly temperature is 30-40°C. Four microhabitats for butterfly sampling were chosen to represent the diverse landscapes of Ajmer. The four ecologically diverse study landscapes were Open Land, Scrub Land which also includes some water bodies, Aravalli Hills, and MDS University Campus, which were located between the latitude and longitude of 26.50°N and 74.68°E.

Open Lands include areas near university with scattered vegetations which are chiefly grasses and herbs. Scrub Lands were dominated by dense scrubs and trees like *Prosopis juliflora*, *Calotropis procera*, *Prosopis cineraria*, *Lantana camara*, *Capparis decidua* and *Acacia senegal*. The rocky terrain of Aravalli Hill regions encompassing the regions of ancient central aravallies in Rajasthan, the chief vegetations are *Acacia arabica*, *Prosopis juliflora*, *Cynodon dactylon* etc. MDS University Campus is area with highest human activities among all and is most diverse in terms of both natural and ornamental flora that include *Azadirachta indica*, *Alstonia macrophylla*, *Acacia acacia*, *Calotropis gigantean*, *Ziziphus nummularia*, *Adhatoda vasica*, *Bougainvillea sp.* and invasive flora such as *Prosopis juliflora*, *Lantana sp.* and *Parthenium sp.* The total study area is 5.3 km² approximately, in which scrub land accounts for 1.5 km², open land for 1.3 km², MDS University campus for 0.5 km², and Aravalli Hills for 2 km².




Figure 1. Four studied microhabitats in study area located at Ajmer district, Rajasthan India.

2.2 Data Collection and Analysis

Sampling was carried out for a period of one year between November 2021 and October 2022. Pollard walk method was used for butterfly surveys (Pollard 1977). At each site, a road transect method was followed to collect the data. Each site was visited twice a month (N=96) and on each day sampling was carried out for 3 hours in morning (07:00 to 10:00 hrs) and 3 hours in evening (16:00 to 19:00 hrs).

The study area was explored during three seasons viz. summer (March-June), monsoon (July-October) and winter (November-February). Twenty permanent transects (N = 20), each of 300m in length were laid in a stratified and random manner across four habitats (N = 5 each) viz. in the study area. The consecutive transects in a habitat were spaced 50m apart from each other. Individuals of adult butterflies were recorded and counted around an imaginary 5m radius while walking with a slow, constant pace in each permanent transect. Ten transects (N = 10) were covered in each sampling day, mostly during cloudless and sunny weather conditions in order to spot maximum butterflies (Subedi *et al.*, 2020).

Comment [mm1]: Optional : This research would be better if there were comparison results between the number of species and individuals present during morning and evening sampling, so that we would be able to estimate that changes in temperature will provide an idea of changes (differences) in the number of species and the number of individuals

The identification of butterflies was done in the field based on morphological and behavioral descriptions with reference to Kehimkar (2016) followed by photography using DSLR Canon-EOS 1300D and Nikon D5300 (55-300mm).

Alpha and beta diversity were calculated using the following indices: Shannon Weiner Index, Simpson Index, Evenness Index, Berger Parker Index of Dominance, Margalef's Index for Richness, Whittaker's Beta Diversity was also compared pairwise for each of the four locations. Microsoft Excel 2010 and PAST 4.06 (Hammer *et al.*, 2001) were used for all statistical analysis.

3. RESULTS AND DISCUSSION

From the four study sites, 54 butterfly species from five families were identified over the course of the investigation (Table 1). With 16 species each (29.6%), Nymphalidae had the highest species richness, followed by Pieridae (14 species, 25.9%), Lycaenidae (11 species, 20.3%), Hesperidae (7 species, 12.9%), and Papilionidae (6 species, 11.1%). Whittaker's Beta Diversity index showed highest value of 0.36842 between University Campus and Aravalli Hills (Table 3).

Table 1: Distribution of the different butterfly species observed at the four study sites.

S.No	Family	Scientific Name	Study sites and number of sightings			
			Aravalli Hills	Scrub Forest	University Campus	Open Land
1	Hesperidae	<i>Hasora chromus</i>	0	10	30	22
2		<i>Sarangesa purendra</i>	18	31	40	22
3		<i>Spialia galba</i>	5	10	40	30
4		<i>Suastus gremius</i>	0	0	0	5
5		<i>Parnara guttata</i>	0	20	30	23
6		<i>Borbo cinnara</i>	0	0	20	11
7		<i>Pelopidas mathias</i>	12	20	31	11
8	Papilionidae	<i>Graphium doson</i>	0	2	3	2
9		<i>Graphium aganemnon</i>	1	5	10	20
10		<i>Pachliopta aristolochiae</i>	0	10	40	44
11		<i>Pachliopta hector</i>	0	10	40	33
12		<i>Papilio polytes</i>	20	25	70	55
13		<i>Papilio demoleus</i>	0	5	30	50
14	Pieridae	<i>Eurema brigitta</i>	0	22	30	27
15		<i>Eurema hecabe</i>	5	30	44	32
16		<i>Eurema laeta</i>	0	12	30	20
17		<i>Catopsilia pomona</i>	7	29	40	33
18		<i>Catopsilia pyranthe</i>	0	32	70	60
19		<i>Colotis amata</i>	0	19	49	33

20		<i>Colotis etrida</i>	20	40	80	44
21		<i>Colotis fausta fausta</i>	2	20	30	13
22		<i>Ixias marianne</i>	0	20	33	14
23		<i>Ixias pyrene</i>	1	2	5	2
24		<i>Pieris canidia</i>	0	2	12	8
25		<i>Cepora nerissa</i>	5	11	28	22
26		<i>Delias eucharis</i>	0	4	1	0
27		<i>Belenois aurota</i>	0	10	55	33
28	Lycaenidae	<i>Catochrysops strabo</i>	1	40	55	22
29		<i>Leptotes plinius</i>	0	15	45	33
30		<i>Tarucus nara</i>	5	20	100	91
31		<i>Zizeeria karsandra</i>	0	0	4	7
32		<i>Zizula hylax</i>	0	10	50	44
33		<i>Azonus uranus</i>	0	0	3	0
34		<i>Freyeria putli</i>	2	20	66	29
35		<i>Freyeria trochylus</i>	0	10	18	11
36		<i>Chilades pandava</i>	0	12	44	20
37		<i>Chilades parrhassius</i>	11	66	90	70
38		<i>Rapala iarbus</i>	0	0	10	0
39	Nymphalidae	<i>Trimala limniace</i>	5	20	5	15
40		<i>Danaus genutia</i>	0	0	3	1
41		<i>Danaus chrysippus</i>	50	59	130	40
42		<i>Euploea core</i>	0	3	5	11
43		<i>Melanitis leda</i>	0	0	1	1
44		<i>Melanitis phedima</i>	0	5	1	2
45		<i>Ypthima asterope</i>	20	100	200	140
46		<i>Acraea violae</i>	0	1	0	0
47		<i>Phalanta phalantha</i>	10	11	5	2
48		<i>Vanessa cardui</i>	50	43	30	37
49		<i>Junonia orithya</i>	77	60	140	108
50		<i>Junonia hierta</i>	50	16	70	33
51		<i>Junonia almana</i>	30	28	120	100
52		<i>Junonia lemonias</i>	23	33	56	43
53		<i>Hypolimnas bolina</i>	0	19	28	25
54	<i>Hypolimnas misippus</i>	0	6	20	10	

Table 2: Alpha diversity indices of butterflies recorded at four study sites.

	Aravalli Hills	Scrub Forest	University Campus	Open Land
Total Sightings	430	998	2190	1564
Species Richness				

Species	24	47	52	50
Margalef's Richness	3.793	6.661	6.631	6.662
Evenness				
Pielou's Evenness Index	0.5965	0.7095	0.6738	0.6969
Dominance				
Dominance_D	0.09157	0.03873	0.03627	0.03673
Berger-Parker	0.1791	0.1002	0.09132	0.08951
Diversity				
Simpson_1-D	0.9084	0.9613	0.9637	0.9633
Shannon_H	2.661	3.507	3.556	3.551

Table 3: Pairwise comparison of Whittaker's Beta Diversity of butterflies recorded in four study sites

	Aravalli Hills	Scrub Forest	University Campus	Open Land
Aravalli Hills	0	0.32394	0.36842	0.35135
Scrub Forest	0.32394	0	0.070707	0.072165
University Campus	0.36842	0.070707	0	0.039216
Open Land	0.35135	0.072165	0.039216	0

Butterflies are an important part of the food chain and in addition, they serve as markers of environmental changes in their surroundings. They have greater sensitivity than a lot of other biological groups (Thomas, 2005). Two species of Lyncaenidae, *Azonus uranus* and *Rapala jarbus* were found only in one study area that was University campus with mixed vegetation. One species of Hesperidae (*Suastus gremius*) was only found at open land with very low vegetation and one species of Nymphalidae (*Acraea violaei*) was only found at scrub forest dominated with scrubby vegetation. Because of high availability of food plants, 24 species (44.4% of total), 3 of Hesperidae, 4 of Lycaenidae, 9 of Nymphalidae, 2 of Papillionidae and 6 of Pieridae were noticed in all study sites, There are 9 species of Nymphalidae that use every study habitat, which suggests that they are strong, energetic flyers that may aid them in looking for resources over wide regions (Eswaran and Pramod, 2005; Krishnakumar *et al.*, 2008). According to Table 2, there is more species richness species in scrub forest with water bodies than in open land with little vegetation followed by university campus with a variety of vegetation, and the Aravalli hills in that order. Alpha species diversity (α) is highest at university campuses with mixed vegetation have high Simpson (0.9637) and Shannon (3.556) indices

values, while Aravalli Hills have the lowest diversity according to both indices (0.9084 and 2.661). However, Aravalli Hills (0.09157) exhibits higher Dominance than other locations.

The highest Whittaker's Beta diversity is between the university campus and the Aravalli hills (0.36842), and the lowest is between the university campus and open land (0.039216) (Table 3).

Under Wild Life Protection Act, 1972 one species *Pachliopta hector* (Crimson rose) of Papilionidae family is classified in Schedule-I. Two species *Hypolimnas misippus* (Danaid eggfly) of family Nymphalidae and *Cepora nerissa* (Common gull) of Pieridae family in Schedule-II and one species *Euploea core* (Common crow) of Nymphalidae in Schedule-IV are valuable from conservation point of view.

4. CONCLUSION

From the study sites, a total of 54 species of butterflies have been documented. Among the five recorded butterfly families in the four studied microhabitats, Nymphalidae appears to be the most diverse not only in abundance but also in species richness (n=16, 29.6%), followed by Pieridae (n=14, 25.9%), Lycaenidae (n=11, 20.3%), while families with lowest species richness are Hesperidae (n=7, 12.9%) and Papilionidae (n=6, 11.1%). The diversity of Nymphalidae and Pieridae is greater in terms of species abundance in the study area, according to our present outcomes, which are followed by Lycaenidae. The two families Hesperidae and Papilionidae showed the least diversity. The information gathered in this study's data could prove invaluable as a resource for estimating how the climate in the area will change in the near future. The depicted butterfly family and species list is not final and comprehensive, so long-term research work with periodic surveys of the vegetation cover and tracking of the diversity of butterflies may be performed in the study area with a focus on ecological aspects. Due to their sensitivity to environmental changes, it is crucial to monitor changes in the climate and human effects. The aforementioned list needs to be updated through further research into the study of new and innovative butterfly diversity. The negative effects of sustainable development and industrialization cannot be completely eliminated. Insects play an important role in the structure of a society by providing services like pest control, nutrient depletion, and decomposition, as well as pollinating different plant species. But we can at least try to reduce them through planting suitable trees, plants, and other vegetation that will help the organisms stay healthy. By making such an attempt, at the very least, the common species won't be on the verge of extinction.

REFERENCES

- Bhardwaj, M., Uniyal, V.P., Sanyal, A.K. & Singh, A.P. 2012. Butterflies communities along an elevational gradient in the Tons valley, Western Himalayas: implications of rapid assessment for insect conservation. *Journal of Asia-Pacific Entomology* 15: 207-217. <https://doi.org/10.1016/j.aspen.2011.12.003>
- Bonebrake, T.C., Ponisio, L.C., Boggs, C.L. & Ehrlich, P.R. 2010. More than just indicators: a review of tropical butterfly ecology and conservation. *Biological Conservation* 143(8): 1831-1841. <https://doi.org/10.1016/j.biocon.2010.04.044>.
- Brown, Jr. K.S. 1991. Conservation of neotropical environments: Insects as indicators. In Collins, N.M. & Thomas, J.A. (Eds), *The Conservation of Insects and their Habitats*, Academic Press, London. pp 349-404.
- Ehrlich, P.R. & Hanski, I. 2004. *On the wings of checkerspots: a model system for population biology*. Oxford University Press, Oxford.
- Eswaran, R. & Pramod, P. 2005. Structure of butterfly community of Anaikatty hills, Western Ghats. *Zoos' Print Journal* 20(8):1939-1942.
- Gasparatos, A., Doll, C.N., Esteban, M., Ahmed, A. & Olang, T.A. 2017. Renewable energy and biodiversity: implications for transitioning to a Green Economy. *Renew Sustain Energy Rev* 70:161-184
- Grøtan, V., Lande, R., Chacon, I.A. & DeVries, P.J. 2014. Seasonal cycles of diversity and similarity in a Central American rainforest butterfly community. *Ecography* 37: 509-516. <https://doi.org/10.1111/ecog.00635>
- Grøtan, V., Lande, R., Engen, S., Sæther, B.E. & DeVries, P.J. 2012. Seasonal cycles of species diversity and similarity in a tropical butterfly community. *Journal of Animal Ecology* 81: 714-723. <https://doi.org/10.1111/j.1365-2656.2011.01950.x>
- Hammer, Ø., Harper, D.A.T. & Ryan, P.D. 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica* 4(1):1-9.
- Kehimkar, I. 2016. *The Book of Indian Butterflies*. Bombay Natural History Society, Oxford University Press, Mumbai.

- Kocher, S.D. & Williams, E.H. 2000. The diversity and abundance of North American butterflies, vary with habitat disturbances and geography. *Journal of Biogeography* 27: 785-794. <https://doi.org/10.1046/j.1365-2699.2000.00454.x>
- Krishnakumar, N., Kumaraguru, A., Thiyagesan, K. & Asokan, S. 2008. Diversity of papilionid butterflies in the Indira Gandhi Wildlife Sanctuary, Western Ghats, Southern India. *Tigerpaper* 35(1):1-8.
- Kunte, K. 1997. Seasonal patterns in butterfly abundance and species diversity in four tropical habitats in northern Western Ghats. *Journal of BioSciences* 22(5): 593-603. <https://doi.org/10.1007/BF02703397>
- Launer, A.E. & Murphy, D.D. 1994. Umbrella species and the conservation of habitat fragments: a case of a threatened butterfly and a vanishing grassland ecosystem. *Biological Conservation* 69: 145-153. [https://doi.org/10.1016/0006-3207\(94\)90054-X](https://doi.org/10.1016/0006-3207(94)90054-X)
- Lee, C. M. & Kwon T. 2014. Change of butterfly communities after clear cutting in Gwangneung Forest. *Korean Journal of Applied Entomology* 53 (4), 347–354.
- Lindenmayer, D.B. & Likens, G.E. 2018. *Effective ecological monitoring*. CSIRO publishing, Australia. 210 pages.
- Maleque, M. A., Maeto, K. I. & Hiroaki, T. 2009. Arthropods as bioindicators of sustainable forest management, with a focus on plantation forests. *Applied Entomology and Zoology* 44 (1), 1–11.
- New, T.R. 2011. Launching and steering flagship Lepidoptera for conservation benefit. *Journal Threatened Taxa* 3(6):1805-1817. <https://doi.org/10.11609/JoTT.o2621.1805-17>
- Pollard, E. 1977. A method for assessing changes in the abundance of butterflies. *Biol Conserv* 12:115–134.
- Sengupta, P., Banerjee, K.K. & Ghorai, N. 2014. Seasonal diversity of butterflies and their larval food plants in the surroundings of upper Neora Valley National Park, a sub-tropical broad leaved hill forest in the eastern Himalayan landscape, West Bengal, India. *Journal of Threatened Taxa* 6(1): 5327-5342. <http://dx.doi.org/10.11609/JoTT.o3446.5327-42>
- Sharma, K., Acharya, B.K., Sharma, G., Valente, D., Pasimeni, M., Petrosillo, I. & Selvan, T. 2020. Land use effect on butterfly alpha and beta diversity in the Eastern Himalaya, India. *Ecological Indicators* 110: 105605. <https://doi.org/10.1016/j.ecolind.2019.105605>

- Sharmila, E.J., Thatheyus, A.J., Susaritha, S. & Snegapriya, M. 2020. Seasonality of butterflies in Alagar Hills reserve forest, India. *Entomon* 45 (1): 53-60. <https://doi.org/10.33307/entomon.v45i1.503>
- Singh, P., Choudhary, R., Sharma, V. & Chandra, S. 2022. Community composition and microhabitat preference by Saurian fauna of MDS University campus and nearby areas of central Aravalli foothills. *International Journal of Ecology and Environmental Sciences*. Volume 4, Issue 2, 2022, Page No. 1-5
- Subedi, B., Stewart, A., Neupane, B., Ghimire, S. & Adhikari, H. 2020. Butterfly species diversity and their floral preferences in the Rupa wetland of Nepal. *Ecology and Evolution* 11: 2086-2099. <https://doi.org/10.1002/ece3.7177>
- Thomas, J.A. 2005. Monitoring change in the abundance and distribution of insects using butterflies and other indicator groups. *Philosophical Transactions of the Royal Society London Biological Sciences* 360(1454):339-357. <https://doi.org/10.1098/rstb.2004.1585>
- Tiple, A.D. 2012. Butterfly species diversity, relative abundance and status in Tropical Forest Research Institute, Jabalpur, Madhya Pradesh, central India. *Journal Threatened Taxa* 4(7):2713-2717. <https://doi.org/10.11609/JoTT.o2656.2713-7>
- Tiple, A.D., Deshmukh, V.P. & Dennis, R.L.H. 2006. Factors influencing nectar plant resource visits by butterflies on a university campus: implications for conservation. *Nota Lepidopteralogica* 28(3/4): 213-224.