

# Estimation of Nitrogen in Plant Leaves Using SPAD

## ABSTRACT

A chlorophyll meter measures the greenness of leaves and can indicate the chlorophyll content and N status. Therefore, by using a chlorophyll meter, information related to the leaf N content could be obtained and this would avoid the need of laborious laboratory analysis. Keeping in this view a field experiment was conducted during winter (Rabi) 2019 growing season at the experimental field of JNKVV, Jabalpur to estimate crop nitrogen (N) and chlorophyll content in wheat using SPAD. The treatment comprised of 3 sources of nutrient M1- Inorganic sources (NPK fertilizers), M2- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and M3- Integrated sources (50% Inorganic + 50% organic) as main treatments and 5 NPK levels S1- control, S2- 100% RDF, S3- 150% RDF, S4- 200% RDF, S5- Based on soil test value for target yield of 6t ha<sup>-1</sup> as sub treatments were replicated thrice in a split plot design. The results showed that the correlation ( $r = 0.94, 0.90, 0.92$  and  $0.94$ , respectively) and coefficient of determination ( $R^2 = 0.96, 0.94, 0.83$  and  $0.94$ , respectively) was significantly and positively correlated with SPAD and nitrogen concentration in leaf. Similarly significant and positive correlation ( $r = 0.84, 0.76, 0.78$  and  $0.71$ , respectively) found with SPAD and total chlorophyll values. Coefficient of determination ( $R^2$ ) SPAD with chlorophyll and leaf N concentration proved to be significant ( $R^2 = 0.75, 0.76, 0.54$  and  $0.85$ , respectively) at different growth stages. It was concluded that SPAD meter are useful for the determination of chlorophyll and total nitrogen content of wheat plant.

*Keywords: SPAD, Chlorophyll, total N, Inorganic, INM, organic*

## 1. INTRODUCTION

Nitrogen (N) is a key constituent of all living cells and is essential for the growth and development of plants [1]. The most important role of N in the plant is its presences in the structure of protein, the most important building substances from which the living material or protoplasm of every cell is made. In addition, nitrogen is also found in chlorophyll, the green colouring matter of leaves. Chlorophyll enables the plant to transfer energy from sunlight by photosynthesis. Therefore, the nitrogen supply to the plant will influence the amount of protein, protoplasm and chlorophyll formed [2]. Nitrogen is one of the major nutritional elements that limits crop yields. Insufficient N supply results in smaller leaf area [3,4] and lower leaf photosynthesis, chlorophyll content and biomass production [4] leading to the loss of yield and quality.

On the other hand, excessive use of N fertilizer increases not only production cost but also environmental pollution [5]. Increased N fertilizer application does not always imply an increase of grain yield; on the contrary, even a decrease in crop yields is seen in poorly managed production systems [6]. In the quest of achieving high yield of wheat, farmers tend to apply nitrogen in excess of the requirements. It is well known that excess N availability for

crops like wheat and barley can adversely affect plant health, yield, values of the final product and environmental pollution [7]. As [8] showed, about 67% of total applied N fertilizer in wheat production could be lost because of natural processes like denitrification, leaching, volatilization (NH<sub>3</sub>), and greenhouse gas (N<sub>2</sub>O, NO or NO<sub>2</sub>) emissions. Nitrogen application at the right amount contributes to higher yields [9,10]. Leaf chloroplasts contain 70% of leaf N concentration [11] thus, the amount of chlorophyll present in plant leaves is often well correlated with leaf N [12,13]. Assessment of crop N requirement is critical for optimum N fertilizer application [14]. The optimum rate and application timing of Nitrogen (N) fertilizer are crucial in achieving a high yield however, conventional laboratory testing of plant nutrients is time-consuming and expensive [15]. Plant tests performed with optical instruments can be a significant simplification of the methods applied to attain precise determination of the nutritional status of plants during a growing season [16]. As leaf nitrogen content is strongly correlated with chlorophyll content the use of Soil Plant Analysis Development (SPAD) meter has been introduced as a popular, fast, and cheap technique to estimate N levels from the measurement of leaf transmittance [15]. Many studies suggest that a positive correlation between N uptake, leaf N concentration, leaf chlorophyll content, and grain yield [17, 18, 19, 20]. In this regard, [21] suggested that the SPAD chlorophyll meter can be used as an alternative to nitrogen nutrition index (NNI) to measure N status in wheat. Therefore, SPAD values have been successfully used for N fertilizer management in rice [22, 23] wheat [22,24] and maize [25]. The device can be used very simply and effectively for decision making in supplementary application of N fertilizer in wheat field. Wheat is very sensitive to insufficient nitrogen and very responsive to N fertilization [26]. The chlorophyll meter is faster than tissue testing for N and allows “fine tuning” of N management to field conditions, and consequently reduces risk of under- or over fertilizing the wheat crop [27]. There is no doubt that the nitrogen management practices can be adopted by crop growers if the practices are affordable, accurate in fertilization, easy to use, and environmentally friendly. The SPAD meter has been demonstrated and proved to be a viable and effective tool in developing and evaluating management recipes across a field [28]. Keeping in this view, the objective of this work was to evaluate the potential for Minolta SPAD 502 chlorophyll meter readings in monitoring leaf chlorophyll content and its relation to N concentration in wheat leaves.

## **2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY**

### **2.1 Experiment details**

A field experiment was conducted during winter (Rabi) 2019 growing season at the experimental field of JNKVV, Jabalpur. (23° 13' North latitude, 79° 57' East longitudes; 393 m elevation). The soil of the experimental site was Typic Haplustert, clay in texture neutral in reaction, non-calcareous, medium in organic carbon content, medium in available nitrogen, phosphorus, and potassium and low in DTPA extractable Zn. The treatment comprised of 3 sources of nutrient M1- Inorganic sources (NPK fertilizers), M2- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and M3- Integrated sources (50% Inorganic + 50% organic) as main treatments and 5 NPK levels S1- control, S2- 100% RDF, S3- 150% RDF, S4- 200% RDF, S5- Based on soil test value for target yield of 6t ha<sup>-1</sup> as sub treatments were replicated thrice in a split plot design. The wheat crop (GW-366) sown on with spacing of 22.5 cm row to row.

The experimental treatments were arranged in split plot design with three replications. Each experimental plot was 4 m long and 3 m wide consisting of three rows with one m inter-row spacing. The plots were kept weed-free and maintained in an optimum soil moisture regime throughout the experimental period. The specific quantity of each fertilizer was calculated based on gross plot size as per treatment. Treatment wise 1/3 quantity of nitrogen and entire

dose of phosphorus and potassium were applied as basal. While, the 1/3 dose of nitrogen was top dressed in two equal splits, after CRI and third irrigation (65 DAS). Urea (46% N), Single super phosphate (16% P<sub>2</sub>O<sub>5</sub>) and muriate of potash (60% K<sub>2</sub>O) were used as sources of nitrogen, phosphorus and potassium, respectively. FYM, VC and biofertilizers were applied as per treatment 15 days before sowing. The observations were recorded from each plot at crown root initiation (CRI) (21 DAS), tillering (45 DAS), flowering (65 DAS) and milking stages (90 DAS) of wheat.

### **2.3 Estimation of Chlorophyll using SPAD-502**

A chlorophyll meter (SPAD-502, Minolta, Japan) was used to obtain readings estimating leaf chlorophyll concentration (SPAD value). Five plants per treatment were selected randomly and SPAD values were recorded from the fully matured leaves counted from the top of the plants, the youngest fully expanded leaf. The observations were recorded from each plot at crown root initiation (CRI) (21 DAS), tillering (45 DAS), flowering (65 DAS) and milking stages (90 DAS).

### **2.4 Nitrogen analysis in plant**

The nitrogen content in plant sample was estimated by micro Kjeldahl digestion and distillation method using KEL PLUS system. For digestion, a known weight of plant sample in the presence of sulphuric acid with catalyst mixture (5:1) under high temperature was digested and distilled in presence of sodium hydroxide, the released ammonia was condensed and absorbed in boric acid with mixed indicator (Bromocresol cresol green and methyl red indicator) to form ammonium borate, the excess of which was titrated with standard sulphuric acid (0.1N) (AOAC, 1995).

### **2.5 Statistical analysis:**

Data were analyzed using SPSS for analysis of variance and Fisher's LSD multiple range test was employed for the means comparisons.

## **3. RESULTS AND DISCUSSION**

### **3.2 SPAD readings at different growth stages**

It can be seen from the data (Table 2) that the SPAD reading, or chlorophyll content index (CCI) was significantly maximum (8.15, 12.38, 28.58 and 30.65, respectively) with the application of inorganic sources of nutrient (M1) over integrated (M3) and organic sources of nutrient (M2) at CRI, tillering, jointing and milking stages, respectively. The SPAD reading of 6.36, 9.71, 25.54 and 25.00 were registered in M3 and it was also found significantly superior over M2 (4.47, 6.74, 17.26 and 16.65, respectively) at each stage, respectively.

However, the application of S3 (150% RDF) and S4 (200% RDF) significantly increased SPAD reading over S2 (100% RDF) and S5 (STV based RDF) at CRI and jointing but at par at tillering. S4 Significant S2 (100% RDF) and S5 (STV based RDF) but at par with S3. At milking. The highest SPAD reading was registered in S4 (200% RDF) which was 7.97, 11.07, 27.45 and 28.02 and 7.29, 11.11, 26.60 and 27.24 with 150% RDF at each stage, respectively.

The interaction effect was found to be significant at CRI stage of wheat. The combination of M1S4 (10.50) was recorded the highest SPAD reading, which was significantly superior over all the combination and followed by M1S3 (9.23), M1S2 (8.40), M1S5 (8.40), M3S4 (7.93)

and M3S3 (7.52) all were at par among themselves. At tillering stage, the combination M1S5 (16.11), M1S4 (15.64), M1S3 (15.28) were found significant over all combination. On comparing the treatment combination at jointing stage, applied to wheat, performance of M1S3 (33.70), M1S4 (33.48), M1S5 (33.45) were found significant over all combinations. However, at milking stage, the interaction of M1S4 (38.92) causes significant different and statistically similar treatment M1S3 (32.44) and M1S5 (32.86), M1S2 (31.86).

Maximum SPAD reading were obtained about jointing stage (65 DAS) then the chlorophyll content gradually decreased till the end of the vegetation season similar results also reported by Juško et al [29]. Maximum SPAD reading found in inorganic fertilizers because there was more nitrogen available to the NPK treated plants, there was more chlorophyll in the leaves, which speeds up photosynthesis [30]. Our results show that the application of nitrogen in inorganic form at the right time enhances chlorophyll content which resulted to higher SPAD values [15]. Nitrogen is part of the enzymes associated with chlorophyll synthesis [31] and the chlorophyll concentration reflects relative crop nitrogen status and yield level [32]. Very close link between chlorophyll and nitrogen content was observed as investigated by many investigators [33, 34]. It is understandable, because nitrogen is a structural element of chlorophyll and protein molecules, and thereby affects formation of chloroplasts and accumulation of chlorophyll in them [35, 36]. Our results supported by Islam et al. 2014 [37] reported that the SPAD values drastically reduced in treatments treated with only organic fertilizers. The values were higher in treatment receiving full dose of NPK through inorganic fertilizers as compared with reduced inorganic fertilizers and/or organically supplied fertilizers. These results agree with that obtained [37] in potato. This might be because N supply in organic treatment is generally restricted for slow N mineralization as compared to crop N demand [38].

### **3.2 Total chlorophyll in leaves (mg g<sup>-1</sup> leaf tissue) at different growth stages**

Table 3 showed that the total chlorophyll of 1.27, 1.92, 2.14 and 1.24 mg g<sup>-1</sup> leaf tissue was significantly more where, inorganic sources of nutrient (M1) was applied followed by integrated sources of nutrients (M3) with value of 1.08, 1.75, 1.78 and 1.04 mg g<sup>-1</sup> leaf tissue over organic sources of nutrient (M2) which recorded lowest total chlorophyll of 0.93, 1.33, 1.41 and 0.87 mg g<sup>-1</sup> leaf tissue at CRI, tillering, jointing and milking stages,.

However, the application of S3 (150% RDF) and S4 (200% RDF) significantly increased total chlorophyll over S2 (100% RDF) and S5 (STV based RDF) at each growth stages. The highest total chlorophyll was recorded with the application of S4 at CRI and tillering stage which was statistically similar with S3 (150% RDF) with the values of 1.26 and 1.84 mg g<sup>-1</sup> leaf tissue at CRI and tillering stage. However, S3 (150% RDF) was highest at jointing and milking stage with the values of 1.35 and 1.86 mg g<sup>-1</sup> leaf tissue. Treatment S2 (100% RDF) was found significant over S5 (STV based RDF) at tillering, jointing and milking stage but at par at CRI stage.

The interaction effect was found to be significant at CRI stage of wheat. The combination of M1S3 (1.61 mg g<sup>-1</sup> leaf tissue) was recorded the highest CCI, which was significantly superior over all the combination and at par with M1S4 (1.54 mg g<sup>-1</sup> leaf tissue). At tillering stage, the combination M1S4 (2.27) was found maximum and significant over all combination and at par with M1S3 (2.16 mg g<sup>-1</sup> leaf tissue). On comparing the treatment combination at jointing stage, applied to wheat, performance of M1S4 (2.58 mg g<sup>-1</sup> leaf tissue), was found significant over all combinations. However, at milking stage, the interaction of M1S4 (1.54 mg g<sup>-1</sup> leaf tissue) cause significant different and statistically similar treatment M1S3 (1.38 mg g<sup>-1</sup> leaf tissue).

The maximum chlorophyll in inorganic sources of nutrients is usually immediately and fast containing all necessary nutrients that are ready for plants [39]. Earlier study [40] indicated that abundant nutrient supplementation increases chlorophyll a concentration in Sorghum bicolor and Zea mays.

However, organic sources of nutrient perform poorly this may be due to disappearance of mineral N in the soil system was caused by microbial immobilisation under organic sources [41] which cause lower availability of N and chlorophyll. Organic fertilizer improves physical, biological, and chemical properties of a soil but the nutrients may not be as readily available to the plants [39].

### **3.3 Total N content in leaf at different growth stages**

Data presented in table 4 showed that the total N content in leaves of 2.39, 2.29, 1.80 and 1.38% was significantly more where, inorganic sources of nutrient (M1) was applied followed by integrated sources of nutrients (M3) with value of 2.26, 2.15, 1.76 and 1.34% over organic sources of nutrient (M2) which recorded lowest total chlorophyll of 2.22, 2.13, 1.60 and 1.19% at CRI, tillering, jointing and milking stages. However, M1 is significant over M3 except milking stage.

However, the application of S3 (150% RDF) and S4 (200% RDF) significantly increased N content over S2 (100% RDF) and S5 (STV based RDF) at CRI, jointing and milking stage. S3 (150% RDF) and S4 (200% RDF) were found at par among themselves. All levels were found at par among themselves.

The interaction effect was found to be significant at jointing and milking stage of wheat. The combination of M1S4 (2.13%) was recorded as the highest N content, which was significantly superior over all the combination and at par with M1S3 (2.04%) at jointing stage. However, at milking stage, the interaction of M1S4 (1.56%) causes significant different and statistically similar treatment M1S3 (1.45%) and M1S5 (1.48%)

The percentage of total N concentration in dried leaf increased as larger and larger N doses in mineral fertilization were applied. The pattern of nitrogen accumulation during the vegetation season deviated from the pattern of chlorophyll synthesis. The highest N accumulation was observed during the first measurement. Subsequent analyses showed lesser N concentration in the leaves. The lowest N concentration was noted later stages, and then it remained roughly at this level till the end of the vegetation season. The fact that the N concentration stayed roughly at the same level throughout the whole period till the end of the vegetation season, while chlorophyll content in the leaves fell, points to the accumulation of nitrogen compounds other than pigments [28].

Maximum N concentration in inorganic fertilizers may be because inorganic fertilizers releases nutrients for the plants instantly and in readily available forms for the plants during its growth, development and reproductive phases where the nutrient demand is at its peak. Higher levels of inorganic fertilizers have been increases the activity of photosynthesis and enzymes which responsible for transformation of energy, carbohydrates, fat metabolism and respiration of plant [42]. The nitrogen containing organic compounds in organic manures especially in FYM are more resistant to decomposition and only about one third of the nitrogen is easily released. The remaining amount of nitrogen persisted in the soil for a long period [43].

### **3.4 Correlation and Coefficient of determination of SPAD readings with total chlorophyll and total N content in leaves at different growth stages**

Pearson correlation coefficients between total N concentration in leaf, total chlorophyll content, SPAD values at CRI, tillering, jointing and milking stages of wheat crop are presented in Table 1 and fig 1. Good correlations were found for both meters for determination of chlorophyll content and N content. The SPAD values were positively correlated with grain yield of wheat at different good stages of plants. The SPAD values were also significantly correlated ( $r = 0.94, 0.90, 0.92$  and  $0.94$ , respectively) with leaf N concentration at different growth stages in this study. Maximum correlation was found in CRI and milking stage followed by jointing and tillering respectively. Coefficient of determination  $R^2$  between chlorophyll content in SPAD units and nitrogen concentration and leaf N concentration proved to be significant ( $R^2 = 0.96, 0.94, 0.83$  and  $0.94$ , respectively respectively). Highest value of Coefficient of determination was found in CRI and followed by milking, jointing and tillering, respectively. A close correlation ( $r = 0.93$ ) was established between the measurements of chlorophyll content using the N-tester and those obtained by laboratory tissue testing, for soybean, rice and spring wheat leaves [44], potato ( $r = 0.97$ ) [45], rice [46]. Also, Piekielek and Fox [47], Smeal and Zhang [48], Waskom et al. [49] confirm that the measurement results obtained using the SPAD optical method reflect the actual chlorophyll content in various crop plants.

Table showed that the relationship between leaf chlorophyll and SPAD value at different growth stages. The SPAD values were also significantly correlated ( $r = 0.84, 0.76, 0.78$  and  $0.71$ , respectively) with leaf chlorophyll at different growth stages in this study. Maximum correlation was found in CRI and jointing followed by tillering and milking stage, respectively. Coefficient of determination  $R^2$  between chlorophyll content in SPAD units and nitrogen concentration and leaf N concentration proved to be significant ( $R^2 = 0.75, 0.76, 0.54$  and  $0.85$ , respectively). Highest value of Coefficient of determination was found in milking and followed by tillering, CRI and jointing, respectively.

#### 4. CONCLUSION

Since SPAD readings are closely related to leaf N content, the SPAD meter can be used to monitor the N status of rice and thereby to adjust the rate of N fertilization to increase N use efficiency [50,51]. Data collected at different growth stages were used to determine when in the season SPAD data can be used to predict leaf N amount and future crop N need. The increasing of SPAD reading values with growth stage could be observed in this study. Although a chlorophyll meter (SPAD) has become a simple, quick and portable diagnostic tool for monitoring leaf N status and improving the timing of N topdressing in wheat.

#### REFERENCES

1. Govindasamy P, Muthusamy SK, Bagavathiannan M, Mowrer J, Prasanth Tej Kumar Jagannadham PTK, Maity A, Halli HM, Sujayanad GK, Vadivel R, Das TK, Rishi Raj, Pooniya V, Babu S, Rathore SS, Muralikrishnan L, Tiwari G. Nitrogen use efficiency—a key to enhance crop productivity under a changing climate. *Frontiers in Plant Science. Sec. Plant Nutrition*. 2023. (14): 1-19.
2. Alexandre C, Mamadou B, Rebouh NY, Valentinovich VV, Polityko PM. Nitrogen Fertilizer Influence on Winter Wheat (*Triticum aestivum* L.) Grain Yield and Grain Quality Production under Non-Chernozem Soils of Central Russia: A Review. *Agrotechnology*. 2022; 11: 288.
3. Fernandez CJ, McInnes KJ, Cothren JT. Water status and leaf area production in water- and nitrogen-stressed cotton. *Crop Science*; 1996; 36: 1224-1233.

4. Zhao D, Oosterhuis DM. Nitrogen application effect on leaf photosynthesis, nonstructural carbohydrate concentrations and yield of field-grown cotton. In D.M. Oosterhuis ed., AAES Special Report. Proceedings of the 2000 Arkansas Cotton Research. 2000; 198: 69-71.
5. Jaynes DB, Colvin TS, Karlen DL, Cambardella CA, Meek DW. Nitrate loss in subsurface drainage as affected by nitrogen fertilizer rate. *Journal of Environmental Quality*. 2001; 30:1305-1314.
6. Kosti'c MM, Tagarakis AC, Ljubi'ci'c N, Blagojevi'c D, Radulovi'c M, Ivoševi'c B, Raki'c D. The Effect of N Fertilizer Application Timing on Wheat Yield on Chernozem Soil. *Agronomy*. 2021; 11:1413.
7. El-Habbal MS, Ashmawy F, Saoudi HS, Abbas IK. Effect of nitrogen fertilizer rates on yield, yield components and grain quality measurements of some wheat cultivars using SPAD-Meter. *Egyptian Journal of Agricultural Research*. 2010; 88: 211-223.
8. Raun WR, Solie JB, Johnson GV, Stone ML, Mullen RW, Freeman KW, Thomason WE, Lukina EV. Improving Nitrogen Use efficiency in cereal grain production with optical sensing and variable rate application. *Agronomy Journal*. 2002; 94 (6).
9. Abedi, T.; Alemzadeh, A.; Seyed Abdolreza, K. Wheat Yield and grain protein response to Nitrogen amount and timing. *Australian Journal of Crop Science*. 2011; 5 (3): 330–336
10. Si Z, Zain M, Mehmood F, Wang G, Gao Y. Effects of Nitrogen Application rate and irrigation regime on growth, yield, and water-Nitrogen use efficiency of drip-irrigated winter wheat in the North China plain. *Agricultural water management*. 2020; 231: 106002.
11. Madakadze IC, Stewart KA, Madakadze RM, Peterson PR, Coulman BE, Smith DL. Field evaluation of the chlorophyll meter to predict yield and nitrogen concentration of switchgrass. *Journal of Plant Nutrients*. 1999; 22: 1001-1010.
12. Kantety RV, Van Santen E, Woods FM, Wood CW. Chlorophyll meter predicts nitrogen status of tall fescue. *Journal of Plant Nutrients* 1996; 19: 881-899.
13. Bullock DG, Anderson DS. Evaluation of the Minolta SPAD-502 chlorophyll meter for nitrogen management in corn. *Journal of Plant Nutrients*. 1998; 21: 741-755
14. Jones JB, Jr. *Hydroponics: A Practical Guide for Soilless Grower* 4; St. Lucie Press: Boca Raton, FL, 1997; 230.
15. Didal VK, Vidyasagar GE Ch, Kumar RM, Surekha K, Reddy SN, Brijbhooshan. Effect of nitrogen management practices on SPAD values and NDVI readings of rice crop. *The Pharma Innovation Journal*. 2022; 11(2): 367-371.
16. Gianquinto G, Goffart JP, Olivier M, Guarda G, Colauzzi M, Dalla Costa L., Delle Vedove G, Vos J, MacKerron DKL. The use of hand-held chlorophyll meters as a tool to assess the nitrogen status and to guide nitrogen fertilization of potato crop. *Potato Research*. 2004. 47: 35-80.
17. Guendouz A, Hafsi M, Khebbat Z, Moumeni L, Achiri A. Evaluation of Grain yield, 1000 kernels weight and chlorophyll content as indicators for drought tolerance in durum wheat (*Triticum durum Desf.*). *Journal of Advanced Studies in Agricultural, Biological and Environmental Sciences*. 2014; 1: 89-92.
18. Goncalves da Silva MA, Muniz AS, Mannigel AR, Porto SMA, Marchetti ME, Nolla A, Grannemann I. Monitoring and evaluation of need for nitrogen fertilizer topdressing for maize leaf chlorophyll readings and the relationship with grain yield. *Brazilian Archives of Biology and Technology*. 2011; 54: 665- 674.
19. El-habbal MS, Ashmawy F, Saoudi HS, Abbas IK, 2011 Effect of nitrogen fertilizer rates on yield, yield components and grain quality measurements of some wheat cultivars using SPAD-meter. *Egyptian Journal of Agricultural Research*. 2010; 88 (1).
20. Maiti D, Das DK, Karak T, Banerjee M. Management of nitrogen through the use of leaf color chart (LCC) and soil plant analysis development (SPAD) or chlorophyll meter in rice under irrigated ecosystem. *The Scientific World Journal*. 2004; 4: 838-846.
21. Prost L, Jeuffroy MH. Replacing the nitrogen nutrition index by the chlorophyll meter to assess wheat N status. *Agronomy for Sustainable Development*. 2007; 27: 321-330.

22. Singh B, Singh Y, Ladha JK, Bronson KF, Balasubramanian V, Singh J, Khind CS. Chlorophyll meter- and leaf color chart-based nitrogen management for rice and wheat in northwestern India. *Agronomy Journal*. 2002; 94: 821-829
23. Ramesh K, Chandrasekaran B, Balasubramanian TN, Bangarusamy U, Sivasamy R, Sankaran N. Chlorophyll dynamics in rice (*Oryza sativa*) before and after flowering based on SPAD (chlorophyll) meter monitoring and its relation with grain yield. *Journal of Agronomy and Crop Science*. 2002; 188: 102-105.
24. Reeves DW, Mask PL, Wood CW, Delaney DP. Determination of wheat nitrogen status with a hand-held chlorophyll meter: Influence of management practices. *Journal of plant nutrition*. 1993; 16: 781-796.
25. Rostami M, Koocheki A, Mahallati MN, Kafi M. Evaluation of chlorophyll meter (SPAD) data for prediction of nitrogen status in corn) *Eurasian journal of agricultural & environmental sciences*. 2008; 3: 79-85.
26. Francis DD, Piekielek WP. Assessing Crop Nitrogen Needs with Chlorophyll Meters. The Site-Specific Management Guidelines series-12, published by the Potash & Phosphate Institute (PPI), USA. 2012.
27. Chunjiang Z, Aning J, Wenjiang H, Keli L, Liangyun L, Jihua W. Evaluation of variable-rate nitrogen recommendation of winter wheat based on SPAD chlorophyll meter measurement, *New Zealand Journal of Agricultural Research*. 2007; 50 (5): 735-741.
28. Juško K, Kościk B, *Electronic journal of polish agricultural universities*, possible use of the chlorophyll meter (SPAD- 502) for evaluating nitrogen nutrition of the Virginia tobacco. *Alina*. 2002;5 (1).
29. Setiawati MR, Aini HF, Suryatmana P, Hindersah R. 2019. Application of inorganic fertilizer and bio-fertilizer on chlorophyll content, pH, and leaves number of pakchoi (*Brassica rapa*L.) in hydroponics. *International Journal of Agriculture, Environment and Bioresearch*. 2019; 4(04): 269-278.
30. Chapman SC, Barreto HJ. Using a chlorophyll meter to estimate specific leaf nitrogen of tropical maize during vegetative growth. *Agronomy Journal*. 1995; 89: 557-562.
31. Blackmer TM, J.S. Schepers. Use of a chlorophyll meter to monitor nitrogen status and schedule fertigation for corn. *Journal of Production Agriculture*. 1995; 8:56–60.
32. Evans JR. Nitrogen and photosynthesis in the flag leaf of wheat (*Triticum aestivum* L.). *Plant Physiology*. 1983; 72: 297-302.
33. Amaliotis D, Therios I, Karatissiou M. Effect of nitrogen fertilization on growth, leaf nutrient concentration and photosynthesis in three peach cultivars. *ISHS, Acta Horticulturae*. 2004; 449: 36-42.
34. Tucker M. Primary Nutrients and Plant Growth. - In: *Essential Plant Nutrients* (SCRIBD, Ed.). North Carolina Department of Agriculturae, 2004.
35. Daughtry CST, Walthall CI, Kim MS, Browndecostoun E, McMurtrey JE. Estimating corn leaf chlorophyll concentration from leaf and canopy reflectance. *Remote Sensing of Environment*. 2000; 74:229-239.
36. Islam MR, Shamsul Haque KM, Akter N, Karim MA. Leaf chlorophyll dynamics in wheat based on SPAD meter reading and its relationship with grain yield, *Scientia Agriculturae*. 2014; 8 (1): 13-18
37. Shukla AK, Ladha JK, Singh VK, Dwivedi BS, Balasubramanian V, Gupta RK, Sharma SK, Singh Y, Pathak H, Pandey PS, Padre AT, Yadav RL. Calibrating the leaf color chart for nitrogen management in different genotypes of rice and wheat in a systems perspective. *Agronomy Journal*. 2004; 96:1606-1621.
38. Khan MA, Shukla AK, Upadhyay NC, Singh OP. Nutrient management strategies for maize-potato-wheat cropping system in typic ustochrept soils. *Potato Journal*. 2005; 32: 167-68.
39. Roba T. Review on: The Effect of Mixing Organic and Inorganic Fertilizer on Productivity and Soil Fertility. *Open Access Library Journal*. 2018; 5 (6) :1-11.

40. Amujoyegbe BJ, Opabode JT, Olayinka A. Effect of organic and inorganic fertilizer on yield and chlorophyll content of maize (*Zea mays* L.) and sorghum *Sorghumbicolour* (L.) Moench. African Journal of Biotechnology. 2007; 6(16): 1869-1873.
41. Recous S, Robin D, Darwis D, Mary B. Soil inorganic N availability: effect on maize residue decomposition. Soil Biology and Biochemistry. 1995; 12:1529-1538.
42. Yadav S, Lal M, Naresh RK, Yadav RB, Yadav AK, Yadav KG, Kumar R, Sharath Chandra M, Pradeep Rajput. Effect of Organic and Inorganic Nutrient Sources on Productivity, Grain Quality of Rice and Soil Health in North-West IGP: A Review. International Journal of Current Microbiology and Applied Sciences. 2019; 8 (12): 2488-2514.
43. Khachi B, Sharma SD, Ghumare V, Kumar P, Mir M. Study on comparative efficacy of bio-organic nutrients on plant growth, leaf nutrient contents and fruit quality attributes of kiwi fruit. Journal of Applied and Natural Science. 2015; 7(1): 175 – 181.
44. Monje OS, Bugbee B. The inherent limitations of non-destructive chlorophyll meters: a comparison of two types of meters. HortScienc. 1992; 27 (1): 69-71.
45. Vos J, Bom M, Hand-held chlorophyll meter: a promising tool to assess the nitrogen status of potato foliage. Potato Research. 1993; 36: 301-308.
46. Kitagawa Y, Okayama K, Hirokawa T. Determination of leaf colour in rice cv. Koshihikari plants with a chlorophyll meter. Bull. Toyama Agriculture Research Centre 1987; 1: 1-7.
47. Piekielek WP, Fox RH, Use of chlorophyll meter to predict sidedress nitrogen requirements for maize. Agronomy Journal. 1992; 84: 59-65.
48. Smeal D, Zhang H. Chlorophyll meter evaluation for nitrogen management in corn. Soil Science Plant Analysis. 1994; 25 (9/10), 1495-1503.
49. Waskom RM, Westfall DG, Spellman DE, Soltanpour PN. Monitoring nitrogen status of corn with a portable chlorophyll meter. Soil Science Plant Analysis. 1996; 27 (384): 561-574.
50. Hussain F, Bronson KF, Singh Y, Singh B, Peng S. Use of chlorophyll meter sufficiency indices for nitrogen management of irrigated rice in Asia. Agron. J. 2000; 92:875-879.
51. Varvel GE, Wilhelm WW, Shanahan JF, Schepers JS. An algorithm for corn nitrogen recommendations using a chlorophyll meter-based sufficiency index. Agron. J. 2007; 99:701-706.

**Table 1. Correlation and Coefficient of variance of SPAD with total chlorophyll and total N content**

Stages	Correlation of SPAD with total chlorophyll and total N content		Coefficient of variance of SPAD with total chlorophyll and total N content	
	Total chlorophyll	Total N content	Total chlorophyll	Total N content
<b>CRI</b>	0.94	0.96	0.84	0.75
<b>Tillering</b>	0.90	0.94	0.76	0.76
<b>Jointing</b>	0.92	0.83	0.78	0.54
<b>Milking</b>	0.94	0.94	0.71	0.85

**Table 2. Effect of different sources and NPK levels on SPAD readings at different growth stages**

M/S	SPAD															
	CRI				Tillering				Jointing				Milking			
	M1	M2	M3	Mean	M1	M2	M3	Mean	M1	M2	M3	Mean	M1	M2	M3	Mean
<b>S1</b>	4.20	3.83	4.17	<b>4.06</b>	2.70	6.60	4.10	<b>4.47</b>	13.95	17.38	15.82	<b>15.72</b>	17.17	13.47	17.86	<b>16.17</b>
<b>S2</b>	8.40	5.03	5.76	<b>6.40</b>	12.15	7.53	10.58	<b>10.09</b>	28.31	18.14	26.48	<b>24.31</b>	31.86	16.13	25.42	<b>24.47</b>
<b>S3</b>	9.23	5.10	7.52	<b>7.29</b>	15.28	7.15	10.90	<b>11.11</b>	33.70	18.49	27.63	<b>26.60</b>	32.44	20.27	29.01	<b>27.24</b>
<b>S4</b>	10.50	5.47	7.93	<b>7.97</b>	15.64	6.11	11.44	<b>11.07</b>	33.48	19.85	29.01	<b>27.45</b>	38.92	16.48	28.65	<b>28.02</b>
<b>S5</b>	8.40	2.90	6.40	<b>5.90</b>	16.11	6.34	11.53	<b>11.33</b>	33.45	12.43	28.29	<b>24.72</b>	32.86	16.90	24.04	<b>24.60</b>
<b>Mean</b>	<b>8.15</b>	<b>4.47</b>	<b>6.36</b>	<b>6.32</b>	<b>12.38</b>	<b>6.74</b>	<b>9.71</b>	<b>9.61</b>	<b>28.58</b>	<b>17.26</b>	<b>25.45</b>	<b>23.76</b>	<b>30.65</b>	<b>16.65</b>	<b>25.00</b>	<b>24.10</b>
<b>SEm±</b>		0.19				0.48				0.90				1.18		
<b>CD(p=0.05)</b>		<b>0.76</b>				<b>1.90</b>				<b>3.56</b>				<b>4.64</b>		
<b>SEm±</b>		0.20				0.45				0.93				1.14		
<b>CD(p=0.05)</b>		<b>0.60</b>				<b>1.31</b>				<b>2.71</b>				<b>3.34</b>		
<b>Int I</b>		0.35				0.77				1.61				1.98		
<b>CD(p=0.05)</b>		<b>1.03</b>				<b>2.27</b>				<b>4.70</b>				<b>5.79</b>		
<b>Int II</b>		0.50				1.19				2.31				2.95		
<b>CD(p=0.05)</b>		<b>1.46</b>				<b>3.48</b>				<b>6.76</b>				<b>8.62</b>		

*M1- Inorganic sources (NPK fertilizers), M2- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and M3- Integrated sources (50% Inorganic + 50% organic) S1- control, S2- 100% RDF, S3- 150% RDF, S4- 200% RDF, S5- Based on soil test value for target yield of 6t ha<sup>-1</sup>*

**Table 3. Effect of different sources and NPK levels on total chlorophyll at different growth stages**

M/S	Total chlorophyll															
	CRI				Tillering				Jointing				Milking			
	M1	M2	M3	Mean	M1	M2	M3	Mean	M1	M2	M3	Mean	M1	M2	M3	Mean
<b>S1</b>	0.74	0.72	0.73	<b>0.73</b>	1.35	1.26	1.26	<b>1.29</b>	1.48	1.19	1.30	<b>1.32</b>	0.92	0.73	0.89	<b>0.85</b>
<b>S2</b>	1.20	0.98	1.06	<b>1.08</b>	2.02	1.36	1.88	<b>1.75</b>	2.14	1.45	1.75	<b>1.78</b>	1.26	0.75	1.14	<b>1.05</b>
<b>S3</b>	1.61	1.05	1.38	<b>1.35</b>	2.16	1.40	2.01	<b>1.86</b>	2.36	1.51	2.01	<b>1.96</b>	1.38	0.90	1.29	<b>1.19</b>
<b>S4</b>	1.54	1.00	1.22	<b>1.26</b>	2.27	1.34	1.92	<b>1.84</b>	2.58	1.72	2.34	<b>2.21</b>	1.54	1.13	0.94	<b>1.20</b>
<b>S5</b>	1.25	0.88	0.98	<b>1.04</b>	1.82	1.27	1.66	<b>1.59</b>	2.16	1.19	1.51	<b>1.62</b>	1.12	0.81	0.92	<b>0.95</b>
<b>Mean</b>	<b>1.27</b>	<b>0.93</b>	<b>1.08</b>	<b>1.09</b>	<b>1.92</b>	<b>1.33</b>	<b>1.75</b>	<b>1.67</b>	<b>2.14</b>	<b>1.41</b>	<b>1.78</b>	<b>1.78</b>	<b>1.24</b>	<b>0.87</b>	<b>1.04</b>	<b>1.05</b>
<b>SEm±</b>		<b>0.029</b>				<b>0.043</b>				<b>0.082</b>				<b>0.028</b>		
<b>CD(p=0.05)</b>		<b>0.113</b>				<b>0.169</b>				<b>0.321</b>				<b>0.109</b>		
<b>SEm±</b>		0.031				0.044				0.044				0.033		
<b>CD(p=0.05)</b>		<b>0.090</b>				<b>0.130</b>				<b>0.129</b>				<b>0.097</b>		
<b>Int I</b>		0.053				0.077				0.076				0.057		
<b>CD(p=0.05)</b>		<b>0.156</b>				<b>0.224</b>				<b>0.223</b>				<b>0.167</b>		
<b>Int II</b>		0.075				0.110				0.177				0.076		
<b>CD(p=0.05)</b>		<b>0.218</b>				<b>0.321</b>				<b>0.517</b>				<b>0.221</b>		

*M1- Inorganic sources (NPK fertilizers), M2- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and M3- Integrated sources (50% Inorganic + 50% organic) S1- control, S2- 100% RDF, S3- 150% RDF, S4- 200% RDF, S5- Based on soil test value for target yield of 6t ha<sup>-1</sup>*

**Table 4. Effect of different sources and NPK levels on total N content (%) in leaves at different growth stages**

M/S	Total N content (%)															
	CRI				Tillering				Jointing				Milking			
	M1	M2	M3	Mean	M1	M2	M3	Mean	M1	M2	M3	Mean	M1	M2	M3	Mean
<b>S1</b>	2.13	2.13	2.15	<b>2.14</b>	2.09	2.04	1.97	<b>2.03</b>	1.17	1.17	1.23	<b>1.19</b>	1.07	1.10	1.14	<b>1.10</b>
<b>S2</b>	2.33	2.18	2.30	<b>2.27</b>	2.26	2.12	2.18	<b>2.19</b>	1.82	1.32	1.66	<b>1.60</b>	1.36	1.22	1.28	<b>1.29</b>
<b>S3</b>	2.52	2.24	2.35	<b>2.37</b>	2.28	2.15	2.19	<b>2.21</b>	2.04	1.95	2.00	<b>1.99</b>	1.45	1.24	1.41	<b>1.37</b>
<b>S4</b>	2.56	2.31	2.33	<b>2.40</b>	2.41	2.20	2.23	<b>2.28</b>	2.13	2.02	2.11	<b>2.09</b>	1.56	1.23	1.50	<b>1.43</b>
<b>S5</b>	2.40	2.25	2.19	<b>2.28</b>	2.38	2.15	2.20	<b>2.24</b>	1.83	1.52	1.83	<b>1.72</b>	1.48	1.17	1.36	<b>1.34</b>
<b>Mean</b>	<b>2.39</b>	<b>2.22</b>	<b>2.26</b>	<b>2.29</b>	<b>2.29</b>	<b>2.13</b>	<b>2.15</b>	<b>2.19</b>	<b>1.80</b>	<b>1.60</b>	<b>1.76</b>	<b>1.72</b>	<b>1.38</b>	<b>1.19</b>	<b>1.34</b>	<b>1.30</b>
<b>SEm±</b>		0.032				0.029				0.034				0.024		
<b>CD(p=0.05)</b>		<b>0.126</b>				<b>0.113</b>				<b>0.136</b>				<b>0.096</b>		
<b>SEm±</b>		0.027				0.032				0.038				0.025		
<b>CD(p=0.05)</b>		<b>0.080</b>				<b>0.093</b>				<b>0.111</b>				<b>0.072</b>		
<b>Int I</b>		0.047				0.055				0.066				0.043		
<b>CD(p=0.05)</b>		<b>NS</b>				<b>NS</b>				<b>0.192</b>				<b>0.126</b>		
<b>Int II</b>		0.077				0.076				0.091				0.062		
<b>CD(p=0.05)</b>		<b>2.13</b>				<b>2.13</b>				<b>2.15</b>				<b>2.14</b>		

*M1- Inorganic sources (NPK fertilizers), M2- organic sources (FYM, vermicompost, biofertilizers Azotobacter and PSB) and M3- Integrated sources (50% Inorganic + 50% organic) S1- control, S2- 100% RDF, S3- 150% RDF, S4- 200% RDF, S5- Based on soil test value for target yield of 6t ha<sup>-1</sup>*




Fig 1a. Relationship between SPAD and chlorophyll content at different growth stages




Fig 1b. Relationship between SPAD and nitrogen content at different growth stages

Fig 1. Relationship of SPAD with chlorophyll and N content at different growth stages