

Original Research Article

Biodiversity and Distribution Patterns of Clams (*Tridacnidae*) on Coral Reefs In The Waters of Indonesia Kayu Angin Bira Island, Seribu Islands

ABSTRACT

This research aims to find out the biodiversity and distribution patterns of clams and the relationship between clam abundance and coral reef conditions in the Waters of Kayu Angin Bira Island, Kepulauan Seribu. The research was conducted between April and July 2022, with field data collection taking place in September 2022 at four chosen stations. These stations were selected based on variations in water depth and were situated in the South, West, North, and East regions of the waters surrounding Kayu Angin Bira Island. The data collection was conducted using survey methods. Coral reef data collection using LIT method and clam observation using belt transect. The results showed that in the waters of Bira Kayu Angin Island, 25 individuals of 2 species of bivalve class were found, namely *Tridacna squamosa*, *Tridacna crocea*. The highest relative abundance of clams is found at the eastern station which is 20% and the lowest relative abundance is found at the southern, western and northern stations which is only 8%. The diversity index of clams at the four stations was obtained in the low category, which ranged from 0.451 - 0.662 and the uniformity index at the four stations ranged from 0.650 - 0.954, which means high population uniformity. The spatial distribution pattern of clams at the four stations is uniform which ranges from 0.4 - 0.74. The relationship between clams and coral reefs is 0.7284, which means that clams have a strong relationship with the condition of coral reefs.

Keywords: Biodiversity, Distribution Patterns, *Tridacnidae*, Coral Reefs, Kayu Angin Bira Island, Seribu Islands

1. INTRODUCTION

The presence of benthic fauna in a body of water is important for the aquatic ecosystem. In addition to being an indicator of the health of the aquatic environment (Bilyard, 1987), benthic fauna also has an important role in supporting capture fisheries activities (Kritzer et al., 2016) and as symbiotic fauna for demersal fauna (Kaiser et al., 1999). One of the benthic fauna is Clams (*Tridacninae* ~~*Tridacnidae*~~ sp.).

Clams play an important role in coral reef ecosystems to maintain ecosystem balance (Neo et al., 2013). The depth-related distribution of clams (*Tridacninae* ~~*Tridacnidae*~~) has an impact on their shape and size. In regions with greater water depth, the clams tend to be smaller compared to those residing in shallower waters (Niar tiningsih et al., 2008). The abundance of clams is influenced by the condition or quality of coral reefs as their habitat (Marsukiet al., 2012).

There are several studies that have been conducted on the abundance of clams in Indonesia, one study conducted by Othman, Goh, & Todd, (2010) revealed that habitat loss, pollution, and increased sea surface temperatures are the causes of declining clam populations. Van Wynsberge et al. (2016) found that human population is closely related to clam abundance. According to Setiyaningsih (2020), clams in Pulau Seribu are under threat of population decline due to the culture and habits of the surrounding

community who consume clams. In addition to human factors, natural factors also affect the abundance of clams. This study aims to measure biodiversity and observe the distribution pattern of clams in the Waters of KayuAnginBiral Island, KepulauanSeribu and analyze the relationship between clam abundance and coral reef conditions in the Waters of KayuAnginBira Island, KepulauanSeribu.

2. METHODS

Research Time and Location




Figure 1. Map of The Research Area

This research was conducted in September 2022. Then, the data processing was carried out until November 2022. The research was conducted in the waters of KayuAnginBira Island, Taman NasionalLautKepulauanSeribu, DKI Jakarta. Data collection took place in the waters of KayuAnginBira Island at four specific station locations: South, West, North, and East. The research stations were chosen using purposive sampling, a technique that involves specific considerations (Sugiyono, 2012). In this study, four stations were selected, each plot at varying depths: 3 meters and 9 meters.

The method used in this research is the Survey Method. A survey method in research involves studying a population as the subject, but it focuses on analyzing data collected from samples taken within that population. This approach helps uncover distributions and relationships between variables (Sugiyono, 2013). The data studied in this research includes clam biodiversity data, coral reef data and water quality data. The data gathered is both quantitative and qualitative information from each observation station. This information includes metrics such as the percentage of coral reef cover, diversity, uniformity,

abundance, and patterns of clam distribution, and the data is analysed to explore the correlation between clam abundance and the condition of the coral reefs.

Comment [HP1]: Kindly mentioned the coordinates of studyzone . Also present in a brief paragraph the three study zone.

Observed Parameters

- **Percentage of Coral Reef Cover**

Data obtained from the field is the data from transects with the LIT method, which is then stored in a computer and processed with the Ms. Excel program. To calculate coral reef cover, the formula was used (English *et al.* 1994).

$$L = \frac{Li}{N} \times 100\%$$

Description:

L = Percentage of Coral cover
Li = Amount of each component
N = Total Components (50m)

The percentage of live coral cover is used to determine the coral reef condition. Coral cover categories based on Gomes and Yap (1984) are shown in (Table 1).

Table 1. Coral reef assessment criteria

Category	Live Coral Cover (%)
Bad	0 - 24,9
Moderate	25 - 49,9
Good	50 - 74,9
Very good	75 - 100

- **Clam abundance**

Abundance is the total number of individual clams found during observation. The abundance index provides an overview of the species composition in the community. The relative abundance of clams can be found using the following equation (Van Balen, 1984):

$$KR = \frac{ni}{N} \times 100\%$$

Description:

KR = Relative Abundance
ni = Number of individuals of species-i
N = Total number of individuals

- **Clam diversity**

The diversity index shows the richness of species in a community and shows the balance in the distribution of numbers per species. The diversity index can be calculated with the Shannon-Wiener index (Magurran, 1988) with the equation:

$$H' = -\sum (Pi) (\ln Pi)$$

Description:

H' = Diversity Index
Pi = ni/N
ni = Number of Individuals

N = Total number of individuals

Criteria:

H' < 1 : Low Diversity
1 < H' < 3 : Medium Diversity
H' > 3 : High Diversity

- **Clam uniformity**

Clam diversity can be seen from the uniformity index. This index is used to determine the presence of species that dominate the community and to determine the distribution of the number of individuals of each type (Odum 1971). This index is obtained by comparing the diversity index with its maximum value as follows:

$$E = \frac{\hat{H}}{H_{max}}$$

Description:

E = Uniformity index
 \hat{H} = Shannon-Wiener diversity index
Hmax = Ln S
S = Number of species

Criteria:

E < 0.4 : Small population uniformity
0.4 < E < 0.6 : Moderate population uniformity
E > 0.6 : High population uniformity

- **Clam distribution pattern**

To determine the distribution pattern of clams in the habitat, the Morisita distribution pattern was used. Morisita distribution index is calculated by the following formula (Brower and Zar, 1990):

$$Id = q \frac{\sum ni(ni-1)}{N(N-1)}$$

Description:

Id : Morisita Scatter Index
q : number of samples

ni : number of individuals of i sample.
N : Total number of individuals obtained

With criteria:

Id < 1 : The distribution pattern of individual species is uniform
Id = 1 : The distribution pattern of individual species is random
Id > 1 : The distribution pattern of individual species is clustered

Correlation Data Analysis

Data analysis in this research was carried out using Bivariate Correlation Analysis processed in Microsoft Excel. Bivariate Correlation Analysis is used to measure the level of relationship between two variables.

The formula used is as follows:

$$r_{xy} = \frac{\sum XY - \sum X \cdot \sum Y}{\sqrt{[n \cdot \sum X^2 - (\sum X)^2][n \cdot \sum Y^2 - (\sum Y)^2]}}$$

Description:

- r_{xy} = Correlation coefficient of two variables (Clam Abundance and Percentage Coral Reef Cover)
n = Number of transects
X = Number of Clams at each station ($x + x_{12} + \dots$)
Y = Coral reef area at each station ($y + y_{12} + \dots$)

The results of the bivariate correlation analysis guidelines according to Sugiyono (2007) in Miala (2015) can be seen in (Table 2):

Table 2. Criteria for correlation value

Interpretation Number	Correlation
0 - 0,199	Very weak
0,20 - 0,399	Weak
0,40 - 0,599	Moderate
0,60 - 0,799	Strong
0,80 - 1,0	Very strong

3. RESULTS AND DISCUSSION

The measurement results of water quality parameters of KayuAnginBira Island

Table 3. Water quality parameters

Station	Temperature (° C)	Salinity (ppt)	pH	Brightness (%)
South	30	35	8	100%
West	30.1	35	8	100%
North	30.2	35	8	100%
East	30	34	8	100%

Based on the observations (Table 3) conducted at four different stations, the data collected were regarding water conditions, coral reef density, and reef cover. The parameters observed included temperature, salinity, brightness, and pH levels. These physical and chemical water parameters were measured with three repetitions conducted simultaneously at each station.

The water temperature at KayuAnginBira Island in (Table 3) at each station has an average of 30.1°C. This temperature range is still in the normal range and is tolerable for aquatic biota. According to Rahman (2006), marine biota can tolerate temperatures ranging from 20-35°C. Salinity measurements at the research site show that the salinity levels in the waters of KayuAnginBira Island ranged from 34-35 ppt. The salinity levels at all four stations are classified as suitable for the growth of clams, such as the findings of Jameson (1976), which states that an optimal salinity range for clams is typically between 25-40 ppt.

The results of brightness level measurements in the waters of KayuAnginBira Island are 100%. Brightness is a critical factor that significantly impacts the survival of clams due to their association with clamsymbionts, zooxanthella, which rely on sunlight for the photosynthesis process. This is further clarified by Romimohtarto et al. (1987), as cited in Niartiningsih (2012), regarding the importance of adequate light for zooxanthella's photosynthesis, making clear and shallow waters essential for clams.

Comment [HP2]: Review the position of figures and tables in the text in accordance with the General Guideline for Authors

Comment [HP3]: Why the methodology of these parameters was not presented in subsection2 ?

Furthermore, based on KEPMEN LH No. 51 tahun 2004 concerning marine biota, the brightness levels at KayuAnginBira Island satisfy the criteria for supporting a healthy biota population. Lastly, considering the pH requirements for bivalves, Hart and Fuller (1974) state that bivalves thrive within a pH range of 5.8 to 8. Thus, water conditions at KayuAnginBira Island appear suitable as a habitat for clams.

Clam Abundance

Referring to KEPMEN LH No. 51 of 2004, the results of observations of the physical and chemical properties of seawater in the waters around KayuAnginBira Island indicate favorable conditions for the survival and growth of clams.




Figure 2. Relative abundance of clams in the waters of KayuAnginBira Island

The figure above illustrates the relative abundance of clams at four different stations, categorized by variations in depth. The highest total relative abundance was observed at the eastern station, specifically at a depth of 3 meters is 20%. The lowest relative abundance was recorded at the southern, western, and northern stations, with only 8%.

In general, the clams found in the waters around KayuAnginBira Island are typically found on both living and dead massive coral substrates, as well as rocky substrates. Among the four stations, clams were more common at a depth of 3 meters compared to a depth of 9 meters. This phenomenon happens because the sunlight that reaches the seafloor at a depth of 3 meters is more ample, while sunlight diminishes significantly at a depth of 9 meters. This observation aligns with the statement of Rosewater (1965), that sunlight plays a crucial role in the photosynthesis of zooxanthellae, which is highly beneficial for clams.

At all four observation stations categorized by depth differences, two types of clams, *T. Crocea* and *T. Squamosa*, were identified. The higher abundance of *T. Crocea* can be attributed to its unique habitat within coral reefs, which makes it challenging for people to harvest this species. *T. Crocea* is typically found attached to hard substrates, fully immersed in the substrate, particularly on massive corals and rocks. This observation aligns with Panggabean (1991) description of *T. Crocea* as a small clam that thrives on hard substrates for complete shell immersion. Conversely, *T. Squamosa* was found in fewer observation locations, with only seven individuals recorded.

T. crocea was the most abundant species found at all stations. *T. crocea* species are more abundant than *T. squamosa* species due to the lack of utilization by the community of *T. crocea* species and the tendency of *T. crocea* species to live immersed in the substrate making it difficult to collect.

T. Crocea emerged as the most abundant species across all stations. Its higher abundance can be attributed to the limited use of *T. Crocea* by the local community and the species' tendency to inhabit substrates that are challenging to access.

On the other hand, *T. Squamosa* is the least commonly found clam species in the waters surrounding KayuAnginBira Island. This observation aligns with Nugroho and Ambaryanto's (2001) statement, which suggests that clam populations in nature are predominantly composed of smaller species like *T. Crocea* and *T. Maxima*, while larger species such as *T. Derasa*, *T. Squamosa*, *H. Hippopus*, and *H. Porcellanus* are quite rare.

Clam Diversity and Uniformity

Based on the research results, the diversity and uniformity indices of clams at the four stations obtained varying results. The diversity and uniformity of clams in the waters of KayuAnginBira Island can be seen in Figure 3.

Based on the data analysis, it's evident that the diversity index in the waters surrounding KayuAnginBira




Figure 3. Diversity and uniformity of clams in KayuAnginBira Island

Island falls within the low category ($H' < 1$), while the uniformity index is categorized as high ($E > 0.6$).

The diversity index across the four observation stations ranged from 0.451 to 0.662, which means that the diversity level is relatively low. This indicates that while productivity may be low, the ecosystem maintains a balanced state with moderate ecological pressure. The diversity index serves as a systematic representation describing the community's structure and facilitates the analysis of information regarding the types and quantities of organisms present. It's worth noting that the greater the number of species discovered, the higher the diversity index, though this value is influenced by the number of individuals of each species, as pointed out by Wilhm and Doris (1986). Krebs (1985) also supports this perspective by indicating that a larger number of evenly distributed individuals will result in a higher diversity index.

While the uniformity index at the four observation stations ranged from 0.650 - 0.954 which means high population uniformity. According to Lind (1979) in Amin (2008) the uniformity index is close to one then the community is stable and the number of individuals between species is the same. In addition, the diversity and uniformity of biota in a body of water is highly dependent on the number of species in the community.

The uniformity index at the four observation stations ranged from 0.650 to 0.954, indicating a high level of population uniformity. As per Lind (1979), as cited in Amin (2008), when the uniformity index approaches one, it signifies a stable community with a balanced distribution of individuals among species. Furthermore, it's important to note that the diversity and uniformity of biota in a body of water are strongly influenced by the number of species present within the community.

The results of research conducted in the waters of KayuAnginBira Island found 2 types of clams, namely *T. Squamosa* and *T. crocea*. *T. squamosa* found in the waters of KayuAnginBira Island has a blackish brown coat with a mixture of white spots. This clam was found at all four stations and totaled 7 individuals. *T. crocea* found a striking and beautiful coat color. *T. crocea* was found in 18 individuals and lived on *coral massive and rock* substrates. This clam has the most abundant population. This is evidenced by the discovery of this type of clam at all four observation stations, namely east, south, west and north.

The research conducted in the waters of KayuAnginBira Island identified two types of clams, specifically *T. Squamosa* and *T. Crocea*. *T. Squamosa*, characterized by a blackish-brown shell with white spots, was found at all four stations, totaling seven individuals. On the other hand, *T. Crocea* exhibited a striking and beautiful shell coloration. This species was encountered in 18 individuals and primarily inhabited massive coral and rock substrates. *T. Crocea* demonstrated the most abundant population, as evidenced by its presence at all four observation stations, namely east, south, west, and north.

Clam Distribution Pattern

To determine the spatial distribution pattern of clams in the waters of KayuAnginBira Island, the Morisita index was calculated. Based on the results of calculations using the Morisita Index shows the distribution pattern of clams in KayuAnginBira Island is uniform. Clam distribution pattern data can be seen in Table 4.

Table 4. Kima Scatter Value

Station	Morisita Index	Distribution Pattern
South	0.4	Uniform
West	0.26	Uniform

North	0.4	Uniform
East	0.74	Uniform

Based on the results presented in Table 4, the Morisita Index value varies across the stations, with the highest value observed at the East station (0.74) and the lowest values recorded at the South and North stations (0.4). The calculations of the Morisita Index at each station consistently indicate a uniform distribution pattern of clams. In all four stations, the Morisita Index is less than one, indicating a uniform pattern of clam distribution within these locations. This uniformity can be attributed to the availability of suitable habitat conditions that allow each species to find its own niche and food source. This observation aligns with Odum's (1971) assertion that uniform distribution patterns emerge when there is a balanced level of competition among individuals or when there exists a positive antagonistic relationship that supports spatial distribution.

The uniform distribution of clams across the four stations is likely influenced by the substrates they inhabit and the availability of sunlight. As Suin (2002) suggests, the physical and chemical factors that are evenly distributed within a habitat, along with the presence of food sources, play a crucial role in determining whether animals live in groups, random patterns, or uniform distributions.

According to Krebs (2003), distribution patterns are a result of various factors, including genetic traits and habitat preferences of individual species. Additionally, other factors such as the spatial and temporal distribution of food resources and competition for habitat resources due to adverse environmental conditions also contribute to the observed distribution patterns.

Relationship between Clams and Coral Reefs

The correlation analysis reveals a strong and positive relationship between clam abundance and the percentage of coral reef cover in the waters surrounding KayuAnginBira Island, with a substantial correlation coefficient of 0.853 (as indicated in Table 2). Notably, the coefficient of determination (R square) obtained from our calculations is 0.7284, equivalent to 72.84%. This implies that the presence of clams is significantly influenced by the coral reef cover, accounting for 72.84% of the variation, while the remaining 27.16% is attributed to other contributing factors. Figure 4 provides a visual representation of this correlation between coral reef cover and clam abundance.




Figure 4. Correlation of percent coral cover and clam abundance

According to Astuti (2017), the correlation coefficient (r) is positive, it signifies a direct and proportional relationship between variables X and Y. In simpler terms, as variable X increases, the value of variable Y also increases, as illustrated in Figure 4. The results of our correlation analysis affirm a very strong relationship, which aligns with Neo's (2013) statement that clams play a pivotal role in maintaining the balance of the coral reef ecosystem, acting as keystone species for coral reef communities. Consequently, clams share a directly proportional relationship with coral reefs.

The decline in clam populations can be attributed to habitat destruction caused by clam poaching techniques that damage coral reefs. Some species of clams live attached to the coral substrate and stuck between the living corals, there are even species of clams that immerse themselves in the coral. To collect clams that live attached to the coral, it is usually done by damaging the coral using crowbars or iron picks so that the coral will be broken and damaged. If this continues, the damage to the coral will become worse and the clam population will automatically decrease.

The decline in clam populations can be caused by habitat destruction resulting from clam poaching techniques that harm coral reefs. Certain clam species attach themselves to coral substrates, nestled among living corals, and some even immerse themselves within the coral structure. The collection of clams attached to coral typically involves damaging the coral using tools like crowbars or iron picks, leading to coral breakage and damage. If this practice continues, it will worsen coral damage and inevitably lead to a reduction in clam populations.

4. CONCLUSION

Based on the results of research on the abundance and diversity as well as spatial distribution patterns of clams in the coral reef waters of KayuAnginBira Island, KepulauanSeribu, the following conclusions were obtained:

- In the waters surrounding KayuAnginBira Island, only two types of clams were discovered: *T. Squamosa* and *T. Crocea*, with a total count of 25 individuals. Across the four stations, categorized by differences in depth, the relative abundance of clams ranged from 8% to 20%. The diversity index falls within the low category, while the uniformity index registered in the high category. The spatial distribution pattern of clams in the waters surrounding KayuAnginBira Island appears to be uniform.
- Based on the correlation of coral reef cover percentage and clam abundance, it a positive relationship, implying that changes in clam abundance, whether high or low, directly correspond to variations in the percentage of coral reef cover.

REFERENCES

1. Amin M, Utojo. 2008. Composition and Diversity of Plankton Species in the Waters of Kupang Bay, East Nusa Tenggara Province. *Toran* Vol. 18 (2): 129 - 135
2. Bilyard, G. R. (1987). The value of benthic infauna in marine pollution monitoring studies. *Marine Pollution Bulletin*, 18(11), 581-585.
3. Brower, JE, Zar JH. (1990). *Field and Laboratory Methods for General Ecology*. Third Edition. Dubuque, Iowa: C. Brown Publishers.
4. English, S., C. Wilkinson and V. Baker. 1994. *Survey manual for tropical marine resources*. - Australian Marine Science Project Living Coastal Resources. Australia, 390 pp.

5. Gomez E.D. and H. Yap. 1984. Monitoring Reef Condition. In Kenchington, R.A. and B. Hudson E.T. Coral Reef Management Hand Book. Unesco Regional Office for Science and Technology for South East Asia. Jakarta, 187-195 pp.
6. Hart, C. W. and Fuller, S. L. 1974. *Pollution Ecology of Estuarine Invertebrates*. London: Academic Press.
7. Jameson, C. S., 1976. Early Life History of Clams *Tridacnacrocea* Lamarck, *Tridacna maxima* (Roding) and *Hipopushiopus*. *Pacific Science*. 30 (3) : 219- 233
8. Krebs, C.J. 1985. *Ecology: The Experimental Analysis of Distributions and Abundance*. 3rd ed. New York.
9. Magurran A.E. 1988. *Ecological Diversity and Its Measurement*. New Jersey (US): Princeton University Press.
10. Marsuki, I.D., Baru, S., Ratna, D.P. 2012. Coral Reef Condition and Clam Abundance in Indo Island Waters. Haluoleo University. *Journal of Indonesian Marine Mina*. Kendari.
11. Neo, M.L. and P.A. Todd. 2013. Conservation Status Reassessment of Clams (*Mollusca: Bivalvia: Tridacninae*) In Singapore Nature In Singapore. National University of Singapore. 6 : 125-13
12. Niartingsih, A., Maghdalena, L., Emma, S., Idris, P. 2008. Juvenile rearing of hawksbill clams (*Tridacnasquamosa*) and lola (*Trochusniloticus*) in monoculture and polyculture at different depths in the waters of Badi Island, Pangkep Regency. COREMAP II Coral Reef Rehabilitation and Management Program. South Jakarta.
13. Nugroho, E.P. and Ambariyanto. (2001). Influence of bottom substrate on the growth of clams (*Tridacnacrocea*). *Marine Science* 23: 237 - 243.
14. Odum, E.P., 1971, *Fundamentals of Ecology*. W.B. Saunders Company, Philadelphia
15. Othman, A. S. Bin, Goh, G. H. S., & Todd, P. A. (2010). The distribution and status of clams (family Tridacnidae) - A short review. *Raffles Bulletin of Zoology*, 58(1), 103-111.
16. Panggabean LMG. 1991. *The Secret Life of Clams: III. Survival*. *Oseana*. 16(2): 35-45.
17. Rahman, A. (2006) Heavy metal content of lead (Pb) and cadmium (Cd) in some crustacean species at Batakan and Takisung Beach, Tanah Laut Regency, South Kalimantan. *Bioscientiae*, 3, pp. 93-101.
18. Romimohtarto, K., Pardomuan, S., M.G. Lily Panggabean, Sutomo. 1987. KIMA Biologi, Resources and Sustainability. Oceanology Research and Development Center-LIPI. Jakarta.
19. Rosewater, J. 1965. The family *Tridacnidae* in the Indo-Pacific. *Indo-Pacific Mollusca*. 1:347-396.
20. Setyaningsih, M., Amali, H and Susilo. 2020. Density Study of Clamshells (Tridacnidae) in Pramuka Island and Dutch Island, Thousand Islands, DKI Jakarta. *Quagga: Journal of Education and Biology* 12(2): 188-193.
21. Suin, N.M. 2002. *Ecological Methods*. Publisher of Andalas University. Padang.
22. Sugiyono, 2013. *Quantitative, Qualitative and R&D Research Methods*. Bandung: Alfabeta.CV
23. Sugiyono, 2012. *Understanding Qualitative Research*. Bandung: Alfabeta.CV

24. Van Wynsberge, S., Andréfouët, S., Gaertner-Mazouni, N., Wabnitz, C. C. C., Gilbert, A., Remoissenet, G., ...Fauvelot, C. (2016). Drivers of density for the exploited clam *Tridacna maxima*: a meta-analysis. *Fish and Fisheries*, 17(3), 567-584. <https://doi.org/10.1111/faf.12127>
25. Wilhm, J. L. and T. C. Dorris. 1968. *Biological Parameters for Water Quality Criteria*. Bio Scientific Publication. London. 18: 477-481.

UNDER PEER REVIEW