

Original Research Article

YIELD ENHANCEMENT OF FIELD PEA (*Pisumsativum*) THROUGH ZINC SOLUBILIZING MICROORGANISM

Abstract

A field experiment was conducted during rabi season 2022 at Instructional farm, School of Agriculture, SAGE University Bhopal (Madhya Pradesh) to evaluate the response of field pea to Zn and ZSB solubilizers. Results revealed that maximum plant height, dry matter accumulation (g), number of branches/plant, number of pods/plant, number of seeds/pod, seed (1060 kg/ha), straw (2130 kg/ha), biological yield (4243 kg/ha), Net return (1639rs/ha) and B:C Ratio (1.12) was obtained with application of 100% recommended dose fertilizer + Zinc and was found statistically different ($p < 0.05$) compared to mean values recorded in 100% ZSB treatment. Thus, all treatments significantly increased growth, yield and economics of field pea compared to the control.

Keywords: Field pea, Zinc, ZSB (Zinc solubilizing bacteria), Growth and yield. .

Introduction:

The main winter grain legume crop, field pea [*Pisum sativum*], is largely restricted to cooler temperate zones. This crop provides food, animal feed, and vegetables. It is also third-most-popular rabi crop after chickpea and arhar identified as significant pulse crop worldwide. Indian chickpea and arhar are followed by the pulse. To increase the production-productivity and productivity and production, of field pea in the study area, there is a need for a proper and better nutrient management practice. Field pea has shown a positive response to fertilizer application. However, consistent use of chemical fertilizers in modern agricultural practices has a negative impact on the population of native beneficial soil microorganisms and health of the soil. We can use Bio fertilizer on behalf of could be used as alternative to chemical because this bio-fertilizers are organic products which have a living cells. Bio-fertilizer plays a major role in unavailable to available form of nutrient to crops (Itelima *et al.* 2018; Ikeh *et al.* 2023) [1-9]. Bio-fertilizers are organic solution that contains microorganisms and growth hormone which boost crop productivity through biological nitrogen fixation, zinc solubilization, and other growth regulators that are also necessary for plants growth and development. Bio fertilizer are more sustainable in managing soil fertility (Ikeh *et al.* 2023) [11]. In this context, nutrient management combined with bio-fertilizers is emerging as a method of fertilization that is commercially viable and environmentally responsible (Kaur and Purewal, 2019)[2]-and (Ikeh *et al.* 2023)[10]

Comment [H1]: State the country

Comment [H2]: In full

Comment [H3]: State the experimental design used and treatments

Comment [H4]:

Comment [H5]:

Materials and Methods

The present investigation was performed in experimental field small plots of SAGE University Bhopal (M.P.) For the objective purpose of this study was to evaluate find out the effect of Zn and Zn solubilizers on growth and yield of field pea variety (plant growth and development were selected varieties of field pea: Prakash IPFD 1-1). Prakash IPFD 1-1 is characterised by large seeds and powdery mildew resistance. Its yellow ovoid shape grains are characterised by very high protein content. Field pea was sown on 23 November 2022. The field experiment was carried out in randomized block design with 6 treatments. The size of one plot was 6 square meter. The experiment was conducted at Instructional Cum Research Farm, School of Agriculture, SAGE University Bhopal (Madhya Pradesh) during rabi season of 2022. Geographically, Bhopal is situated at central region of India at 23.2599° N latitude and 77.4126° E longitude. Bhopal is known as city of lakes. The city is situated in the lowlands near the vindhya mountain range on the malwa plateau. Recommended dose of nitrogen, phosphorus and potassium @16.6g at basal dose in four treatments and ZSB (Zinc solubilizing bacteria) are applied common to all treatments. The result revealed that use of recommended dose of zinc 30-40 mg /ha. Sowing was done on 23rd November, 2022-11-22, while harvesting was done manually on 24th March, 2023-03-2023. Observation was recorded at 30, 60, 90 DAS and at harvesting of crop. Data collection Several observations were taken from each randomly selected plot at 30, 60, 90 days after planting (DAS) and randomly at harvesting during entire growth period of field pea. Sample Observed plants were tagged properly for further days. The following observations were taken on growth and yield attributes of field pea. That is are plant height (cm), number of branches/plant, number of pods/plant, number of seeds/pod, seed yield. Randomly five plant were selected for measuring height (cm) with the help of meter scale, number of branches/plant, number of pods/plant, number of seeds/pods and mean value was calculated. 1000 seed weight was taken in gram. Seed, straw and biological yield was observed in kg/ha and harvest index was computed. Net returns and B: C ratio were computed to evaluate the economics of the field experiment.

Result and Discussion

The observation of plant height was recorded at 60 and 90 DAS. The result revealed that plant height was minimum upto 30 DAS but it was increased during 60 DAS and at harvest due to increase in the age of crop. The plant height value ranges from 65.20 cm. to 84.34 cm. at harvest. Significantly, maximum plant height value was 84.34 cm was recorded at harvest under T₂: recommended dose fertilizer + Zn 100% (9gm.) and which was found superior when compared to the among the treatments. The Under-recommended dose fertilizer + Zn 100% (19.68 gm.) produced resulted in significantly higher dry matter accumulation (g/plant) in crop at harvest compared the other over rest of the treatments (Table 1). Data (Table 2) shows that application of Zn and ZSB Levels, significantly increased number of branches per /plant, number of pods per /plant and number of seeds per /pod. Treatment recommended dose fertilizer + Zn 100% resulted in maximum branches per/plant (2.46), pods per/plant (16.61) respectively and seeds per/pod

Comment [H6]:

Comment [H7]:

Comment [H8]: State how the land was prepared

Comment [H9]:

Formatted: Superscript

Formatted: Superscript

(6.85) against over rest of the treatments. Results revealed that application of recommended dose of fertilizer (RDF) + Zn (100%) T₂ resulted in significantly higher seed, straw and biological yield over control. Maximum seed (1060 kg/ha), straw (2130 kg/ha) and biological (4243 kg/ha) yield was recorded with application of recommended dose fertilizer + Zn 100% respectively (Table 2). Treatment recommended dose fertilizer + Zn 100% was statistically higher significant with Zinc solubilizing bacteria (ZSB) (100%). This might have increased number of pods/plant, seeds/pod, seed, straw and biological yield. The economical perusal of data (Table 2) showed that among Zn and ZSB Solubilizer fertilizer Treatment recommended dose fertilizer + Zn 100% resulted in highest and significant followed by T₃. These results confirm with the earlier finding of Venkatrao *et al.* (2017) [8] and Kuniya *et al.* (2018) [4]. The beneficial effect of zinc solubilizing bacteria on yield attributing characteristics have also been recorded by Kothyari *et al.* (2017) [3] and Raut (2018) [5] respectively. Test weight was found non-significant with respect to various Zn fertilizers and their solubilizers' levels. The economical perusal of data (Table 2) showed that among Zn fertilizer treatments, Recommended dose of fertilizer + 100% Zinc resulted in highest and significant net return (1639 ₹/ha) and cost-benefit ratio (1.12). Similar findings were also reported by Serawat *et al.* (2018) [6] and Singh *et al.* (2018) [7].

Table.1 Yield enhancement of fieldpea through Zinc solubilizing microorganisms on growth and yield attributes (at harvest).

Treatments	Plant Height (cm) at harvest in g	Dry matter Accumulation (g/plant)	Number of Branches/plant	Number of Pods/plant	Seed/pods	Test weight (g)
Zn and ZSB Levels						
control (T ₁)	65.20	13.12	1.04	9.94	4.42	175.36
Recommended Dose Fertilizer + Zn (100%) (T ₂)	84.34	19.68	2.46	16.61	6.85	189.54
Zinc solubilizing bacteria (ZSB) (100%) (T ₃)	80.53	19.61	2.15	15.53	6.24	185.62
Recommended dose fertilizer + Zn +ZSB (75%) (T ₄)	75.20	17.82	1.78	13.78	5.78	182.41
Recommended dose fertilizer + Zinc +ZSB (50%)	71.48	15.7	1.70	12.63	5.23	179.45

Recommended dose fertilizer + Zinc +ZSB (25%) (T ₆)	69.91	14.9	1.25	11.5	4.5	176.69
SE(m)	1.52	0.46	0.07	0.31	0.18	3.84
CD at 5%	4.65	1.34	0.23	0.95	0.57	NS




Fig 1: Graphical representation of the treatments

Table.2 Yield enhancement of field pea through zinc solubilizing microorganism on Seed, straw, biological yield and harvest index.

Treatments	Seed yield (kg/ha.)	Straw Yield (kg/ha.)	Biological yield (kg/ha.)	Harvest index (%)	Net return (Rs./ha)	B:C ratio
Zn and ZSB Levels						
Control (T ₁)	250	560	1005	30.06	820	0.52
Recommended Dose fertilizer + Zn (100%) (T ₂)	1060	2130	4243	32.26	1639	1.12
Zinc solubilizing bacteria (ZSB) (100%) (T ₃)	830	1655	3298	32.09	1460	1.03
Recommended dose fertilizer + Zn +ZSB (75%) (T ₄)	700	1390	2680	31.45	1234	1.01
Recommended dose fertilizer + Zn +ZSB (50%) (T ₅)	480	940	1860	31.15	1140	0.95

Recommended dose fertilizer + Zn +ZSB (25%) (T ₆)	380	735	1438	30.62	960	0.76
SE(m)	20.4	40.7	95.3	0.29	161.34	0.02
CD at 5%	61.1	120.3	285.4	0.86	484.05	0.07

Conclusion

~~This study compared to control plot various treatment which was T₂(Zn + 100 RDF) was found a higher significant followed by T₃(Zinc solubilizing bacteria 100%) Zn and ZSB Solubilizes treatment shown resulted to in significant increase in the growth parameters, yield attributes and yields of field pea crop and ultimately resulted in higher net return and B:C ratio. Therefore, recommended Dose fertiliser + Zn (100%) was recorded to field pea farmers in the study area, for high yield and economic benefits to farmers, recorded significantly higher growth, yields, net return and B: C ratio of field pea crop over rest of the treatments.~~

References

1. Itelima JU, Bang WJ, Onyimba IA, Egberé OJ. A review: biofertilizer; a key player in enhancing soil fertility and crop productivity. *Journal of Microbiology and Biotechnology Report*. 2018;2(1):22-28.
2. Kaur P, Purewal SS. Biofertilizers and their role in sustainable agriculture. *Biofertilizers for Sustainable Agriculture and Environment Soil Biology*. 2019;55:285- 300.
3. Kothiyari HS, Yadav LK, Jat R, Gurjar PC. Influence of Biofertilizers on Plant Growth and Seed Yield of Pea (*Pisumsativum* [L.]). *International Journal Current Microbiology and Applied Science*. 2017;6(11):1810- 1817.
4. Kuniya N, Chaudhary N, Patel S. Effect of sulphur and zinc application on growth, yield attributes, yield and quality of summer clusterbean (*Cyamopsistetragonoloba* [L.]) in light textured soil. *International Journal of Chemical Studies*. 2018;6(1):1529-1532.
5. Raut AD, Durgude AG, Kadlag AD. Effect of zinc solubilizing bacteria on zinc use efficiency and yield of summer groundnut grown in Entisol. *International Journal of Chemical Studies*. 2019; 7(1):1710-1713.
6. Serawat A, Sharma Y, Serawat M, Kapoor A, Jakhar RK. Effect of Phosphorus and Sulphur on Growth attributes and Yield of mung bean (*Vignaradiata* [L.]). *International Journal of Current Microbiology and Applied Sciences*. 2018;7(8):2674-2683.
7. Singh N, Joshi E, Sasode DS, Sikarwar RS, Rawat GS. Liquid Biofertilizer and Inorganic Nutrients Effect on Physiological, Quality Parameters and Productivity of Kharif Groundnut (*Arachishypogaea* [L.]) *International journal Current Microbiology and Applied Science*. 2018;7(09):729-735.

Formatted: Normal, Justified, Indent: Left: -0.69", Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0" + Indent at: 0"

Formatted: Font: Not Italic, English (United States)

8. Venkatarao CV, Naga SR, Yadav BL, Koli DK, Rao IJ. Effect of phosphorus and biofertilizers on growth and yield of mungbean [Vignaradiata (L.) Wilczek]. *International Journal of Current Microbiology and Applied Sciences*. 2017;6(7):3992-3997.

9. Ikeh, A.O., Okocha, I.O., Umekwe, P.N., Amanze, A.N and Ikeh, C.E. Effect of foliar application of cow dung extract on growth and yield of waterleaf (*Talinum triangulare* Jacq.) in an ultisol. *Journal of Current Opinion in Crop Science*, 2023; 4(3), 103-111.

10. Ikeh, A. O., Ndaeyo, N.U and Ikeh, C.E, Effects of integrated fertilization on soil sustainability and cassava (*Manihot esculenta* Crantz) yield in an ultisol. *Journal of Current Opinion in Crop Science*, 2023; 4(2): 89-102.

11. Ikeh, A.O., J. O. Orji, J. O., Sampson, H. U. and Akata, O. R. (2023). Effect of Oil Palm Bunch Refuse Ash in Sustainable Production of Egusi-Melon (*Colocynthiscitrullus*) in an Ultisol. *Asian Journal Agricultural and Horticultural Research*, 2023;10 (3) 138-148. DOI: 10.9734/AJAHR/2023/v10i324.

7-12.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Normal, Justified, Indent: Left: -0.69", Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0" + Indent at: 0"

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, English (India)

Formatted: Font: Not Italic, English (United States)

Formatted: Indent: Left: -0.69", No bullets or numbering

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Justified, Indent: Left: -0.69", Space After: 0 pt, Line spacing: single, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0" + Indent at: 0"

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: List Paragraph, Left, No bullets or numbering

Formatted: Justified, Indent: Left: -0.69", Space After: 0 pt, Line spacing: single, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0" + Indent at: 0"

Formatted: Font: Not Italic

Formatted: Font:

Formatted: Font: 12 pt

Formatted: Indent: Left: -0.69", Line spacing: single