

IMPACT OF SEAWEED EXTRACT ON GROWTH OF CARROT (*Daucus carota* . L)var. Super kuroda

ABSTRACT: A field experiment was conducted during *rabi* 2022 on carrot variety ‘Super kuroda’ in sandy loam soil at Horticultural Garden, Professor Jaya Shankar Telangana State Agriculture University, Rajendranagar, Hyderabad. To study the effect of seaweed extract on the growth parameters of carrot. The experiment was laid out in randomized block design (RBD) with eight treatments and replicated thrice. The treatments include various sources of bio stimulants *viz.*, Seaweed extract liquid, seaweed granules, vermicompost and vermiwash given as foliar application on three stages *viz.*, first spray on 2-3 leaves stage and there after at every 20 days interval and as soil application twice at the time of sowing as well as 30 DAS. The various growth parameters were recorded. The results revealed that on application of RDF + soil application of sea weed extract twice at time of sowing and at 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water significantly increased the growth parameters *viz.*, plant height, number of leaves per plant, chlorophyll content.

Key words: carrot, sea weed extract, plant height, number of leaves per plant, chlorophyll content.

1. INTRODUCTION :

Plant bio stimulants, or agricultural bio stimulants, are a diverse group of substances that can find application in farming to improve crop productivity, and the crop’s ability to withstand biotic and abiotic stress. These bio stimulants are typically derived from organic sources and contain bio-active compounds. Common components found in bio stimulants include mineral elements, humic substances (HSs), vitamins, amino acids, chitin, chitosan, poly- and oligosaccharides and seaweed extract (Kauffman *et al.*, 2007). In recent trends, seaweed has gained popularity. Seaweed extracts are important in sustainable agriculture because of its composition as it contains phytohormones such as auxins, gibberrelins, cytokinins, and betaines (Crouch and van Staden, 1993) and it also contains inorganic compounds like nitrogen, phosphorous, potassium, calcium, iron, magnesium, zinc, sodium and sulphur (Rayorathet *al.*, 2009). Hence, it plays a crucial role in sustainable agriculture due to its biodegradable, non-polluting and non-hazardous nature for both humans and animals, and does not harm the environment, (Dhargalkar and Pereira, 2005)

Carrot is scientifically known as *Daucus carota*, belongs to umbelliferae family. It is the most popular and widely consumed vegetable worldwide. They are known for remarkable nutritional benefits and they have a very low fat index which makes them be included in dietary practices. Carrot is abundant in essential nutrients, including vitamin A, vitamin C, dietary fibers, minerals like

calcium, iron, and amino acids. It has anti-carcinogenic properties Zaini *et al* . (2011). Keeping the above points in view the experiment was planned to study the effect of seaweed extract on growth parameters of carrot.

2. Material and methods:

The present experiment was conducted during *rabi* season 2022 in sandy loam soils at Horticultural Garden, Professor Jayashankar Telangana state Agriculture University, Rajendra nagar, Hyderabad.. The experiment was laid out in randomized block design comprising of eight treatments and replicated thrice **T₁** - RDF, **T₂** - RDF+ soil application of sea weed extract twice at the time of sowing and at 30 DAS, **T₃** - RDF+ foliar spray of sea weed extract thrice 2.5 ml l⁻¹ of water, **T₄** - RDF+ soil application of sea weed extract twice at time of sowing and at 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water, **T₅** - RDF+ vermicompost 5 t ha⁻¹, **T₆** - RDF+ foliar spray of water 3 times, **T₇** - RDF+ foliar spray of 10% of vermiwash, **T₈** - Soil application of sea weed extract twice at the time of sowing and at 30 DAS + foliar spray of sea weed extract thrice 2.5 ml l⁻¹ of water. Super kuroda variety was sown on November 3rd 2022 with a spacing of 30 cm x 5 cm. The RDF used was 50 : 40 : 50 kg ha⁻¹ N, P₂O₅ and K₂O respectively. For the treatment, commercially available seaweed liquid “Sagarika” and granules “ Sagarika Gold ” were used and are manufactured by IFFCO. Spraying was done through a knapsack sprayer using 500 liters of water per hectare. Biometric observations were taken on tagged five representative plants selected at random from each treatment of the net plot and the mean values were presented.

3. RESULTS AND DISCUSSION :

3.1 Plant height (cm)

All treatments had shown an impact on plant height of carrot, with gradual improvement observed as the days after sowing (DAS) increased. The effects of seaweed extract on carrot plant height at 30, 60 and at 90 DAS are summarized in Table 1.

At 30 DAS, there was no significant difference was noticed among the various treatments with respect to the plant height . In contrast, notable disparity in plant height was observed at 60 DAS. The highest plant height (18.43) was recorded with RDF + soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water . This treatment was statistically on par with RDF+ foliar spray of seaweed extract thrice 2.5 ml l⁻¹ of water and RDF + soil application of seaweed extract at time of sowing and at 30 DAS. The

next most effective treatments viz., RDF + foliar spray of 10 % of vermiwash, RDF + vermicompost 5 t ha⁻¹, RDF alone and RDF + foliar spray of water 3 times. The lowest plant height (9.18) was application of soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water

Similarly, at 90 DAS, the treatment that led to the highest plant height (25.63) was RDF + soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water . This was statistically similar to RDF+ foliar spray of seaweed extract thrice 2.5 ml l⁻¹ of water (24.63) and RDF + soil application of seaweed extract at time of sowing and 30 DAS (24.55). The lowest plant height was observed in the treatment soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water (15.43).

The increase in plant height can be attributed due to the provision of essential primary, secondary, and trace nutrients through the application of seaweed extract, similar results by Lee, (2010).

3.2 Number of leaves per plant:

Much like the observation for plant height at 30 days after sowing (DAS), there were no significant difference in the number of leaves per plant when seaweed extract was applied. However, at 60 and 90 DAS, there was an increase in the number of leaves per plant. The highest number of leaves were observed in the treatment RDF + soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water . This was statistically similar to the treatments RDF+ foliar spray of seaweed extract thrice 2.5 ml l⁻¹ of water and RDF + soil application of seaweed extract at time of sowing and 30 DAS. the treatment RDF + foliar spray of 10% vermiwash, RDF + vermicompost 5 t ha⁻¹, RDF alone, and RDF + foliar spray of water three times showed comparable results. Conversely, the lowest number of leaves per plant was observed in soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray of seaweed extract thrice at 2.5 ml l⁻¹ of water.

The increase in the number of leaves may be ascribed due to the presence of micro and macro nutrients, as well as phytohormones found in seaweed extract. These findings align with the results of Mooney and Vastaden, (1985) and Blunden (1991).

3.3 Chlorophyll content (SPAD):

Initially, at 30 DAS there were no discernible difference was observed among the treatments with respect to chlorophyll content at 30 DAS. However, at 60 and 90 DAS, significant impact of seaweed extract was noticed. The highest chlorophyll content was observed in T4 RDF + soil application of sea weed extract twice at time of sowing and 30 DAS + foliar

spray thrice 2.5 ml l⁻¹ of water. Treatments with RDF+ foliar spray of seaweed extract thrice 2.5 ml l⁻¹ of water and RDF + soil application of seaweed extract at the time of sowing and 30 DAS were statistically on par with RDF + soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water. The next best treatments were RDF+ foliar spray of seaweed extract thrice 2.5 ml l⁻¹ of water and RDF + soil application of seaweed extract at time of sowing and 30 DAS. The treatment RDF + foliar spray of 10% vermiwash, RDF + vermicompost 5 t ha⁻¹, RDF alone, and RDF + foliar spray of water three times . The treatment of lowest chlorophyll content was observed oil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray of seaweed extract thrice at 2.5 ml l⁻¹ of water.

The significant increase in chlorophyll content can be attributed due to an elevated production of chloroplasts, a reduction in the breakdown of chlorophyll, and a delayed aging process in plants. This was observed in studies conducted by Janninet *et al.* (2013). Furthermore, cytokinin, a plant hormone, plays a protective role in preserving chloroplasts (as demonstrated by Zavaleta-Mancera *et al.*, 2007) and promoting chloroplast division Okazaki *et al.* (2009).

Treatments	Plant height (cm)		
	30 DAS	60 DAS	90 DAS
T ₁ - RDF	9.15	13.39	23.13
T ₂ - RDF + soil application of seaweed extract at time of sowing and 30 DAS	9.39	17.70	24.55
T ₃ - RDF+ foliar spray of seaweed extract thrice 2.5 ml l ⁻¹ of water	9.35	17.77	24.63
T ₄ -RDF + soil application of sea weed extract twice at time	9.35	18.43	25.63

of sowing and 30 DAS + foliar spray thrice 2.5 ml l ⁻¹ of water			
T ₅ - RDF + vermicompost 5 t ha ⁻¹	9.26	13.53	23.43
T ₆ - RDF + foliar spray of water 3 times	9.10	13.08	22.82
T ₇ - RDF + foliar spray of 10 % of vermiwash	9.26	14.73	23.69
T ₈ - soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l ⁻¹ of water	9.18	9.21	15.43
SEm±	0.07	0.31	0.43
CD (P = 0.05)	NS	1.0	1.3

Table no 1. Impact of seaweed extract on plant height

Treatments	Number of leaves		
	30 DAS	60 DAS	90 DAS
T ₁ - RDF	3.67	7.67	10.33
T ₂ - RDF + soil application of seaweed extract at time of sowing and 30 DAS	4.00	9.67	11.62
T ₃ - RDF+ foliar spray of seaweed extract thrice 2.5 ml l ⁻¹ of water	4.00	10.00	11.80

T ₄ -RDF + soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l ⁻¹ of water	4.00	10.67	12.11
T ₅ - RDF + vermicompost 5 t ha ⁻¹	3.66	8.33	10.67
T ₆ - RDF + foliar spray of water 3 times	3.66	7.00	7.33
T ₇ - RDF + foliar spray of 10 % of vermiwash	4.00	9.00	10.80
T ₈ - soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l ⁻¹ of water	3.45	4.49	6.46
SEm±	0.19	0.15	0.34
CD (P = 0.05)	NS	0.5	1.0

Table no 2. Impact of seaweed extract on number of leaves per plant in carrot.

Table 3 Impact of seaweed extract on chlorophyll content of carrot leaves

Treatments	Chlorophyll content (SPAD)		
	30 DAS	60 DAS	90 DAS
T ₁ - RDF	3.67	7.67	10.33
T ₂ - RDF + soil application of seaweed extract at time of sowing and 30 DAS	4.00	9.67	11.62

T ₃ - RDF+ foliar spray of seaweed extract thrice 2.5 ml l ⁻¹ of water	4.00	10.00	11.80
T ₄ -RDF + soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l ⁻¹ of water	4.00	10.67	12.11
T ₅ - RDF + vermicompost 5 t ha ⁻¹	3.66	8.33	10.67
T ₆ - RDF + foliar spray of water 3 times	3.66	7.00	7.33
T ₇ - RDF + foliar spray of 10 % of vermiwash	4.00	9.00	10.80
T ₈ - soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l ⁻¹ of water	3.45	4.49	6.46
SEm±	0.19	0.15	0.34
CD (P = 0.05)	NS	0.5	1.0

Conclusion

Results of the experiment revealed that application of RDF + soil application of sea weed extract twice at time of sowing and 30 DAS + foliar spray thrice 2.5 ml l⁻¹ of water improves the plant growth parameters of the carrot. Hence, it could be recommended to the farmers.

References:

- Blunden, G., Jenkins, T., Liu, Y., 1997. Enhanced leaf chlorophyll levels in plants treated with seaweed extract. *Journal of Applied Phycology*. 8: 535–543.
- Crouch, I.J., Van Staden, J., 1993. Evidence for the presence of plant growth regulators in commercial seaweed products. *Plant Growth Regulation*. 13: 21–29.
- Dhargalkar, V.K. and Pereira, N. 2005. Seaweed: promising plant of the millennium. *Science and culture*. 71(3-4): 60-66.
- Jannin, L., Arkoun, M., Etienne, P., Laîné, P., Goux, D., Garnica, M., Fuentes, M., Francisco, S.S., Baigorri, R., Cruz, F., 2013. Brassica napus growth is promoted by *Ascophyllum nodosum* (L.) Le Jol. seaweed extract: microarray analysis and physiological characterization of N, C, and S metabolisms. *Journal of Plant Growth Regulation*. 32: 31–52.
- Kauffman, G. L., Kneivel, D. P., and Watschke, T. L. (2007). Effects of a biostimulant on the heat tolerance associated with photosynthetic capacity, membrane thermostability, and polyphenol production of perennial ryegrass. *Crop Science* .

- Lee, J., 2010. Effect of application methods of organic fertilizer on growth, soil chemical properties and microbial densities in organic bulb onion production. *Scientia Horticulturae*, 124(3): 299-305.
- Mooney, P.A. and Van Staden, J., 1986. Algae and cytokinins. *Journal of plant physiology*. 123(1). 1-21.
- Okazaki, K., Kabeya, Y., Suzuki, K., Mori, T., Ichikawa, T., Matsui, M., Nakanishi, H., Miyagishima, S. 2009. The plastid division 1 and 2 components of the chloroplast division machinery determine the rate of chloroplast division in land plant cell differentiation. *Plant Cell*. 21: 1769–1780.
- Zaini, R., Clench, M.R. and Maitre, C.L. (2011) Bioactive Chemicals from Carrot (*Daucus carota*) Juice Extracts for the Treatment of Leukemia. *Journal of Medicinal Food*, 14: 1303-1312.
- Zavaleta-Mancera, H.A., Lopez-Delgado, H., Loza-Tavera, H., Mora- Herrera, M., Trevilla-Garcia, C., Vargas-Suarez, M., Ougham, H., 2007. Cytokinin promotes catalase and ascorbate peroxidase activities and preserves the chloroplast integrity during dark- senescence. *Journal of Plant Physiology*. 164: 1572–1582.