

## Original Research Article

# Effect of Nutrient Management Practices on Carbon Pools Following 13 Years of Cropping With Soybean Based Cropping Systems In Vertisols of Central India

### ABSTRACT

Organic manure application has its significant impact on the soil health. Low organic matter in tropical soils is a major factor contributing to their poor productivity. Soil properties have been continuously influenced by the management practices and land uses, in which a latter one has been identified as profound influence on soil properties especially on soil organic carbon. A thirteenth year experiment on soybean based cropping system in a vertisol of central india under organic farming was used for this investigation. An investigation was carried out on “Soil organic carbon dynamics under long-term nutrient management in soybean based cropping system” at the Indian Institute of Soil Science, Bhopal on an on-going research project on organic farming. The effect of organic, integrated and inorganic nutrient management was assessed in three cropping systems viz. soybean (JS 335)-wheat (Malwa Shakti), soybean-mustard (Pusa Bold) and soybean-gram (JG 130) on aggregate size fractions, carbon content in aggregate as well as soil organic carbon pools dynamics on a split plot experimental design with three replications. The study relevant to dynamics of soil organic carbon pools revealed higher content of soil organic carbon, labile carbon, water soluble carbon, soil microbial biomass carbon (SMBC) as well as dehydrogenase activity (DHA) that varied between 0.86 and 1.04 percent; 440 and 538 mg kg<sup>-1</sup>, 52.97 and 70.43; 288 and 375 mg kg<sup>-1</sup>, 88 and 137 µg TPF g<sup>-1</sup> soil d<sup>-1</sup>, respectively in surface 0-15 cm soil under organic nutrient management.

Keyword- Carbon Pools, nutrient management, cropping systems

## 1. Introduction

Long term experiments are primary source of information determining the effect of cropping systems, soil management, fertilizer use and residue utilization on changes on SOC (Leigh and Johnston, 1994). Long-term application of farmyard manure and vermi-compost in crop production can contribute to both sustainable food production and mitigation of greenhouse gas emissions through soil C sequestration.

Large scale implementation of the organic manure amendments will help in enhancing the capacity of carbon sequestration and promote food security in the region (Parmar *et al.*, 2015). In response, new management strategies have emerged, including soil-focused approaches such as no-tillage, which aim to improve soil regulating and supporting ecosystem services by reducing soil disturbance (Williamset *al.*, 2016) **Not In References NIR**. Aggregation results from the rearrangement of the soil constituent particles, flocculation and cementation and is arbitrated by SOC, biota, ionic bridging, clay and oxide content. The aggregate dynamics vary among different crops, crop rotations and cover crops (Jarecki and Lal, 2003). Stability of aggregate is the measure of the structural stability of soils (Six *et al.*, 2000). Any reduction in soil aggregate stability is a powerful early indicator of the onset of land degradation. Favorable soil aggregation is important to improve soil fertility and quality with particular emphasis on SOC sequestration, increasing agronomic productivity, enhancing porosity and decreasing erodibility. Vegetation cover can influence soil aggregate stability because of contribution of organic matter through litter and plant root turnover (Angers and Caron, 1998), root exudates rizhodeposition (Gysselet *al.*, 2005; Pohl *et al.*, 2009). Hence, there is strong need to evaluate the effect of different nutrient inputs on soil health in Vertisols, as these soils occupy about 35% of the area under cultivation in India. Also, soybean-wheat, is the predominant cropping system in deep Vertisols of central India along with soybean-chickpea and soybean-mustard cropping systems. Keeping above in view the limited information available on the effect of nutrient management practices in prominent cropping systems in central India, the present investigation was carried out to assess the effect of nutrient management practices under different cropping systems in black soils of central India.

## **2. Material and Methods**

### **2.1 Experimental site, climate and soil characteristics**

The field experiment was conducted during *kharif* season of 2017-18. at the research farm, ICAR-IISS, Bhopal, Madhya Pradesh The on-going experiment on Network project on organic farming was used for this investigation. The study area falls under semi-arid and sub-tropical zone characterized by hot summer and cold winter. Mean annual precipitation is about 1146 mm, most of which is received during

the monsoon period of July to September. The average maximum temperature during summer is 35°C, while the average minimum temperature during winter is 4°C.

## 2.2 Treatments detail

The on-going experiment on Network project on organic farming was used for this investigation. Soybean (cv JS 335), wheat (cv Malwa Shakti), mustard (cv Pusa Bold) and chickpea (cv JG-130) were sown during kharif and rabi seasons at a spacing of 45X5; 22.5; 45X10 and 30X10 cm as per standard cultural practices in split plot design. The recommended dose of fertilizer for these crops was 30:60:20; 80:40:40; 60:40:0 and 30:40:0 for N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O, respectively. The plot size of the experiment is 90 sq meters.

Soil samples (0-5, 5-15 and 15-30 cm) were collected and homogenized from long-term experiment on organic farming at Indian Institute of Soil Science under soybean-wheat, soybean-mustard, soybean-gram cropping systems. Visible litter and roots were picked out by hand. Part of the soil samples were air-dried at room temperature and gently passed through a < 2 mm sieve for determining different soil organic carbon pools, biological properties and other soil properties.

Soil organic carbon content (%) in soil samples was determined in soil passed through 0.2 mm sieve by (Walkley and Black 1934) method. A known weight of soil was treated with an excess volume of standard K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> in the presence of concentrated H<sub>2</sub>SO<sub>4</sub>. The excess of K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> not reduced by the organic matter was titrated back against a standard solution of ferrous ammonium sulphate, in the presence of diphenylamine indicator.

Labile carbon in soil samples was analysed by using procedure outlined by dilute slightly alkaline KMnO<sub>4</sub> which reacts with the most readily oxidizable (active) forms of soil carbon as per procedure described by (Blair *et al.*, 1995). and (modified by Weil *et al.*, 2003). The results were expressed as mg kg<sup>-1</sup>.

The water soluble carbon in soil was extracted by shaking 10 g soil in 20 ml distilled water for one hour as described by (McGill *et al.*, 1986). Then water soluble organic carbon was estimated in the extract by using Shimadzu make TOC (Total Organic Carbon) analyser. The result was expressed as mg kg<sup>-1</sup>.

Microbial biomass was determined by the incubation and fumigation technique (Jenkinson and Powlson, 1976). The soil samples were subjected to chloroform fumigation, which causes cell walls to lyse and denature. The soil was

then extracted with 0.5 M K<sub>2</sub>SO<sub>4</sub>. The readily oxidizable carbon contained in the extract was measured through standard chemical procedures.

Biological activity of a soil is the function of number of organisms present in soil coupled with their physiological efficiency. Dehydrogenase activity was measured and expressed as the rate of formation of TPF from TTC as per method outlined by (Casida 1977). The result was expressed as µg TPF g<sup>-1</sup> soil d<sup>-1</sup>.

### 3. RESULTS AND DISCUSSION

#### 3.1 Easily oxidizable carbon

The data on water based carbon (WBC) or easily oxidizable carbon has been presented in Table.1. The easily oxidizable carbon in surface 0-5 cm soil depth varied between 0.72 and 1.13 percent under different cropping systems and nutrient management practices. The organic nutrient management registered highest organic carbon content under different cropping systems with a mean value of 1.05 percent. The WBC was significantly lower in integrated and inorganic nutrient management where it was found to be 0.86 and 0.72 percent respectively. Among cropping systems soybean – wheat cropping system registered highest organic carbon content which was statistically at par with soybean gram but statistically higher over soybean mustard cropping system. In 5-15 cm soil depth, soil organic carbon varied between 0.90 and 0.96 percent under organic sources of nutrients with a mean value of 0.91 percent. Integrated sources of nutrients resulted in significantly lower soil organic carbon as compared to organic source of nutrient and ranged between 0.79 and 0.83 among different cropping systems with a mean value of 0.81 percent. Application of inorganic sources of nutrients resulted in lowest value of soil organic carbon with a mean value of 0.62 percent. The effect of different cropping systems was not found significant with respect to WBC in 5-15 cm soil depth. Soil organic carbon varied between 0.54 and 0.58 percent in 15-30 cm soil depth under organic sources of nutrients with a mean value of 0.56 percent. Jha *et al.* (2014) (NIR) observed an increment of 49.1% in TOC with the application of FYM @ 15 t ha<sup>-1</sup> Y<sup>-1</sup> along with recommended dose of NPK on long term basis (38 years) over control in a Vertisol of Jabalpur under soybean-wheat sequence. Similarly, also reported 105 and 71% higher TOC in long term organic farming practice over absolute control and recommended dose of NPK fertilizers, respectively under soybean-wheat cropping system. Manna *et al.* (2012) (NIR) and Lakaria *et al.* (2012<sub>b</sub>) (NIR) also found increase in

WBC with the application of FYM alone Lakaria *et al.* (2012<sub>a</sub>) or in combination with recommended NPK fertilizers over absolute control and sole NPK fertilizer application. The results of this study are in close agreement with these findings. The higher C accumulation in the Vertisol may be attributed to their high silt+clay content which increase the C stabilization capacity (Six *et al.*, 2002; Jha *et al.*, 2012<sub>a</sub>).

**Table.1 Effect of nutrient management on easily oxidizable carbon (%) under different cropping systems**

Treatment	Soybean-Wheat	Soybean-Mustard	Soybean-Gram	Mean
<b>0-5 cm</b>				
Organic	1.13	1.00	1.04	1.05
Integrated	0.90	0.83	0.85	0.86
Inorganic	0.72	0.73	0.73	0.72
Mean	0.91	0.85	0.87	
LSD (p=0.05)	CS	NM	CS X NM	
	0.04	0.06	NS	
<b>5-15 cm</b>				
Organic	0.96	0.90	0.86	0.91
Integrated	0.83	0.80	0.79	0.81
Inorganic	0.63	0.61	0.63	0.62
Mean	0.81	0.77	0.76	
LSD (p=0.05)	CS	NM	CS X NM	
	NS	0.03	NS	
<b>15-30 cm</b>				
Organic	0.58	0.55	0.54	0.56
Integrated	0.48	0.48	0.45	0.47
Inorganic	0.48	0.44	0.42	0.45
Mean	0.51	0.49	0.47	
LSD (p=0.05)	CS	NM	CS X NM	
	NS	0.03	NS	

CS- cropping system, NM – nutrient management

### 3.2 Water soluble carbon content

Water soluble carbon content in different soil depths have been presented in Table.2. In 0-5 cm soil depth, the water soluble carbon varied widely among the different nutrient management options. Under organic sources of nutrient application the water soluble carbon ranged between 60.86 and 70.43 mg kg<sup>-1</sup> with a mean value of 64.52 mg kg<sup>-1</sup>. With integrated system of nutrient application the water soluble carbon ranged between 50.78 and 58.47 mg kg<sup>-1</sup> with a mean value of 53.89 mg kg<sup>-1</sup>. Least water soluble carbon in 0-5 cm was recorded in inorganic nutrient management where it varied between 43.49 to 48.45 mg kg<sup>-1</sup> among cropping systems with a mean value of 46.03 mg kg<sup>-1</sup>. In 5-15 cm soil depth, the water soluble carbon decreased as compared to surface layer and it varied between 42.61 and 56.57 mg kg<sup>-1</sup> among the cropping systems. Under organic sources of nutrient application the water soluble carbon ranged between 52.97 and 56.57 mg kg<sup>-1</sup> with a mean value of 54.37 mg kg<sup>-1</sup> across the plots. Integrated system of nutrient application resulted in slightly lower water soluble carbon but a significant decrease could be recorded only with inorganic nutrient management (45.07 mg kg<sup>-1</sup>). There was no significant difference between the cropping systems with respect to water soluble carbon in 5-15 cm soil depth. Water soluble carbon varied widely among the different nutrient management options in 15-30 cm soil depth. It was registered the highest in organic treatment where ranged between 47.50 and 50.19 mg kg<sup>-1</sup> with a mean value of 48.82 mg kg<sup>-1</sup> across the cropping systems (Table 2). Integrated nutrient management resulted in significantly lower water soluble carbon which varied between 41.11 and 45.18 mg kg<sup>-1</sup> with a mean value of 42.88 mg kg<sup>-1</sup>. Under only inorganic addition of nutrients the water soluble carbon has been found to vary from 37.85 to 45.15 mg kg<sup>-1</sup> with a mean value of 39.75 mg kg<sup>-1</sup>. Lakaria *et al.* (2012<sub>c</sub>) also found that the WSC was ranged from 30.9 mg kg<sup>-1</sup> to 101.6 mg kg<sup>-1</sup> under different land use whereas Jha *et al.* (2012<sub>b</sub>) recorded WSC in the range of 13.8 to 101.6 mg kg<sup>-1</sup> under different treatments in Vertisols.

**Table.2 Effect of nutrient management on Water soluble carbon (mg kg<sup>-1</sup>) under different cropping systems**

Treatment	Soybean-Wheat	Soybean-Mustard	Soybean-Gram	Mean
-----------	---------------	-----------------	--------------	------

<b>0-5 cm</b>				
Organic	70.43	62.27	60.86	64.52
Integrated	58.47	50.78	52.43	53.89
Inorganic	48.45	46.15	43.49	46.03
Mean	59.11	53.06	52.26	
LSD (p=0.05)	CS	NM	CS X NM	
	NS	4.34	NS	
<b>5-15 cm</b>				
Organic	56.57	53.57	52.97	54.37
Integrated	53.03	51.75	48.86	51.21
Inorganic	42.61	49.62	42.98	45.07
Mean	50.74	51.64	48.27	
LSD (p=0.05)	CS	NM	CS X NM	
	NS	3.32	NS	
<b>15-30 cm</b>				
Organic	50.19	48.78	47.50	48.82
Integrated	41.11	45.18	42.34	42.88
Inorganic	37.85	41.50	39.89	39.75
Mean	43.05	45.15	43.25	
LSD (p=0.05)	CS	NM	CS X NM	
	NA	3.69	NA	

CS- cropping system, NM – nutrient management

### 3.3 Labile carbon

Potassium permanganate oxidizable carbon (labile carbon) determined in different cropping systems with nutrient management options has been presented in Table 3. In 0-5 cm soil depth, labile carbon content also varied widely among the different nutrient management options. The content of labile carbon under different sources of nutrient application varied between 417 to 538 mg kg<sup>-1</sup> with a mean value of 515 mg kg<sup>-1</sup>. With integrated nutrient management practice the labile carbon varied

between 479 and 484 mg kg<sup>-1</sup>. Among different cropping system no significant difference could be observed in labile carbon content, however, the interaction among nutrient management and cropping systems was significant. In 5-15 cm soil depth, labile carbon content also varied widely among the different nutrient management options. Labile carbon under organic sources of nutrient varied from 440 to 470 mg kg<sup>-1</sup> with a mean value of 450 mg kg<sup>-1</sup>. It was decreased with addition of only integrated nutrient it varied between 423 and 446 mg kg<sup>-1</sup> with a mean value of 431 mg kg<sup>-1</sup>. Labile carbon In 15-30 cm soil depth, was found the lowest where it varied between 341 and 391 mg kg<sup>-1</sup> with a mean value of 374 mg kg<sup>-1</sup>. Integrated system of nutrient resulted in lower labile carbon as compared to organic nutrient management and it varied between 329 and 365 mg kg<sup>-1</sup> with a mean value of 344 mg kg<sup>-1</sup>. Hassink (1995) ;Lakaria et al. (2012b) NIR and Jha et al. (2012b) also observed the KMnO<sub>4</sub>-C content between 463 and 621 mg kg<sup>-1</sup> and 311.8 and 555.5 mg kg<sup>-1</sup> under soybean-wheat rotation in a Vertisol. Further, the results showed that the continuous application organic manures significantly improved soil KMnO<sub>4</sub>-C content as compared to application of chemical fertilizers, sole biodynamic and absolute control .The organic treatments increased KMnO<sub>4</sub>-C by 72-81% and 64-73% over control and RDF, respectively. Lakaria et al. (2012b) NIR and Jha et al. (2012b) also reported 12% and 34%; 12% and 99% higher KMnO<sub>4</sub>-C under organic farming than conventional farming and control, respectively.

**Table.3 Effect of nutrient management on Labile carbon (mg kg<sup>-1</sup>) under different cropping systems**

Treatment	Soybean-Wheat	Soybean-Mustard	Soybean-Gram	Mean
<b>0-5 cm</b>				
Organic	538	508	499	515
Integrated	479	481	484	481
Inorganic	417	421	466	435
Mean	478	470	483	
LSD (p=0.05)	CS	NM	CS X NM	

	NS	17	30	
	<b>5-15 cm</b>			
Organic	470	440	441	450
Integrated	446	425	423	431
Inorganic	363	404	385	384
Mean	427	423	416	
LSD (p=0.05)	CS	NM	CS X NM	
	NS	14	24	
	<b>15-30 cm</b>			
Organic	391	389	341	374
Integrated	365	338	329	344
Inorganic	342	318	309	323
Mean	366	348	326	
LSD (p=0.05)	CS	NM	CS X NM	
	13	10	17	

---

CS- cropping system, NM – nutrient management

### 3.4 Soil microbial biomass carbon

Soil microbial biomass carbon (SMBC) as influenced by different nutrient management practices under three cropping systems have been presented in Table.4. In 0-5 cm soil depth, the SMBC among the different nutrient management options varied between 224 and 375 mg kg<sup>-1</sup>. Under organic sources of nutrient application the SMBC ranged between 336 and 375 mg kg<sup>-1</sup> with a mean value of 354 mg kg<sup>-1</sup>. With integrated system of nutrient application SMBC ranged between 254 and 312 mg kg<sup>-1</sup> with a mean value of 284 mg kg<sup>-1</sup>. Application of only organic nutrients In 5-15 cm soil depth, the SMBC varied widely where it ranged between 288 and 307 mg kg<sup>-1</sup> with a mean value of 297 mg kg<sup>-1</sup> under organic nutrient management. With integrated nutrient management the SMBC values range was between 244 and 247 mg kg<sup>-1</sup> with a mean value of 245 mg kg<sup>-1</sup>. The lowest SMBC

was recorded with inorganic nutrient sources and it was found to vary from 179 to 191 mg kg<sup>-1</sup> with a mean value of 185 mg kg<sup>-1</sup> (Table 4). In 15-30 cm soil depth, the SMBC decreased sharply irrespective of the nutrient management practices. Under organic sources of nutrient application the SMBC ranged between 69 and 72 mg kg<sup>-1</sup> among the cropping systems with a mean value of 70 mg kg<sup>-1</sup>. The integrated and inorganic nutrient management led to significant decrease in the SMBC. All the cropping systems were at par with respect to SMBC values. Resulted in lowest SMBC (224 to 247 mg kg<sup>-1</sup>) with a mean value of 238 mg kg<sup>-1</sup>. Lakaria et al. (2012c) NIR found that SMBC was ranged from 113 mg kg<sup>-1</sup> to 430.7 mg kg<sup>-1</sup> under different land use whereas Jha et al. (2012b) recorded SMBC in the range of 88.9 to 430.7 mg kg<sup>-1</sup> under different treatments in Vertisols. The results also showed that the SMBC were higher under the treatments receiving the organic source of nutrient application.

**Table 4 Effect of nutrient management on SMBC (mg kg<sup>-1</sup>) under different cropping systems**

Treatment	Soybean-Wheat	Soybean-Mustard	Soybean-Gram	Mean
<b>0-5 cm</b>				
Organic	375	336	351	354
Integrated	312	254	287	284
Inorganic	247	224	244	238
Mean	312	271	294	
LSD (p=0.05)	CS	NM	CS X NM	
	16	17	NS	
<b>5-15 cm</b>				
Organic	307	294	288	297
Integrated	247	244	244	245
Inorganic	191	185	179	185
Mean	249	241	237	
LSD (p=0.05)	CS	NM	CS X NM	

	7	8	NS	
	15-30 cm			
Organic	72	69	69	70
Integrated	56	55	54	55
Inorganic	48	45	44	46
Mean	58	56	56	
LSD (p=0.05)	CS	NM	CS X NM	
	NS	5	NS	

---

CS- cropping system, NM – nutrient management

### 3.5 Dehydrogenase enzymes activity (DHA)

The data on activity of dehydrogenase enzymes in soil under different cropping systems as influenced by nutrient management has been presented in Table 5. In surface soil (0-5 cm) DHA under different nutrient management varied between 124 and 137  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$  under organic sources of nutrients with a mean value of 131.8  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$ . A significant decrease in DHA was recorded with the application of integrated and inorganic source of nutrients. With the integrated use of nutrients the DHA ranged between 104 and 117  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$  with a mean value of 113  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$  among different cropping systems. Inorganic sources of nutrients resulted in lowest dehydrogenase activity (106 and 117) with a mean value of 92  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$ . Highest DHA was recorded under soybean – wheat cropping system followed by soybean – mustard and soybean - gram. In 5-15 cm soil depth, DHA varied between 86 and 89  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$  under organic sources of nutrients while it decreased further under integrated sources of nutrients (64 and 71  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$ ). Application of only inorganic nutrients further decreased DHA (50 and 54  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$ ). All the cropping systems were statistically at par with respect to DHA in 5-15 cm soil depth. DHA in 15-30 cm depth were decreased sharply and different nutrient management treatment showed impact on its activity. Organic sources of nutrients resulted in mean value of 28  $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$ . It was further decreased significantly under integrated and inorganic nutrient management. The soil enzyme activities are positively significantly

correlated with TOC, active, slow and passive pools of carbon, soil respiration, microbial biomass and soil available nitrogen (Casida *et al.*, 1964; NIR; Juma and Tabatabai, 1977; Burns, 1978; and Graham *et al.*, 2005). The organic farming practices are known to increase the microbial enzyme activities in soil as reported by Marinari *et al.* (2006). Results of the present investigation are in accordance with other researchers who also observed higher enzyme activities with the application of FYM (Zaller and Koepke, 2004; Mandal *et al.*, 2007; Bhattacharyya, 2012).

**Table. 5** Effect of nutrient management on dehydrogenase activity (DHA) ( $\mu\text{g TPF g}^{-1} \text{ soil d}^{-1}$ ) under different cropping systems

Treatment	Soybean-Wheat	Soybean-Mustard	Soybean-Gram	Mean
<b>0-5 cm</b>				
Organic	137	134	124	132
Integrated	117	117	104	113
Inorganic	97	90	89	92
Mean	117	114	106	
LSD (p=0.05)	CS	NM	CS X NM	
	8	7	NS	
<b>5-15 cm</b>				
Organic	89	86	88	88
Integrated	70	64	71	69
Inorganic	54	50	51	52
Mean	71	67	70	
LSD (p=0.05)	CS	NM	CS X NM	
	NS	4	NS	
<b>15-30 cm</b>				
Organic	29	27	26	28
Integrated	21	24	21	22

Inorganic	16	15	16	16
Mean	22	22	21	
LSD ( $p=0.05$ )	CS	NM	CS X NM	
	NS	3	NS	

---

CS- cropping system, NM – nutrient management

## Conclusion

Thus, present investigation on long term nutrient management through organic inputs, integrated improved soil aggregation and carbon pools is comparable to inorganic nutrient management. The effect of these nutrient management practices was more pronounced in the surface layer (0-5 to 5-15 cm) as compared to sub the surface layer (15-30cm). The soil organic carbon pool was significantly improved with 100% organic treatment followed by integrated nutrient management and the least being the inorganic fertilizer application. It decreased sharply in the 15-30cm soil layer although treatment effects were the same. In the lower layers, soybean – wheat cropping system maintained significantly higher labile carbon as compared to soybean- mustard and soybean-gram.

## References

- Angeres, D.A. and Caron, J. (1998) Plant-induced soil changes: processes and feedbacks. *Biogeochem.*, 42: 55-72.
- Blair, S.N. (1995) Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men., 12, 273(14):1093-8.
- Burns, R.G. (1978) Enzyme activity in soil: some theoretical and practical considerations. In: Burns, R.G. (ed) Soil enzymes. Academic, New York, pp 295–340.
- Casida, L.E. Jr. (1977) Microbial metabolic activity in soil as measured by dehydrogenase determinations. *Applied Envir. Microbiol.*, 34(6):630-636.
- Graham, M.H., Haynes, R.J., Meyer, J.H. (2005) Soil organic matter content and quality: effects of fertilizer applications, burning and trash retention on a long-term sugarcane experiment in South Africa. *Soil BiolBiochem.*, 34:93-102.

- Gyssels, G., Poesen, J., Bochet, E. and Li, Y. (2005) Impact of plant roots on the resistance of soils to erosion by water: a review. *Progress in Physical Geography: Earth and Envir.*, 29 (2): 189-217.
- Bhattacharyya, T. (2012) Soil dehydrogenase activity in agro-ecological sub regions of black soil regions in India *Geoderma* 197-198:186-192.
- Hassink, J. (1995) Density fractions of soil macroorganic matter and microbial biomass as predictors of C and N mineralization. *Soil Biol. Biochem.*, 27:1099-1108.
- Jarecki, M. and Lal, R. (2003) Crop management for soil carbon sequestration. *Critical review in Pl. Sci.*, 22: 1-32.
- Jenkinson, D.S. and Powlson D.S. (1976) The effects of biocidal treatments on metabolism in soil—V: A method for measuring soil biomass. *Soil Biology and Biochemistry*, 8: 209-213.
- Jha, P., De, A., Lakaria, B. L., Biswas, A. K., Singh, M., Reddy, K. S. and Subba Rao, A. (2012<sub>b</sub>) Soil carbon pools, mineralization and fluxes associated with land use change in Vertisols of central India. *National Acad. Sci. Letters*, 35: 475-483.
- Jha, P., De, A., Lakaria, B.L., Biswas, A.K., Singh, M., Reddy, K.S. and Rao, A.S. ( 2012<sub>a</sub>) Soil Carbon Pools, Mineralization and Fluxes Associated with Land Use Change in Vertisols of Central India. *National Academy Science Letters*, 35(6):475– 483.
- Jha, P., Garg, N., Lakaria, B.L., Biswas, A.K., Subba Rao, A. (2012<sub>b</sub>) Soil and residue carbon mineralization as affected by soil aggregate size. *Soil & Tillage Research*, 121: 57–62.
- Juma, N.G., Tabatabai, M.A. (1977) Effects of trace elements on Dehydrogenase activity in soils. *Soil Sci. Soc. Am. J.*, 41:343–346.
- Leigh, R.A., Johnston, A.E. (1994) *Long-term Experiments in Agricultural and Ecological Sciences*. CAB International, Wallingford, UK.
- Mandal, A., Patra, A.K., Singh, D., Swarup A. and Masto, R.E. (2007) Effect of long-term application of manure and fertilizer on biological and biochemical activities in soil during crop development stages. *Bioresour. Technol.*, 98: 3585-3592.
- Marinari, S., Mancinelli, R., Campiglia, E. and Grego, S. (2006) Chemical and biological indicators of soil quality in organic and conventional farming systems in Central Italy. *Ecol. Indic.*, 6: 701–711.

- McGill, W.B., Cannon, K.R., Robertson, J.A. and Cook, F.D. (1986) Dynamics of soil microbial biomass and water soluble organic C in Breton L after 50 years of cropping to two rotations. *Can. J. Soi. Sci.*, 66:1-19.
- Parmar, D.K., Thakur, D.R., Jamwal, R. S. (2015) Effect of long term organic manure application on soil properties, carbon sequestration, soil - plant carbon stock and productivity under two vegetable production systems in Himachal Pradesh. *Journal of Environ. Biol.*, 37: 333-339.
- Six, J., Paustian, K., Elliott, E. T. and Combrink, C. (2000) Soil organic matter I. Distribution of aggregate-size classes and aggregate-associated carbon. *Soil Science Society of America Journal*, 64:681-689.
- Pohl, M., Alig, D., Körner, C. and Rixen, C. (2009) Higher plant diversity enhances soil stability in disturbed alpine ecosystems. *Pl. Soil*, 324: 91-102.
- Weil, R. R., Islam, K.R., Stine, M. A., Gruver, J. B. and E. Susan, Samson-Liebig (2003) Estimating active carbon for soil quality assessment: A simplified method for laboratory and field use. *American Journal of Alternative Agriculture*, 18: 3-17.
- Walkley, A. and Black, I. A. (1934) An examination of degtjareff method for determination of soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.*,34: 29-38.
- Zaller, J.G. and U. Koepke, (2004) Effect of traditional and biodynamic farmyard manure amendment on yields, soil chemical, biochemical and biological properties in a long-term field experiment. *Biol. Fertil. Soils*, 40: 222-229.