

Revisiting Linear Width: Rethinking the Relationship Between Single Ideal and Linear Obstacle

Abstract:

The study of graph width parameters holds significant importance in the fields of graph theory and combinatorics. Among these parameters, linear-width stands out as a well-established and esteemed measure. The notions of single Ideal and Linear obstacle act as obstacles to achieving optimal linear-width in a connectivity system. In this succinct paper, we offer an alternative proof establishing the equivalence between single ideal and linear obstacle.

Keyword: Linear width, Single ideal, Linear obstacle, Connectivity system

1. Introduction

The exploration of width parameters holds significant importance in the realms of graph theory and combinatorics, as evidenced by the plethora of publications dedicated to this subject (e.g., [4, 5-8, 10-15, 17-31]). Among these parameters, branch-width, a well-studied concept, has received considerable attention in numerous papers. Likewise, linear width, a constrained version of branch-width, has been thoroughly investigated in a multitude of publications. Therefore, the study of both branch-width and linear width assumes critical significance.

The concept of single Ideal, introduced in reference [1], serves as a modeling tool for fundamental mathematical "ideal" in Boolean algebra and topology. In the context of a connectivity system, single Ideal represents the dual concept of linear width (also see reference [2,3]). Additionally, the notion of linear obstacle on a connectivity system corresponds to the dual concept of linear width [6,26].

Building upon these findings, it is established that single Ideal and linear obstacle are equivalent. However, in this concise paper, we present an alternative proof of their equivalence. While the level of novelty may be modest, our objective is to make a valuable contribution to the advancement of research in areas such as Linear width.

2. Definitions in this paper

This section provides mathematical definitions for each concept.

2.1 Symmetric Submodular Function and connectivity system

The definition of a symmetric submodular function is provided below. However, it is important to note that although symmetric submodular functions can generally take real values, this paper specifically focuses on the subset of functions that take only natural numbers.

Definition 1: Let X be a finite set. A function $f: X \rightarrow \mathbb{N}$ is called symmetric submodular if it satisfies the following conditions:

- $\forall A \subseteq X, f(A) = f(X \setminus A)$.
- $\forall A, B \subseteq X, f(A) + f(B) \geq f(A \cap B) + f(A \cup B)$.

A symmetric submodular function possesses the following properties. This lemma will be utilized in the proofs of lemmas and theorems presented in this paper.

Lemma 1 [10]: A symmetric submodular function f satisfies:

1. $\forall A \subseteq X, f(A) \geq f(\emptyset) = f(X)$,
2. $\forall A, B \subseteq X, f(A) + f(B) \geq f(A \setminus B) + f(B \setminus A)$.

In this brief paper, a connectivity system is defined as a pair (X, f) consisting of a finite set (an underlying set) X and a symmetric submodular function f . Throughout this paper, we use the notation f to refer to a symmetric submodular function, a finite set (an underlying set) X , and natural numbers k, m . A set A is said to be k -efficient if $f(A) \leq k$.

2.2. Single Ideal on a connectivity system (X, f)

The definition of a single ideal on a connectivity system (X, f) is given below.

Definition 2 [2]: Let X represent a finite set and f denote a symmetric submodular function. In a connectivity system (X, f) , the set family $S \subseteq 2^X$ is called a single ideal of order $k+1$ if the following axioms hold true:

(IB) For every $A \in S$, $f(A) \leq k$.

(IH) If $A, B \subseteq X$, A is a proper subset of B , and B belongs to S , then A belongs to S .

(SIS) If A belongs to S , $e \in X$, $f(\{e\}) \leq k$, and $f(A \cup \{e\}) \leq k$, then $A \cup \{e\}$ belongs to S .

(IW) X does not belong to S .

In this short paper, we also consider the following additional axiom:

(IE) For each k -efficient subset A of X , exactly one of A or $(X \setminus A)$ is in S .

It has been shown in literature [2] that the linear width of (X, f) is at least $k+1$ if and only if there exists a single ideal on (X, f) of order $k+1$ that satisfies axiom (S4).

2.3 Linear obstacle on a connectivity system (X, f)

The definition of Linear obstacle is shown below. This concept is deep relation to (k, m) -obstacle in literature [6].

Definition 3 [26]: Let X represent a finite set and f denote a symmetric submodular function. In a connectivity system (X, f) , the set family $O \subseteq 2^X$ is called a linear obstacle of order $k+1$ if the following axioms hold true:

(O1) $A \in O$, $f(A) \leq k$,

(O2) $A \subseteq B \subseteq X$, $B \in O$, $f(A) \leq k \Rightarrow A \in O$,

(O3) $A, B, C \subseteq X$, $A \cup B \cup C = X$, $A \cap B = \emptyset$, $f(A) \leq k$, $f(B) \leq k$, $|C| \leq 1 \Rightarrow$ either $A \in O$ or $B \in O$.

3. Result: Equivalence between Single ideal and linear obstacle

The result of this short paper is below.

Theorem 1. Let X represent a finite set and f denote a symmetric submodular function.

Assuming that $f(\{e\}) \leq k$ for every $e \in X$, S is a single ideal of order $k+1$ on (X, f) satisfying the additional axiom (IE) if and only if S is a linear obstacle of order $k+1$ on (X, f) .

Proof of Theorem 1:

(\Rightarrow) If S is a single ideal of order $k+1$ satisfying the additional axiom (IE), then S is a linear obstacle of order $k+1$.

Axiom (O1) is clearly true. Axiom (O2) follows from axiom (IE).

To show axiom (O3), it is clear from axiom (IE) when $|C| = 0$. When $|C| = 1$, it is obvious from axiom (IE) if either $C \subseteq A$ or $C \subseteq B$. Therefore, consider the case where both $C \not\subseteq A$ and $C \not\subseteq B$ hold. Assume, without loss of generality, that $A \notin S$ and $B \notin S$, or $A \in S$ and $B \in S$. Here, we use the fact that either $A \notin S$ or $A \in S$ holds, following from axiom (IE).

When $A \notin S$ and $B \notin S$, we have $(X \setminus A) = B \cup C \in S$. Since $f(B) \leq k$ and $B \subseteq B \cup C$, axiom (IH) implies $B \in S$, leading to a contradiction. When $A \in S$ and $B \in S$, we have $(X \setminus A) = B \cup C \notin S$. On the other hand, from axiom (SIS), we have $B \cup C \in S$, which leads to a contradiction.

(\Leftarrow) If S is a linear obstacle of order $k+1$, then S is a single ideal of order $k+1$ satisfying the additional axiom (IE).

Axiom (IH) and (IB) is obvious.

To show axiom (IE), assume $f(A) \leq k$. Since $A \cup (X \setminus A) = X$ and $A \cap (X \setminus A) = \emptyset$, either $A \in S$ or $(X \setminus A) \in S$ follows from axiom (O3).

To show axiom (SIS), assume $A \in S$ and $f(A \cup \{e\}) \leq k$. Then, we have $f((X \setminus A) \cap (X \setminus \{e\})) = f(A \cup \{e\}) \leq k$, implying $f((X \setminus A) \cap (X \setminus \{e\})) \leq k$. Since $A \in S$ and axiom (O1) hold, we have $f(A) \leq k$. From $A \cup ((X \setminus A) \cap (X \setminus \{e\})) \cup \{e\} = X$ and $A \cap ((X \setminus A) \cap (X \setminus \{e\})) = \emptyset$, either $A \in S$ or $(X \setminus A) \cap (X \setminus \{e\}) \in S$ follows from axiom (O3). Since $A \in S$, we obtain $A \cap (X \setminus \{e\}) \notin S$. Using the previously shown axiom (IE), we obtain $A \cup \{e\} \in S$.

To show axiom (IW), assume $X \in S$, which leads to a contradiction. Using Lemma 1, we obtain $f(X) = f(\emptyset) \leq k$. Using the previously shown axiom (IE) with $A = X$ and $B = \emptyset$, either $X \in S$ or $\emptyset \in S$ follows. Since $X \in S$, we obtain $\emptyset \notin S$, contradicting axiom (IH) which implies $\emptyset \in S$. This proof is completed.

Reference

- [1] Daniel Bienstock. Graph searching, path-width, tree-width and related problems (a survey). Reliability of Computer and Communication Networks, Vol. DIMACS. Series in Discrete Mathematics and Theoretical Computer Science, pp. 33–50, 1989.
- [2] Takaaki Fujita and Koichi Yamazaki. Linear-width and single ideal. The 20th Anniversary of the Japan Conference on Discrete and Computational Geometry, Graphs, and Games, pp. 110–111, 2017.
- [3] Fujita, T. and Yamazaki, K. (2019) Equivalence between Linear tangle and Single Ideal. Open Journal of Discrete Mathematics, 9, 7-10. doi: 10.4236/ojdm.2019.91002.
- [4] Yamazaki, Koichi. "Tangle and maximal ideal." WALCOM: Algorithms and Computation: 11th International Conference and Workshops, WALCOM 2017, Hsinchu, Taiwan, March 29–31, 2017, Proceedings 11. Springer International Publishing, 2017.
- [5] Yamazaki, Koichi. "Inapproximability of rank, clique, boolean, and maximum induced matching-widths under small set expansion hypothesis." Algorithms 11.11 (2018): 173.
- [6] Fedor V Fomin and Dimitrios M Thilikos. On the monotonicity of games generated by symmetric submodular functions. Discrete Applied Mathematics, Vol. 131, No. 2, pp. 323–335, 2003.
- [7] Hicks, Illya V., and Boris Brimkov. "Tangle bases: Revisited." Networks 77.1 (2021): 161-172.
- [8] Hall, Dennis. "A characterization of tangle matroids." Annals of Combinatorics 19 (2015): 125-130.
- [9] Elbracht, Christian, Jakob Kneip, and Maximilian Teegen. "Obtaining trees of tangles from tangle-tree duality." arXiv preprint arXiv:2011.09758 (2020).
- [10] Jim Geelen, Bert Gerards, Neil Robertson, and Geoff Whittle. Obstructions to branch-decomposition of matroids. Journal of Combinatorial Theory, Series B, Vol. 96, No. 4, pp. 560–570, 2006.
- [11] Diestel, Reinhard, Philipp Eberenz, and Joshua Erde. "Duality theorems for blocks and tangles in graphs." SIAM Journal on Discrete Mathematics 31.3 (2017): 1514-1528.
- [12] Paul, Christophe, Evangelos Protopapas, and Dimitrios M. Thilikos. "Graph Parameters, Universal Obstructions, and WQO." arXiv preprint arXiv:2304.03688 (2023).
- [13] Erde, Joshua. "Directed path-decompositions." SIAM Journal on Discrete Mathematics 34.1 (2020): 415-430.
- [14] Diestel, Reinhard. "Ends and tangles." Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg. Vol. 87. No. 2. Berlin/Heidelberg: Springer Berlin Heidelberg, 2017.
- [15] Yamazaki, Koichi, et al. "Isomorphism for graphs of bounded distance width." Algorithmica 24 (1999): 105-127.
- [16] BOŹYK, Łukasz, et al. On objects dual to tree-cut decompositions. Journal of Combinatorial Theory, Series B, 2022, 157: 401-428.
- [17] Diestel, Reinhard, Philipp Eberenz, and Joshua Erde. "Duality theorems for blocks and tangles in graphs." SIAM Journal on Discrete Mathematics 31.3 (2017): 1514-1528.
- [18] Thilikos, Dimitrios M. "Algorithms and obstructions for linear-width and related search parameters." Discrete Applied Mathematics 105.1-3 (2000): 239-271.
- [19] Robertson, Neil, and Paul D. Seymour. "Graph minors. X. Obstructions to tree-decomposition." Journal of Combinatorial Theory, Series B 52.2 (1991): 153-190.
- [20] Robertson, Neil, and Paul D. Seymour. "Graph minors. II. Algorithmic aspects of tree-width." Journal of algorithms 7.3 (1986): 309-322.

- [21] Thilikos, Dimitrios M., and Sebastian Wiederrecht. "Approximating branchwidth on parametric extensions of planarity." *arXiv preprint arXiv:2304.04517* (2023).
- [22] Takaaki Fujita. Reconsideration of Tangle and Ultrafilter using Separation and Partition. *ArXiv*. 2023.
- [23] Oum, Sang-il, and Paul Seymour. "Testing branch-width." *Journal of Combinatorial Theory, Series B* 97.3 (2007): 385-393.
- [24] Hliněný, Petr, and Sang-il Oum. "Finding branch-decompositions and rank-decompositions." *SIAM Journal on Computing* 38.3 (2008): 1012-1032.
- [25] Dieng, Youssou, and Cyril Gavoille. "On the Treewidth of Planar Minor Free Graphs." *Innovations and Interdisciplinary Solutions for Underserved Areas: 4th EAI International Conference, InterSol 2020, Nairobi, Kenya, March 8-9, 2020, Proceedings*. Cham: Springer International Publishing, 2020.
- [26] Fujita, Takaaki. "Alternative Proof of Linear Tangle and Linear Obstacle: An Equivalence Result." *Asian Research Journal of Mathematics* 19.8 (2023): 61-66.
- [27] Fujita, Takaaki. "Proving Maximal Linear Loose Tangle as a Linear Tangle." *Preprints*. (2023).
- [28] Fujita, Takaaki. "Filter for Submodular Partition Function: Connection to Loose Tangle." *Preprints*. (2023).
- [29] Bienstock, Daniel, et al. "Quickly excluding a forest." *J. Comb. Theory, Ser. B* 52.2 (1991): 274-283.
- [30] Fujita, Takaaki. Revisiting the Relationship Between Blockage and Linear Tangle. *Preprints*. Submitted.
- [31] Grohe, Martin. "Tangled up in blue (a survey on connectivity, decompositions, and tangles)." *arXiv preprint arXiv:1605.06704* (2016).