

Original Research Article

Multivariate Analysis in *Triticum dicoccum* species for salinity Tolerance

Formatted: Strikethrough

Abstract

Salinity stands out as a crucial abiotic stress factor that significantly impacts both crop yield and quality. Wheat, considered a moderately salt-tolerant crop, offers a wealth of variability and diversity within its species, presenting an accessible avenue for enhancing wheat's salt tolerance. Consequently, this study investigated the extent of ~~diversity~~ variability in morphological traits, under salt conditions (?) ~~in salt tolerance~~ among *Triticum dicoccum* germplasm accessions. In both saline and control conditions, the genotypes were divided into three and nine separate clusters respectively, showing that there was a significant level of genetic variability among the genotypes by multivariate analysis. It was observed that biomass (~~85.19 %~~) was the largest contributor (85.19 %) to the divergence in the saline situation. Among the yield parameters, tillers per meter followed by thousand-grain weight and spike length were observed to be important. PCA under saline condition, indicated that traits like biomass, number of tillers, plant height, grain yield, days to 50 percent flowering and days to maturity were important traits for selecting salt tolerant genotypes. Thirteen genotypes were salt-tolerant, according to the Stress susceptibility index (SSI). The stress tolerance index (STI) was defined as a useful tool for determining the high yield and potential stress tolerance potential of genotypes. Based on the stress tolerance index, only eight ~~8~~ genotypes were said to be tolerant.

Formatted: Strikethrough

Comment [R11]: How many accessions were evaluated ?

Comment [R12]: unclear objective. objective does not match the title. best in title: analysis of variability in..

Formatted: Strikethrough

Comment [R13]: For what ? in control or salt condition? The traits before named are character components of grain yield..

Formatted: Underline

Formatted: Underline

Formatted: Underline

Formatted: Underline

Formatted: Underline

Formatted: Strikethrough

Keywords: Dicoccum wheat, saline, diversity, principal component and stress index

Introduction

The nutrient-rich cereal wheat is farmed in a wide range of conditions; it covers over 217 million hectares globally, ranking first among all crops in terms of acreage, and produces around 731 million tonnes per year (USDA, 2018). Wheat (*Triticum aestivum* L.) is one of the world's most frequently farmed cereal crops and one of the world's most important staple foods for about 2.5 billion people. Cereals are essential in satisfying the expanding global food demand, especially in underdeveloped nations where cereal-based farming is the only source of nutrients and calories. (Alexandratos and Bruinsma, 2012; Shiferaw *et al.*, 2013). Wheat is the most significant food crop in North Africa, West Asia, and Central Asia, accounting for more than half of the calories consumed. In terms of calories and dietary consumption, wheat, along with rice, is one of the most important sources of protein in the least-developed and middle-income nations. Why study *Triticum dicoccum*? desirable characteristics?

Formatted: Strikethrough

Formatted: Strikethrough

Comment [R14]: How many accessions were evaluated ?

Biological and abiotic stress, often known as environmental stresses, have a significant impact on wheat grain yield (Shamsi *et al.*, 2011; Khavarinejad and Karimov, 2012). Salinity is one of the most important abiotic stresses affecting crop yield and quality (Greenway and Munns, 1980), affecting around 7 percent of the world's total land area (Flowers *et al.*, 1997). Salt affects a higher percentage of cultivated land, with 23 percent of cultivated land being saline and 20 percent of irrigated land suffering from secondary salinization. Furthermore, there is a perilous tendency of a 10 percent annual rise in the saline area all over the world (Ponnamieruma, 1984). In India, around 6.73 million hectares of land are affected by salt, with 3.77 million hectares affected by sodicity and 2.96 million hectares affected by salinity, respectively (Mondal *et al.*, 2010).

Wheat morpho-physiological and agronomic traits play a key role in assessing the value of each trait in enhancing yield, therefore these qualities were employed in breeding projects that resulted in higher yields and the introduction of commercial varieties that can survive salt stress. The selection of genotypes with all conceivable yield contributing traits is the most critical requirement in any crop improvement program. The main factor to consider when making a selection is genotype variability for grain yield and yield component characteristics. Further, such information may be great to set the future path for the salt tolerance breeding program in wheat.

Materials and methods

The present study included pre-tested 60 dicoccum wheat germplasm accessions under both saline (natural saline soils) and control plot, out of which, 46 local germplasm accessions, 4 advanced breeding lines, and ten checks (Table-1) were evaluated in alpha lattice design with four blocks and two replications. Each block consisted of 15 genotypes with two rows per genotype and 3- meter length with a spacing of 20 cm between rows. The present investigation was carried out during *rabi* 2020-21 at Ugar Sugars Pvt. Ltd, Ugar Khurd, Tq- Chikkodi, Dt-Belagavi, Karnataka, which is situated in the northern transitional tract of Karnataka with 16°38' N latitude and 74° 49' E longitude at an altitude of 537 m above mean sea level (AMSL). Two independent experiments were carried out in both salt stress and controlled condition. Under saline condition, pH of less than 8 and EC of more than 4 dS/m and under control condition pH of 6-8 and EC of less than 4 dS/m was maintained. Table 2 represents the pH and EC dS/m (1 soil:2.5 water) of top layer (0-20 cm) and bottom layer (20-40 cm) of the control and saline field.

Morphological traits like germination percent, days to fifty percent flowering, days to maturity and plant height, physiological traits (SPAD and NDVI) at booting, anthesis and grain filling stages, yield and yield attributes *viz.*, number of productive tillers per meter row, spike length (cm), number of grains per spike, number of spikelets per spike, harvest index, biomass and thousand-grain weight (g) were measured under the study. The statistical analysis for genetic divergence was done using Mahalanobis-D² statistics (Mahalanobis, 1936) and the clustering of genotypes was done using the Tocher method (Rao, 1952). The character contribution towards genetic divergence was computed using the method given by Singh and Chaudhary (1977). Principle component analysis was done

Comment [R15]: ANOVA was used ? univariate and multivariate analyses? test of comparison of averages?

using the method suggested by Kovacic (1994). Stress indices viz., Stress Susceptibility Index (Fischer and Maurer, 1978) and Stress Tolerance Index (Fernandez, 1992) were calculated to classify genotypes into salt tolerant and salt susceptible.

Comment [R16]: How were calculated the stress indices?

Table 1: List of genotypes used

Germplasm lines			Advanced breeding lines
DIC-1	DIC-39	DIC-74	DDK 50033
DIC-4	DIC-43	DIC-76	DDK 50505
DIC-9	DIC-44	DIC-77	DDK 50444
DIC-12	DIC-45	DIC-83	DDK 50507
DIC-13	DIC-46	DIC-88	
DIC-14	DIC-47	DIC-91	Checks
DIC-15	DIC-48	DIC-92	Kharchia 65 (Bread wheat, only salt tolerant donor)
DIC-17	DIC-49	DIC-93	KRL 99 (Bread wheat, Salt tolerant released variety, CCSRI, Karnal)
DIC-18	DIC-50	DIC-101	KRL 3-4 (Bread wheat, Salt tolerant released variety, CCSRI, Karnal)
DIC-19	DIC-68	DIC-102	IC 0408331 (Bread wheat, salt tolerant genetic stock, NBPGR)
DIC-21	DIC-70	DIC-103	
DIC-22	DIC-71	DIC-104	
DIC-23	DIC-72	DIC-105	DDK 1029 (Dicocum wheat, high yielding variety)
DIC-26	DIC-73	DIC-106	
DIC-107	DIC-94	DIC-95	UAS 334 (Bread wheat, high yielding variety)

Table 2: Soil pH and Electrical conductivity of saline and control plot at different stages of crop

Crop stage	Soil layer	Saline		Control	
		pH	EC	pH	EC
Sowing	Top layer (0-20 cm)	8.01	6.56	8.02	2.89
	Bottom layer (20-40 cm)	7.98	6.51	8.01	3.01
Booting	Top layer (0-20 cm)	7.8	6.11	7.7	2.21
	Bottom layer (20-40 cm)	7.89	5.89	7.86	2.31
Grain filling	Top layer (0-20 cm)	8.02	6.02	7.95	3.32
	Bottom layer (20-40 cm)	7.77	6.11	7.99	3.11
Harvesting	Top layer (0-20 cm)	7.81	6.31	8.03	3.1
	Bottom layer (20-40 cm)	7.62	6.29	7.56	2.98

Results and discussion

Salinity has affected all the traits under study, and grain yield has reduced significantly, statically significant?. Traits like spikelets per spike, days to maturity, and days to fifty percent flowering are least affected by salt stress (Table 3). All the traits showed a significant difference for all the traits under observation under both saline and control condition. Salt-tolerant wheat varieties are considered to be a viable and effective strategy for reducing the harmful effects of salt stress (Ashraf and O'Leary, 1996; Flowers and Yeo, 1995; Singh and Rana, 1985). Many efforts have been made in the past to develop salinity-tolerant wheat genotypes. Using traditional breeding procedures, successful results have been reached in the form of varieties such as KRL 19, KRL 1-4, KRL 210, and others (Shahbaz and Ashraf, 2013), no such efforts are made to breed

Comment [R17]:

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Table 3: ANOVA for alpha lattice design for different traits under saline and control condition

Sl.no	Trait	Saline					Control				
		Mean	CV	SE	SS	MSS	Mean	CV	SE	SS	MSS
1	GY	19.88	7.57	10.65	300780.77	5013.01	36.23	0.54	13.78	909783.54	15163.05
2	TPM	93.81	2.21	1.47	126717.16	2111.95	122.02	2.00	1.72	78828.97	1313.82
3	BM	84.01	0.25	14.56	8196229.52	136603.82	114.24	13.17	10.64	11144858.62	185747.64
4	SPS	19.43	4.81	0.66	6589.48	109.82	20.27	5.48	0.79	10811.57	180.19
5	SL	6.63	1.42	0.07	144.67	2.41	8.58	4.77	0.29	195.92	3.27
6	GPS	29.28	6.20	1.28	3666.52	61.11	34.23	5.71	1.38	5732.57	95.54
7	PH	81.35	1.79	1.03	15328.41	255.47	94.79	3.47	2.32	8652.79	144.21
8	GP	92.89	4.21	2.76	1307.11	21.79	94.44	4.02	2.68	826.10	13.77
9	TGW	29.92	1.52	0.32	2456.77	40.95	35.42	1.58	0.40	724.39	12.07
10	DFF	60.01	0.81	0.35	3500.49	58.34	62.15	0.63	0.28	3426.25	57.10
11	DM	107.37	0.37	0.28	4329.60	72.16	109.52	0.37	0.28	4360.24	72.67
12	HI	25.25	7.97	1.42	10426.69	173.78	33.38	15.95	3.76	9622.01	160.37
13	NDVI-I	0.53	3.10	0.01	1.23	0.02	0.65	13.92	0.06	0.84	0.01
14	NDVI-II	0.59	3.19	0.01	1.30	0.02	0.65	14.30	0.07	0.79	0.01
15	NDVI-III	0.62	17.95	0.08	1.04	0.02	0.60	19.03	0.08	0.98	0.02
16	SPAD-I	43.50	13.15	4.04	1818.11	30.30	43.88	11.61	3.60	2203.52	36.73
17	SPAD-II	37.71	12.31	3.28	1646.48	27.44	41.23	10.46	3.05	1905.92	31.77
18	SPAD-III	36.69	19.55	5.07	1823.46	30.39	41.48	12.29	3.60	2128.23	35.47

GY- Grain yield (q/ha), TPM-Tillers per meter, BM- Biomass (q/ha), SPS- Spikelets per spike, SL-Spike length (cm), GPS-Grains per spike, PH-Plant height (cm), GP- Germination percentage (per cent), TGW-Thousand grain weight (g), DFF- Days to 50 per cent flowering, DM-Days to maturity, HI-Harvest index, NDVI- anthesis stagebooting and grain filling stages, SPAD- Chlorophyll content before anthesis, booting and grain filling stages. [Test means comparisons](#)

?

for a salt-tolerant dicoccum wheat genotype or to identify a salt tolerant dicoccum wheat genotype. However, a significant increase in wheat grain production under salinity stress has yet to be achieved.

The sum of squares of the difference between the mean values obtained between any two genotypes for all morpho-physiological, yield, and yield attributing traits in sixty wheat genotypes under two environments *viz.*, saline and control condition were used for the final grouping of the genotypes. The genetic diversity analysis was carried out using Tocher's (Rao, 1952) approach, which was used to arrange 60 wheat genotypes into various clusters by considering the estimated D^2 value as the square of the generalized distance. The pattern of distribution of genotypes into various clusters under saline and control condition was presented in Tables 4 and 5 respectively. Under saline condition, three clusters were created by combining all 60 wheat genotypes in such a way that the genotypes in each cluster had a lower D^2 value than the genotypes in the other clusters. Cluster I had the most genotypes *i.e.*, 58 genotypes, followed by Cluster II and III the ones that were solitary with genotypes DIC 18 and DIC 94 respectively. Nine clusters were formed under control condition, cluster I was the largest with 16 genotypes, followed by cluster III and cluster V with 12 genotypes each, cluster II had 7 genotypes which included the checks. Cluster IV had six genotypes and Cluster VI had four genotypes, cluster VII, VIII, IX were solitary clusters with genotypes KRL 210, UAS 334 and DIC 103 respectively.

The divergence analysis showed that all sixty genotypes were grouped into three clusters in saline condition and seven clusters, under control condition. A maximum number of genotypes were grouped into cluster I under both saline and control condition. Genotypes namely DIC 18 of cluster II and DIC 94 of cluster III in saline environment were showing high inter-cluster distance. Hybridization between the genotypes of these clusters might yield desirable segregants (Table 6). These results are supported by those of Mundiyyara *et al.* (2013), Vinod *et al.* (2014), Singh *et al.* (2015), Ibrahim and Arafa (2020) and Srivijay *et al.* (2022)

The highest intra-cluster distance was obtained by cluster I (167.61), and between clusters II and III highest inter-cluster distance (863.27) was obtained under saline condition, followed by clusters I and III (589.92), represented in Table 6. Under non-stress condition (Table 7), cluster IV (44.50) showed the highest intra-cluster distance, clusters IX and VII were showing the highest inter-cluster (222.80) distance, followed by cluster IX and cluster II (211.02). The cluster means for all the characters under saline conditions are represented in Table 8. The solitary cluster III had early flowering having the genotype DIC-94 (55 days), high grain yield (2,735 kg/ha) and the highest mean values for biomass (18,977 kg/ha), number of tillers per meter (157.00), spike length (11.41cm), germination percent (96.50 percent), plant height (104.73 cm), NDVI at anthesis stage (0.70), lowest NDVI at booting stage (0.37), chlorophyll content at anthesis stage (34.68), number of spikelets per spike (15.00) and least harvest index of 14.40 percent were recorded by this cluster. Under control condition (Table 9), cluster IX with genotype DIC 103 had the highest grain yield (5428.62 kg/ha), biomass (12965.64

kg/ha), late flowering (69 days), late maturing (118 days), highest harvest index (42.06 %) and highest NDVI at grain filling

UNDER PEER REVIEW

Table 4: Clustering of wheat genotypes based on morpho-physiological, yield and yield attributing traits under saline condition

Cluster	Number of genotypes	Genotypes
Cluster I	1	DIC-1, DIC-4, DIC-9, DIC-12, DIC-13, DIC-14, DIC-15, DIC-17, DIC-19, DIC-21, DIC-22, DIC-23, DIC-26, DIC-39, DIC-43, DIC-44, DIC-45, DIC-46, DIC-47, DIC-48, DIC-49, DIC-50, DIC-68, DIC-70, DIC-71, DIC-72, DIC-73, DIC-74, DIC-76, DIC-77, DIC-83, DIC-88, DIC-91, DIC-92, DIC-93, DIC-95, DIC-99, DIC-101, DIC-102, DIC-103, DIC-104, DIC-105, DIC-106, DIC-107, DDK 50033, DDK 50505, DDK 50444, DDK 50507, KRL 210, KRL 19, UAS 304, UAS 334, UAS 428, DDK 1029, HD 2009, Kharchia 65, KRL 3-4, IC 0408331
Cluster II	1	DIC-18
Cluster III	1	DIC-94

Table 5: Clustering of wheat genotypes based on morpho-physiological, yield and yield attributing traits under control condition

Cluster	Number of genotypes	Genotypes
Cluster I	16	DIC-1, DIC-4, DIC-18, DIC-23, DIC-26, DIC-12, DIC-15, DIC-50, DIC-72, DIC-91, DIC-46, DIC-47, DIC-76, DDK 50033, DIC-83, DIC-93
Cluster II	7	KRL 3-4, IC 0408331, UAS 304, HD 2009, Kharchia 65, KRL 19, UAS 428
Cluster III	12	DIC-13, DIC-19, DIC-22, DIC-44, DIC-68, DIC-70, DIC-74, DIC-95, DIC-99, DIC-102, DDK 50505, DDK 1029
Cluster IV	6	DIC-14, DIC-43, DIC-94, DIC-39, DIC-17, DIC-71
Cluster V	12	DIC-2, DIC-45, DIC-48, DIC-49, DIC-77, DIC-92, DIC-101, DIC-104, DIC-105, DIC-106, DIC-107, DDK 50507
Cluster VI	4	DIC-73, DIC-88, DDK 50444, DIC-9
Cluster VII	1	KRL 210
Cluster VIII	1	UAS 334
Cluster IX	1	DIC-103

Table 6: Intra and inter-cluster D^2 values in wheat genotypes under saline condition

	Cluster I	Cluster II	Cluster III
Cluster I	167.61		
Cluster II	313.33	0.00	
Cluster III	589.92	863.27	0.00

Table 7: Intra and inter-cluster D^2 values in wheat genotypes under control condition

	Cluster I	Cluster II	Cluster III	Cluster IV	Cluster V	Cluster VI	Cluster VII	Cluster VIII	Cluster IX
Cluster I	33.46								
Cluster II	121.21	36.88							
Cluster III	62.88	78.11	37.87						
Cluster IV	53.79	106.30	64.74	44.18					
Cluster V	67.63	167.30	113.63	81.41	44.50				
Cluster VI	100.58	62.88	56.75	94.01	153.03	40.96			
Cluster VII	136.01	51.14	90.81	130.53	185.17	75.55	0.00		
Cluster VIII	71.58	79.80	58.08	76.65	111.12	86.55	83.88	0.00	
Cluster IX	99.13	211.02	150.29	125.55	64.01	191.64	222.80	146.63	0.00

Table 8: Cluster means for various quantitative traits in wheat genotypes under saline situation

	GY	TPM	BM	SPS	SL	GPS	PH	GP	TGW
Cluster I	1969.64	91.88	8318.46	19.54	6.53	29.15	80.78	92.77	29.89
Cluster II	2351.50	144.50	2732.50	17.50	7.88	33.50	91.30	96.50	30.86
Cluster III	2735.00	157.00	18977.00	15.00	11.41	32.50	104.72	96.50	30.94

	DFF	DM	HI	NDVI-I	NDVI-II	NDVI-III	SPAD-I	SPAD-II	SPAD-III
Cluster I	60.06	107.44	25.17	0.53	0.59	0.62	43.57	37.80	36.61
Cluster II	62.00	106.64	41.03	0.65	0.65	0.70	35.01	35.70	36.92
Cluster III	55.00	104.01	14.41	0.37	0.70	0.61	47.59	34.67	41.15

GY- Grain yield (q/ha), TPM-Tillers per meter, BM- Biomass (q/ha), SPS- Spikelets per spike, SL-Spike length (cm), GPS-Grains per spike, PH-Plant height (cm), GP- Germination percentage (per cent), TGW-Thousand grain weight (g), DFF- Days to 50 per cent flowering, DM-Days to maturity, HI-Harvest index, NDVI- anthesis stage booting and grain filling stages, SPAD- Chlorophyll content before anthesis, booting and grain filling stages.

Table 9: Cluster means for various quantitative traits in wheat genotypes under control situation

	GY	TPM	BM	SPS	SL	GPS	PH	GP	TGW
Cluster I	3940.76	117.06	12199.62	15.91	7.89	31.56	96.47	94.38	36.32
Cluster II	2480.07	112.56	9990.45	35.56	9.74	34.63	86.52	93.94	35.30
Cluster III	3138.67	116.42	11040.57	18.00	8.51	33.46	94.82	94.46	34.79
Cluster IV	3653.37	157.08	12281.30	19.33	9.08	39.75	99.50	94.08	35.89
Cluster V	4786.99	131.04	12743.90	17.50	8.66	36.38	100.49	95.08	35.55
Cluster VI	2383.67	125.63	7994.25	13.63	7.52	27.75	89.38	94.63	34.22
Cluster VII	2429.39	81.00	6599.81	51.00	11.08	49.50	76.33	92.50	31.33
Cluster VIII	3727.00	74.50	11160.66	33.00	7.79	31.50	78.00	94.50	34.54
Cluster IX	5428.62	99.50	12965.64	17.00	9.95	37.50	94.17	95.00	34.52

	DFF	DM	HI	NDVI-I	NDVI-II	NDVI-III	SPAD-I	SPAD-II	SPAD-III
Cluster I	65.62	113.31	35.38	0.67	0.67	0.61	43.79	40.95	40.51
Cluster II	52.06	98.75	25.79	0.59	0.60	0.57	49.02	44.69	45.44
Cluster III	63.13	110.62	30.35	0.68	0.68	0.58	42.90	40.54	41.24
Cluster IV	61.32	108.96	32.01	0.58	0.62	0.56	40.90	38.01	41.18
Cluster V	63.91	111.14	39.18	0.63	0.61	0.62	42.87	40.68	41.38
Cluster VI	64.36	112.94	30.88	0.63	0.63	0.67	42.95	39.93	37.76
Cluster VII	52.84	95.20	37.95	0.72	0.71	0.60	44.46	42.75	42.22
Cluster VIII	52.98	97.62	33.98	0.74	0.74	0.63	46.16	53.00	42.35
Cluster IX	68.92	118.04	42.06	0.70	0.68	0.74	46.99	44.45	44.13

GY- Grain yield (q/ha), TPM-Tillers per meter, BM- Biomass (q/ha), SPS- Spikelets per spike, SL-Spike length (cm), GPS-Grains per spike, PH-Plant height (cm), GP- Germination percentage (per cent), TGW-Thousand grain weight (g), DFF- Days to 50 per cent flowering, DM-Days to maturity, HI-Harvest index, NDVI- anthesis stage booting and grain filling stages, SPAD- Chlorophyll content before anthesis, booting and grain filling stages.

stage (0.74). Cluster VII with genotype KRL 210 was showing the highest mean values for number of spikelets per spike (51), spike length (11.08 cm) and highest grains per spike (49.50).

The percent contribution of each character towards total genetic divergence under saline and control condition is presented in Table 10. It was observed that under saline condition, biomass (85.19 %) was the largest contributor to the divergence. Days to maturity (6.28 %), tillers per meter row (4.32 %) ranking III and IV contributed the highest to the diversity among various yield attributes. On the contrary, the contribution of other yield attributes such as thousand-grain weight (1.69 %), spikelets per spike (0.33 %), spike length (0.93 %) and grains per spike (0.05 %) was less. Among the morphological traits, days to flowering 0.27 percent to diversity. The NDVI reading at the booting stage (0.16 %) and anthesis (0.11) were observed to be important. Under control condition, it was observed that among various groups of traits in general, the contribution of grain weight (66.12 %) was found highest, followed by NDVI at the anthesis stage (12.24 %), days to maturity (10.49 %), tillers per meter (5.14 %), days to 50 percent flowering (3.39 %) and spikelets per spike (1.97 %) respectively.

The utility of the D^2 statistic is enhanced by its applicability to estimate the relative contribution of various characters to the total genetic divergence. The feature that contributes the most to divergence is utilized to place a higher emphasis on selecting a cluster for further selection and selecting parents for hybridization. It has been noted that under saline condition, biomass (85.19 %) was the largest contributor to the divergence. This suggests that genotype DIC-94 belonging to cluster III exhibiting a high mean for biomass, grain yield, tillers per meter, spike length, thousand-grain weight, plant height, and germination percent can be a potential source for hybridization program to obtain transgressive segregants. These results are in concordance with Sunilyadav *et al.* (2014) wherein they observed a higher contribution of thousand-grain weight, tillers per meter, days to maturity, days to flowering, and grain yield towards diversity.

Under the saline condition (Table 11), only 7 principal components were having Eigen values greater than one (>1) contributing 80.49 percent of the total cumulative percentage with component-1 having the maximum percentage of 25.97 percent, followed by 14.63 percent contribution by component 2, followed by 10.25 percent (PC3), 9.03 percent (PC4), 7.69 percent (PC5), 6.89 percent (PC6) and 6.02 percent (PC7). All measured traits were positively loaded onto PC1 and PC2.

Principal component analysis was constructed to identify superior genotypes for a stressed environment. Principal component analysis simplifies complex data by transforming the number of correlated variables into a smaller number of variables called principal components. The first principal component accounts for maximum variability in the data with respect to succeeding components (Leilah and Al-Khateeb, 2005). The distribution of genotypes with the contribution of characters towards genetic divergence (Biplot). Therefore, a perusal of PCA under saline condition (Fig 1&2), indicated that traits like biomass, number of tillers, thousand-grain weight, plant height, grain yield, days to 50 percent flowering and days to maturity were important traits for selecting salt tolerant

genotypes. Similar results were obtained by Al-Ashkar *et al.* (2019), Dadshani *et al.* (2019) and Singh *et al.* (2015) for traits like the number of tillers, number of grains, days to maturity, grain yield and harvest index.

Table 10: Per cent contribution of morpho-physiological traits, yield and yield attributes to total genetic diversity under control and saline condition

Sl. No.	Source	Per cent contribution	
		Saline condition	Control condition
1	GY		66.12
2	TPM	4.32	5.14
3	BM	85.19	0.05
4	SPS	0.33	1.97
5	SL	0.93	
6	GPS	0.05	0.11
7	PH	0.66	
8	GP		
9	TGW	1.69	0.38
10	DFF	0.27	3.39
11	DM	6.28	10.49
12	HI		
13	NDVI-I	0.16	
14	NDVI-II	0.11	12.24
15	NDVI-III		
16	SPAD-I		
17	SPAD-II		0.11
18	SPAD-III		

Table 11: Eigen values for principal components under saline and control condition

Component	Total	Saline Eigenvalues	
		Per cent variance	Cumulative per cent
1	4.68	25.97	25.97
2	2.63	14.63	40.60
3	1.85	10.25	50.85
4	1.63	9.03	59.89
5	1.38	7.69	67.57
6	1.24	6.89	74.46
7	1.08	6.02	80.49
8	0.85	4.73	85.22
9	0.74	3.95	89.17
10	0.57	3.18	92.35
11	0.48	2.68	95.03
12	0.44	2.28	97.30
13	0.17	0.96	98.27
14	0.15	0.81	99.08
15	0.11	0.61	99.69
16	0.04	0.20	99.89
17	0.02	0.09	99.98
18	0.00	0.02	100.00

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough




Fig1: PCA biplot undersaline condition




Fig 2: PCA biplot for variables undersaline condition

The majority of wheat genotypes are vulnerable to salt stress, resulting in low and unpredictable yields. Because current screening approaches for salinity-related traits are time-consuming and impracticable for most researchers, it is critical to estimate genotype yield under saline and control condition, and tolerance indices are currently being used to quantify the effect of salt on yield. These salt tolerance indices indicate whether the genotypes are salt-tolerant or susceptible to the saline condition. To investigate suitable stress resistance indices for screening of genotypes under salt condition, the grain yield of cultivars under both non-stress and stress condition were measured for calculating different sensitivity and tolerance indices. Among them, stress susceptibility (Table 12) and stress tolerance index (Table 13) have been considered for grouping of genotypes. These indices categorized the sixty wheat genotypes into three categories viz., tolerant, moderately tolerant, and susceptible. The tolerant category of SSI included 14 genotypes. Among 13 tolerant genotypes, eight dicoccum wheat genotypes (DIC-9, DIC-13, DIC-68, DIC-73, DIC-88, DIC-94, DIC-95, DDK 50505), along with five checks viz., KRL 210, UAS 334, Kharchia 65, KRL 3-4, and IC 0408331. Based on the stress tolerance index (STI), only 10 genotypes viz., DIC-26, DIC-48, DIC-92, DIC-101, DIC-104, DIC-107, DDK 50507 and UAS 334 were said to be tolerant.

The stress susceptibility index was evaluated based on the yield ratio of each variety in stressed conditions to non-stressed conditions as compared with the proportions in total varieties. Thus, tolerant cultivars identified with high yield or low in both conditions can have the same amount of SSI, so the selection process based on this index led reformers to make a mistake (Naeimi *et al.*, 2008). So, according to the stress-sensitive index, it was observed that 13 genotypes viz., eight dicoccum wheat (DIC-9, DIC-13, DIC-68, DIC-73, DIC-88, DIC-94, DIC-95, DDK 50505) and five checks (KRL 210, UAS 334, Kharchia 65, KRL 3-4, IC 0408331) as a tolerant. A similar classification of genotypes using SSI was done for 97 bread-wheat genotypes by Sardouie *et al.* (2014). The stress tolerance index (STI) was defined as a useful tool for determining the high yield and stress tolerance potential of genotypes (Fernandez, 1992). Based on the stress tolerance index, only 8 genotypes were identified as salt tolerant viz., seven dicoccum wheat genotypes (DIC-26, DIC-48, DIC-92, DIC-101, DIC-104, DIC-107, DDK 50507) and check UAS 334. Mevlut and Sait (2011) indicated that the genotypes with >0.9 STI value, usually have high-yielding potential under stress condition even greater differences in two different conditions. Hence, there is still scope for both the indices and generating super salt tolerant varieties.

With respect to dicoccum wheat genotype (DIC-88) based on SSI and few genotypes (DIC-101, DIC-92, DIC-48, DIC-26, DDK 50507) based on STI accept the genotype DIC 101 which is the most promising considering both SSI and STI can be involved in biparental mating to complement the mechanisms operating in both categories to improve both salt tolerance and yield simultaneously. It is clear from the present study that there is a lot of scope for improvement of all these genotypes using the potential donors identified. Genome 'A' has the next highest genes or QTLs followed by 'D' genome of hexaploid wheat. Mechanisms like tissue level compartmentalization *i.e.*, accumulation of toxic ions in the lower leaves of the plant is noted in dicoccum wheat. Cell level compartmentalization in the cell vacuoles was noted by Munns and Tester (2008).

Formatted: Underline

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Underline

Table 12: Grouping of wheat genotypes based on Stress susceptibility index (SSI)

Range of SSI	<u>Drought tolerance category</u>	Number of genotypes	Genotypes
<0.5	<u>Tolerant?</u>	13	Dicoccum wheat (8): DIC-9, DIC-13, DIC-68, DIC-73, DIC-88, DIC-94, DIC-95, DDK 50505 Check (5): KRL 210, UAS 334, Kharchia 65, KRL 3-4, IC 0408331
0.5 to 1.0	Moderate	26	Dicoccum wheat (24): DIC-12, DIC-14, DIC-15, DIC-17, DIC-18, DIC-19, DIC-22, DIC-23, DIC-26, DIC-39, DIC-44, DIC-47, DIC-48, DIC-50, DIC-70, DIC-71, DIC-74, DIC-76, DIC-91, DIC-99, DIC-101, DIC-102, DIC-107, DDK 50444 Check (2): KRL 19, UAS 304
>1.0	Susceptible	21	Dicoccum wheat (18): DIC-1, DIC-4, DIC-21, DIC-43, DIC-45, DIC-46, DIC-49, DIC-72, DIC-77, DIC-83, DIC-92, DIC-93, DIC-103, DIC-104, DIC-105, DIC-106, DDK 50033, DDK 50507 Check (3): UAS 428, DDK 1029, HD 2009

Formatted: Underline, Highlight

Formatted: Underline

Formatted: Highlight

Table 13: Grouping of wheat genotypes based on stress tolerance index (STI)

Range of STI	<u>Drought tolerance category</u>	Number of genotypes	Genotypes
>0.9%	Tolerant	8	Dicoccum wheat (7): DIC-26, DIC-48, DIC-92, DIC-101, DIC-104, DIC-107, DDK 50507 Check: UAS 334
0.8-0.9%	Moderate	11	Dicoccum wheat (10): DIC-17, DIC-18, DIC-21, DIC-45, DIC-49, DIC-71, DIC-77, DIC-91, DIC-94, DIC-105 Check: KRL 210
<0.8%	Susceptible	41	Dicoccum wheat (33): DIC-1, DIC-4, DIC-9, DIC-12, DIC-13, DIC-14, DIC-15, DIC-19, DIC-22, DIC-23, DIC-39, DIC-43, DIC-44, DIC-46, DIC-47, DIC-50, DIC-68, DIC-70, DIC-72, DIC-73, DIC-74, DIC-76, DIC-83, DIC-88, DIC-93, DIC-95, DIC-99, DIC-102, DIC-103, DIC-106, DDK 50033, DDK 50505, DDK 50444 Check (9): KRL 19, UAS 304, UAS 428, DDK 1029, HD 2009, Kharchia 65, KRL 3-4, IC 0408331

Formatted: Highlight

Conclusion

The study's findings indicate that dicoccum wheat germplasms possess significant potential as sources of salt tolerance. Additionally, a handful of promising accessions could be officially recognized as national genetic resources or developed into new wheat varieties. This research underscores the untapped possibilities offered by dicoccum wheat species in addressing global challenges related to hunger and growing populations, particularly in regions characterized by unproductive, saline soils.

Formatted: Strikethrough

Formatted: Underline

References

- Al-Ashkar I, Alderfasi A, El-Hendawy S, Al-Suhaibani N, El-Kafafi S, Seleiman MF. Detecting salt tolerance in doubled haploid wheat lines. *Agronomy*. 2019;9(4):211.
- Alexandratos N, Bruinsma J. World agriculture towards 2030/2050: the 2012 revision.
- Ashraf M, O'Leary JW. Responses of some newly developed salt-tolerant genotypes of spring wheat to salt stress. Yield components and ion distribution. *J. of Agron. and Crop Sci.* 1996;176: 91–101.
- Dadshani S, Sharma RC, Baum M, Ogbonnaya FC, Léon J, Ballvora A. Multi-dimensional evaluation of response to salt stress in wheat. *PLoS one*. 2019;14(9):0222659.
- Fernandez GCJ., Effective selection criteria for assessing plant stress tolerance. In: Kuo CG (ed) *Proceedings of the International Symposium on Adaptation of Vegetables and other Food Crops in Temperature and Water Stress*, Publication, Tainan, Taiwan. 1992;257-270.
- Fischer R A, Maurer R. Drought resistance in spring wheat cultivars. I. Grain responses. *Australian J. Agric. Res.* 1978;29:897–912.
- Flowers TJ, Garcia A, Koyama M, Yeo AR. Breeding for salt tolerance in crop plants, the role of molecular biology. *Acta Physiol. Plant.* 1997;19(4):427-433.
- Flowers TJ, Yeo AR. Breeding for salinity resistance in plants. *Aust. J. Plant Physiol.* 1995;22: 875-884.
- Greenway H, Munns. Mechanism of salt tolerance in non-halophytes. *Annu. Rev. Plant. Physiol.* 1980;31:146-190.
- Ibrahim HM, Arafa SA. Effect of Chitosan on Growth, Yield and Certain Salinity Stress-Related Metabolites in Two Barley Cultivars Contrasting in Salt Tolerance. *J. Plant Prod. Sci.* 2020;11(10):899-906.

- Khavarinejad MS, Karimov M. Study of Genetic diversity among spring wheat genotypes in drought stress by advanced statistical analysis. *Int. J. Agron. Plant Prod.* 2012;3(12):590-598.
- Kovacic Z. Multivariate analysis. Faculty of Economics. University of Belgrade, Serbia. 1994:293.
- Leilah AA, Al-Khateeb SA. Statistical analysis of wheat yield under drought conditions. Elsevier. 2005;61:483-496.
- Mahalanobis PC. on the generalised distance in statistics. *Proceed. Nat. Instit. Sci.* 1936;2:49-55.
- Mevlut A, Sait C. Evaluation of drought tolerance indices for selection of Turkish oat (*Avena sativa* L.) landraces under various environmental conditions. *Zemdirbyste Agriculture.* 2011;98(2):157-166.
- Mondal AK, Sharma RC, Singh G, Dagar JC. Computerised database on salt affected soils in India. *Tech. Bull.* 2010;2/2010. CSSRI, Karnal.
- Mundiyyara R, Kerkhi SA, Jakhar ML, Ganga R. D2 analysis in bread wheat (*Triticum aestivum* L.). *J. Plant Sci. Res.* 2013;29(2):215-219.
- Munns R, Tester M. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* 2008;59:651-681
- Naeimi M, Akbari GA, Shiranirad AH, Modares SS, Sadat NS, Jabari H. Evaluation of drought tolerance in different canola cultivars based on stress evaluation indices in terminal growth duration. 2008;83-98.
- Ponnamieruma PN. Role of cultivars tolerance in increasing rice production on saline land. In R. C. Staples & G. H. Toenniessen (Eds.), *Salinity tolerance in plants strategies for crop improvement.* Wiley, New York. 1984;255-71.
- Rao CR. *Advanced Statistical Methods in Biometrical Research.* John Wiley and Sons, Inc., New York. 1952;357-363.
- SardouieNasab S, MohammadiNejad G, Nakhoda B. Field screening of salinity tolerance in Iranian bread wheat lines. *Crop Sci.* 2014;54 (4):1489-1496.
- Shahbaz M, Ashraf M. Improving salinity tolerance in cereals. *Crit. Rev. Plant Sci.* 2013;32(4):237-249.
- Shamsi K, Petrosyan M, Noor-mohammadi G, Haghparast A, Kobraee S, Rasekhi B. Differential agronomic responses of bread wheat cultivars to drought stress in the west of Iran. *Afr. J. Biotechnol.* 2011;10(14):2708-2715.
- Shiferaw B, Smale M, Braun HJ, Duveiller E, Reynolds M, Muricho G. Crops that feed the world 10. Past successes and future challenges to the role played by wheat in global food security. *Food Secur.* 2013;5(3):291-317.

- Singh KN, Rana RS. Genetic variability and character association in wheat varieties grown on sodic soil. Indian J. of Agri Sci. 1985;55:723-726.
- Singh R K, Chaudhary BD. Biometrical methods in quantitative genetic analysis, Kalyani Publishers, New Delhi. pp. 1977;57-58.
- Singh S, Sengar RS, Kulshreshtha N, Datta D, Tomar RS, Rao VP, Garg D, Ojha A. Assessment of multiple tolerance indices for salinity stress in bread wheat (*Triticum aestivum* L.). J. Agric. Sci. 2015;7(3):49.
- Srivijay Malipatil, SS Biradar, SA Desai, SS Gundlur. Genetic variability studies for yield and its attributes under salt affected soils in wheat species. J. Farm Sci., 2022;35(2):169-173.
- Sunilyadav K, Singh A K, Baghel SS, Mahesh J. Assessment of Genetic variability and Diversity for Yield and its Contributing Traits among CIMMYT Based Wheat Germplasm. J. Wheat Res. 2014;6(1):154-159.
- USDA, United States Department of Agriculture. Available at: <http://www.fas.usda.gov>2018
- Vinod K, Devendra KP, Saiprasad SV. Genetic divergence analysis in durum wheat (*Triticum durum* Desf.). Int. J. Cur. Res. 2014;6(6):7001-7005.