

Optimization, a potential production, GC-MS and characterization of dark green pigment from new local isolate *Streptomyces nigra* strain GH12

Abstract

Due to increasing health awareness among consumers and the identification of novel pharmacological properties of diverse natural pigments, the market for natural pigments is currently seeing rapid growth. *Streptomyces. nigra* GH12, a new Actinomycete isolate identified by 16S rRNA used for dark green pigment production. Manipulations of the factors affecting pigment production are the most effective way to have maximum yield of pigment. Starch 2.5% (w/v), ammonium nitrate 0.2% (w/v), dibasic sodium phosphate 0.1% (w/v), inoculum size 6% (v/v), initial pH level of 8, 150 rpm rotation speed, 37 °C temperature, and 9 days of incubation were found the optimum growth conditions for the highest green pigment production. The extraction of green pigment was performed efficiently with ethanol solvent. Results showed no effect on the pigment content at 40, 50 and 60°C. On heating pigment extract at 70, 80, 90 and 100°C for 60 min, retention of pigment remained as high as 97.4, 93, 86 and 79% of the total pigment density, respectively, even with continuous treatment at 100°C for more 60 min. The results indicated that the pigment extracts exhibited their greatest stabilities at high pH values, i.e. 8.0, 8.5, 9.0 and 10.0, during the entire holding time. The pigment was characterized by full UV-visible spectroscopic screening, where the potential absorption maximum of the target green pigment extract was recorded in the range of 300–360 nm, with a λ_{max} at 340 nm. Moreover, the chemical composition diversity of the pigment extract was further investigated by GC/MS analysis that revealed the presence of 57 metabolites in their silylated form.

Keywords: Actinomycetes, green pigment, culture conditions, extraction, UV and GC/MS characterization

1. Introduction

Pigments are defined as finely divided and usually water-insoluble colorants, absorbing and reflecting the visible light to show different colors [1]. They are of larger molecular weights, less water-soluble, and less transparent than dyes. The usage of pigments has a long history, dating back to the beginning of ancient civilizations (ancient China, India, and Egypt), whereby natural plants, insects, and minerals were used to dye textiles, color foods, manufacture paints, color the body in religious ceremonies, and more[2]. Pigments are used to manufacture various products because they can enhance the natural color or replace color lost during the manufacturing process, generating greater consumer appeal by adding a novel sensory aspect. In the field of colors development and production, a recent worldwide trend large-scale research is being done to replace synthetic dyes with natural colors. The colors derived from plants or microorganisms suggest a certain level of safety. Pigments' toxicological issues are less severe than those associated with their synthetic counterparts due to their previous history and usage habits [3,4]. Three primary sources for a sustainable eco-friendly production of bio-safe natural pigments include plants, animals, and microbes [5]. As a result, one of the exciting and developing fields of study that shows promise for a range of industrial applications is microbial pigment synthesis[6-8]. Additionally, some microbial pigments have been found to be stable to light, heat, and pH, contain pro-vitamin A, and exhibit anticancer action [9]. However, from an industrial standpoint, it is important to create a high-tech and economical harnessing for the widespread

manufacture of different microbial pigments [10]. Natural pigments are now used more frequently in cosmetics, pharmaceuticals, food, and textiles [11,12]. Additionally, it is important to note that the natural pigment market is expected to grow at a 12.4 CAGR to reach USD 3.5 billion in the food industry by 2027 [13]. One of the most intriguing genera among microorganisms that generate pigments is *Streptomyces*, which has a high capacity for reproduction and the ability to make melanin, one of the most prevalent pigments in the market.[14,15] .The intriguing genetic dispersion of this particular species of actinomycetes makes it appealing for replication in the biotechnology sector [16,17]. Additionally, *Streptomyces* are well known for producing a variety of bioactive compounds, including antibiotics, anti-inflammatory, antioxidants, and cytotoxins, through their secondary metabolism, which is highly abundant and structurally diverse [18,19]. Numerous generated compounds were colored and given the bioactivity potential showed by *Streptomyces* strains [20,21]. Additionally, many of the pigmented substances produced by *Streptomyces* could be a promising avenue for discovering bioactive pigments. According to Elattaapy and Selim, 2020 [22] chemical, physical, and physiological parameters, such as anaerobic environments and temperatures, have an impact on pigment formation. Actinomycetes produce pigments that contain bioactive agents like antibacterial, antioxidant, and anticancer agents. There is a lot of work being done to identify and use the microbial pigments and their producers due to the growing demand for biological pigments over synthetic colorants. Actinomycetes are regarded as the ideal targets since pigment production appears to be more prevalent in them than in any other microbial group. Accordingly, this research directed to find optimized production conditions and characterization of a green pigment biosynthesized by a new local *Streptomyces* isolate, namely *S. nigra* strain GH12.

2. Materials and Methods

2.1. Sample collection

Rhizosphere soil samples were procured from a depth of 8–10 cm using a sterile spatula. The samples were brought to the laboratory and processed further immediately.

2.2. Pre-treatment of the soil sample

The soil samples were pretreated in order to reduce the proportion of other microorganisms differ from Actinomycetes. The soil samples were dried at about 50–60°C for 5–10 min. [22].

2.3. Isolation and screening of Actinomycetes

In order to isolate Actinomycetes from different rhizosphere soil samples gathered from different locations in cultivated fields from Aswan, Red sea, Alfayoum and/or Kafr Elshykh governorates. The samples were dried at 60°C for 1 h and then 1g of the respective sample was suspended in 9 ml of distilled water taken in a pre-autoclaved sterile test tube. The suspension was then serially diluted up to 10^{-5} . The diluted suspension (100µl) of 10^{-3} , 10^{-4} , and 10^{-5} was spread plated on Actinomycetes isolation agar (AIA). The medium contained (g/L) starch (20), KNO_3 (2.0), K_2HPO_4 (1.0), CaCO_3 , (3.0), MgSO_4 (0.5), NaCl (0.5), FeSO_4 , (0.01), and agar (20) at pH (7.8). The culture was sterilized by autoclave at 121°C for 15 min and incubated at 28°C for 5 days. All the plates were screened for Actinomycetes colonies based on morphology and pigmentation. The actinomycete colony that showed significant coloration was sub-cultured onto fresh medium by streaking until a pure culture was obtained.

2.4. Molecular identification of isolate

Genomic DNA extraction

A single colony of each bacterium was cultured in a 100 ml conical flask (Pyrex, USA) containing 20 ml of TSB by shaking in an orbital shaker (Thermo Fisher Scientific, USA) at 180 rpm for 18 h at 37°C. Bacterial culture then subjected to genomic DNA extraction using GeneJET Genomic DNA Purification Kit (Thermo scientific, USA) according to the manufacturer's instructions. Genomic DNA was used for 16s rDNA gene amplification.

2.5. 16s rDNA amplification and sequencing

The fragment of 16s rDNA gene was amplified in a reaction volume of 25 µl containing 12.5 µl of PCR Master Mix 2x conc. (Thermo scientific, USA). One microliter of each forward and reverse primer sets 8 F/1492 R, 9.5 µl of sterile deionized water, and 1 µl of the bacterial genomic DNA were mixed. The reaction was achieved in Gene Amp PCR system 9600 (Applied Biosystems, Bedford, MA, USA). A temperature cycling program [23] was applied with a slight modification by a hot initiation as 1 cycle of 95°C for 10 min, 40 cycles of 95°C for 1 min, 52°C for 1 min, 72°C for 2 min, and a final extension of 72°C for 10 min. PCR product was analyzed on a 1% agarose gel by electrophoresis using a 100-bp ladder DNA marker (Invitrogen, California, USA). The gel was visualized and photographed using TMXR + Gel Documentation System (Bio-Rad, California 94547, USA). Purified PCR product was subjected to sequencing by Sanger sequencing method using sequencer 3500 genetic analyzer, big dye X terminator kit (Thermo Fisher, USA) for forward and reverse directions in biomedical laboratory of colors (Clinilab, Egypt). BLASTn was used to detect the evolutionary relationship with other relatives (<https://www.ncbi.nlm.nih.gov/blast/Blast.cgi>). Multiple sequence alignment of 16s rRNA sequence from this study plus 16s rRNA sequences retrieved sequences from Gene Bank was performed using MUSCLE algorithm [24] available in MEGA X [25]. The evolutionary history was inferred using the neighbor-joining method [26].

2.6. Inoculum seed preparation

Five days old Actinomyces culture slants were obtained by addition 10 ml sterilized water, where the growth was crushed with culture loop. 1% of spore suspension was used to inoculate 250-ml conical flasks containing 50 ml of Pigment production medium constituted (g/L) starch, 20, KNO₃, 2.0, K₂HPO₄, 1.0, MgSO₄, 0.5, NaCl, 0.5, CaCO₃, 3.0, FeSO₄, 0.01, pH, 7.8. Afterward, incubation was done in a rotary shaker at 150 rpm for 5 days at 30°C. The culture growth was used to inoculate the experimental flasks at 2% v/v.

2.7. Standardization of culture conditions for optimum pigment production

The effect of various cultural conditions were studied like pH (6, 6.5, 7, 7.5, 8, 8.5 and 9), different carbon, nitrogen, and phosphorous sources and their concentrations, different inoculum size (2-8) %, different culture medium volume (25-125) ml of broth medium in 250 ml conical flask, different heavy metals ions (Zn⁺⁺, Mn⁺⁺, Cu⁺⁺ and Cr⁺⁺⁺), different incubation temperatures (25, 28, 31, 34, 37 and 40°C) and different incubation period (3-12 days). The pigment production was studied separately by inoculating bacterial suspension of *S. nigra* GH12.

2.8. Extraction of the pigment

At the end of fermentation period, the whole biomass of culture filtered off through a Whatman No. 1 filter paper. The filtrate was then centrifuged at 10,000 rpm for 10 min. The supernatants were considered the source of extracellular pigment. The analysis of pigment production was conducted by measuring the absorbance of the filtered extract at 340 nm using

spectrophotometer (UV spectrophotometer V-630, JASCO, Japan). Extraction of the pigments was carried out using different solvents ethanol, methanol, acetone, propanol, hexane, water and supernatant samples was taken and mixed well. The mixture was centrifuged at 8,000 rpm for 10 min; the supernatant was monitored at 340 nm to check the optical density. Finally, the extract was lyophilized and used for further characterization studies [27, 28].

2.9. Characterization of the pigment by GC/MS analysis

Sample derivatization

To the dried residue, add 100 μ l of derivatization reagent (80 μ l BFSTA+20 μ l TMCS) and incubate for 1hr at 65°C and inject into GC/MS. The most critical point is to avoid any water or moisture during derivatization step especially the silylating step is highly vulnerable. Chemical composition of the sample was performed using Trace GC-TSQ mass spectrometer (Thermo Scientific, Austin, TX, USA) with a direct capillary column TG-5MS (30 m x 0.25 mm x 0.25 μ m film thickness). The column oven temperature was initially held at 50°C for 5 min, then increased by 5°C /min to 250 °C, followed by holding for 2 min, raised to the final temperature 300°C at a rate of 30°C /min and finally maintained for 2 min. The injector and MS transfer line temperatures were kept at 270, 260°C respectively; Helium was used as a carrier gas at a constant flow rate of 1 ml/min. The solvent delay was 4 min and diluted samples of 1 μ l were injected automatically using Auto sampler AS1300 coupled with GC in the split mode. EI mass spectra were collected at 70 eV ionization voltages over the range of m/z 50–650 in full scan mode. The ion source temperature was set at 200 °C. The components were identified by comparison of their mass spectra with those of WILEY 09 and NIST 14 mass spectral database [29].

3. Results and discussion

3.1. Isolation

Actinomycetes are group of prokaryotic micro-organisms gram-positive filamentous bacteria. Actinomycetes are widely distributed in the natural habitats, like different cultivated lands, riveres and seas waters as well as various natural eco-system. A number of 225 Streptomyces colonies were randomly chosen and isolated from different soil sources of Egypt on the basal starch nitrate medium. Out of the 225 Streptomyces isolates, *S. nigra* GH12 strain was sharp pigment producer with green pigmentation, which then selected for further studies (Figs. 1&2).




Figure 1. (a) Morphology of isolated colony; (b) Pigment production of *S. nigra* GH12



Figure 2. Diffusible green pigment in the media produced by *S. nigra* GH12

3.2. Molecular identification of the Actinomycetes isolate

The Actinomycetes isolate's ribosomal internal transcribed spacer (ITS1-8.5SITS2) rDNA was effectively amplified using conventional PCR, and it was detected as a clear band of about 600 bp (Fig. 3). The NCBI Gene Bank database accepted the *Streptomyces* isolate identified in the current study and registered it with accession number OQ145630. The analysis of the *S. nigra* GH12 phylogeny was done using the sequence of the ITS region. By comparing this isolate to the ITS recorded sequences that were retrieved from NCBI, it was possible to identify the isolate at the genus and species levels. The obtained phylogenetic tree showed that GH12 isolate belongs to the genus of *Streptomyces*, had a tight relationship with each other. The cluster showed a close relationship in the same clade, with a bootstrap value of 100%, as presented the cluster (Fig. 3).




Figure 3. Phylogenetic tree of *S. nigra* GH12 based on 16S rRNA gene sequences. The 16S rRNA sequences were obtained from Gene Bank

3.3. Optimum culture conditions for highest pigment production

In general, the production of pigment is related to cell growth and is influenced by nutritional influences (such as carbon and nitrogen sources), microbiological parameters (such as spore, seed, and inoculum ages), and environmental variables (such as spore, seed, and inoculum ages). In terms of environmental conditions, many important parameters should be examined and improved [2]. These parameters include medium's humidity, the substrates' physical and structural characteristics, temperature, pH and agitation [30] (de Castro and Sato 2015). Due to the low moisture content, only a few microorganisms, primarily yeasts and fungi, are able to perform this cultivation procedure. Submerged fermentation (SmF), which depends on liquid culture for most microbes, is the second cultivation method. It is significantly influenced by variables such as temperature, pH, and agitation [31].

3.4. Determination of suitable initial pH value

Results in Figure 4 illustrated that the new local *S. nigra* GH12 strain produced its maximum pigment concentration at pH 8, the concentration production of green pigment is highly affected by low or high values than pH 8. The pH of fermentation medium may be altering the color of pigment produced by the same organisms. Despite the growth rate not being affected in the same way, the results show that the initial pH has a substantial impact on pigment synthesis [22] (Elattaapy and Selim, 2020). According to Sethi *et al.*, [32] *Penicillium purpurogenum* BKS9 produced the most red pigment at a pH of 6.0. Additionally, *Penicillium aculeatum* ATCC 10409 produced the most yellow pigment when grown at pH 6.5, according to [33]. The highest biomass and pigment synthesis by *Penicillium sp.* were noted at an initial pH of 9.0, according to Gunasekaran and Poorniammal's 2008 research. On the other hand, Méndez *et al.*, 2011 [34] found that *P. purpurogenum* GH2 produced the most red pigment at pH 5 and 24°C.




Figure 4. Effect of different initial pH value on the green pigment production by *S. nigra* GH12

3.5. Selection of proper carbon source

Data presented in figure (5) showed that insoluble starch was more stimulating carbon source for secretion of the green pigment in the growth medium. However, derivatives of starch, maltose, glucose and galactose, lactose, mannitol can produce about 25-40 % comparable to starch. Other various carbon sources failed to stimulate the green pigment production by *S. nigra* GH12 i.e

sucrose, farctouse, xylose and dextran (Fig. 5). Santos-Ebinuma *et al.*, 2013 [35] reported similar results for *P. purpurogenum* that has shown the capability to utilize starch and sucrose for the best pigment production. In addition, starch was the best for red pigment production by *Paecilomyces sinclairii* [36,37]. Mannitol was used as the best carbon source for production of red microbial pigments [38] Moreover, Sankhyayan *et al.*, 2019 [39] found that lactose and fructose strongly decreased pigment production.




Figure 5. Effect of different carbon sources on the green pigment production by *S. nigra* GH12

3.6. Effect of starch concentration on the green pigment production by *S. nigra* GH12

The green pigment secretion in the growth medium inoculated with *S. nigra* GH12, was increased by increasing the introduced starch in the growth medium from 0.5% up to 2.5% (w/v) then affected negatively by adding more starch in the medium (Fig. 6). Starch is more widely applied as carbon source for pigment production by Actinomycetes, while glucose and fructose were reported as more suitable carbon source for pigment production by *Sarcina* sp. and *Exiguobacterium aurantiacum* FH [40,41].




Figure 6. Effect of starch concentration on the green pigment production by *S. nigra* GH12.

3.7. Effect of inorganic nitrogen source on the green pigment production

It is well known the importance of nitrogen for growth of all alive organisms and its essential for metabolism that affected greatly by the presence of nitrogen source in the growth medium. The consumption of the nitrogen source and the metabolism associated to a favorable metabolic way both are necessary for the formation of pigments [35]. Previous studies have proved that the effect of a nitrogen source on pigment production is strain dependent, while using one of nitrogen sources could promote pigment production, using another nitrogen source could inhibit pigment production by the same isolate [22]. For instance, ammonium nitrate is more suitable than other inorganic nitrogen sources for more green pigment secretion in the growth medium inoculated with *S. nigra* GH12 (Fig. 7).




Figure 7. Influence of inorganic nitrogen source on green pigment production by *S. nigra* GH12

3.8. Effect of different concentration of ammonium nitrate on the green pigment production

Figure 8 illustrated the green pigment secretion in the growth medium inoculated with *S. nigra* GH12 contained the optimum above level of starch and adding different concentration of ammonium nitrate as a suitable inorganic nitrogen source. It seems that 2.0 g/l of ammonium nitrate is the more suitable concentration to give the high green pigment production and above or below 2.0 g/l decrease pigment production. The presence high nitrogen concentration in fermentation medium has terminal effect. Joshi, *et al.*, 2011 [41] found also potassium nitrate is the best for microbial pigment production by *Sarcina* sp isolated from water. Furthermore, potassium nitrate (0.5%) was found also the more probe nitrogen source for production of pigment by *Pseudomonas* sp.




Figure 8. Influence of different concentrations of ammonium nitrate on the green pigment production by *S. nigra* GH12

3.9 Influence of organic nitrogen source and its concentration on the green pigment production

Four organic nitrogen sources i.e. casein, malt extract, yeast extract and peptone at three different concentrations of 0.5, 1.0 and 1.5 % were examined (Fig. 9). The presence of 0.5 g/l malt extract in the medium was the more suitable organic source for production the green pigment by *S. nigra* GH12. This result was similar to the findings obtained by Elattaapy and Selim, [22] had enhanced red pigment production using malt extract by *Penicillium* sp. Many researchers reported that the presence of organic source in the culture media resulted in higher accumulation of pigment [42] during studying the optimization factors for maximum pigment production by *Rhodotorula glutinis*, *Exiguobacterium* sp. and *Sarcina* sp. [41]. According to Santos-Ebinuma *et al.*, 2013 [35] yeast extract followed by malt extract were both effective in inducing *P. purpurogenum* to produce pigment.




Figure 9. Influence of organic nitrogen sources on green pigment production by *S. nigra* GH12

3.10. Effect of phosphorous source on the green pigment production

Among the five phosphorous sources examined, i.e. as supplement in growth medium, the dibasic sodium phosphate was the preferred source of phosphate to produce the maximum green

pigment concentration by *S. nigra* GH12 in the growth medium. However, sodium dihydrogen phosphate was lowest one (Fig. 10).




Figure 10. Effect of phosphorous source on green pigment production by *S. nigra* GH12

3.11 Effect of phosphorous concentration on the green pigment production

Different five concentrations from dibasic sodium phosphate were tested in the fermentation medium of *S. nigra* GH12. The obtained results revealed that the maximum relative concentration of the green pigment at the concentration of 1.0 (g/L).




Figure 11. Effect of phosphorous level on the green pigment production by *S. nigra* GH12

3.12 Effect of inoculums size (v/v) on the green pigment concentration production

Study of inoculums percent was taken into consideration, when the pigment produced from microorganisms [43,44] Fig .12 illustrated that the inoculation the medium with 6 % (v/v) inoculums recorded the maximum green pigment production by *S. nigra* GH12 than other lower or higher sizes of inoculums. The maximum pigment synthesis was recorded by Elattaapy and Selim, 2020 [22] at a spore concentration of 106/ml. The ideal inoculum size for pigment formation, according to [45] was 3 ml of 9x10⁴ spores/ml. The ideal inoculum size for pigment formation, according to Velmurugan *et al.*, 2011 [46] was 4 ml of spore suspension. With 108 spores/ml, Santos-Ebinuma *et al.*, 2013 [35] has produced the higher pigment concentration with 108 spore/ml. Arora *et al.*, 2012 [46] reported that high inoculum size led to increases biomass and decreases pigment production, due to the inhibition the affiance of utilization of essential nutrients of culture medium by increased biomass.




Figure 12. Effect of inoculum size (v/v) on the green pigment production by *S. nigra* GH12

3.13. The effect of medium volume

Different volumes of pigment production medium from the range of 25-125 ml were introduced in 250 ml conical flasks capacity. The findings of this study demonstrated that the introduction 75 ml of the growth medium in 250 ml conical flasks capacity was more suitable to produce maximum concentration of green pigment (Fig. 13). The culture volume affects the amount of oxygen available for organism growth, utilization nutrient present in culture, and consequently biosynthesis of pigment.




Figure 13. Effect of culture volume (v/v) on the green pigment production by *S. nigra* GH12

3.14. Effect of metal ions addition

Metal ions are very essential for biosynthesis pathways in cells as enzymes, coenzymes and bioactive compounds production. Data plotted in figure 14 revealed that Cr^{+++} ion is very essential in the culture medium for green pigment production by *S. nigra* GH12 followed by Mn^{++} .




Figure 14. Effect of metal ions on the green pigment production by *S. nigra* GH12

3.15. Effect of incubation temperature on the green pigment production

Temperature is a one of vital factors for growth, nutrients metabolism and scan dry bioactive agent production, as well. It is showed that the green pigment concentration secreted in the growth medium by *S. nigra* GH12 strain was optimum at an incubation temperature of 37°C (Fig 15). Above temperature 37°C the pigment affected sharply negative. Joshi *et al.*, 2011 [41] reported 35°C as an optimum temperature for the pigment production by *Sarcina sp.* cultivated on apple pomace. However, according to Zahan *et al.*, 2023 [47], *Penicillium minioluteum* ED24 pigment synthesis requires an incubation temperature of roughly 30°C. Furthermore, according to Afshari *et al.*, 2015 [33] temperature affects the metabolic activity of fungi and their growth. The regulation of enzymatic activity inside fungal cells may be influenced by the temperature of the incubation process. The highest pigment production obtained at the optimum temperature because the enzyme activity for pigment synthesis seems to be enhanced at this temperature.




Figure 15. Effect of incubation temperature on the green pigment production by *S. nigra* GH12

3.16. Effect of incubation period on the green pigment production

On verification of figure 16, it is clear that the density of green pigment liberated by *S. nigra* GH12 in fermentation medium increased by increasing the age of fermentation period till ninth day then achieved a steady state. After 8 days, Elattaapy and Selim, 2020 [22] observed their highest pigment synthesis; nevertheless, as more incubation time passed, the production

markedly reduced. Santos-Ebinuma *et al.*, 2013 [35] found that *Penicillium purpurogenum* DPUA 1275 produced the most pigment after 12 days of incubation, but Chadni *et al.*, 2017 [48] found that *Talaromyces verruculosus* produced the most pigment after 24 days of incubation. On the other hand, many authors have reported shorter incubation times. Méndez *et al.*, 2011 [34], found that *P. purpurogenum* GH2 produced the highest yield of red pigment after 150 hours (6.25 days) of incubation, and Babitha *et al.*, 2006 [43], found that *Monascus purpureus* produced the most pigment after 6 days [49], however, achieved maximum pigment production by the same method. Previous studies reveal that the optimum incubation time for maximum pigment production varies from one strain to another.




Figure 16. Effect of incubation period on the green pigment production by *S. nigra* GH12

3.17. Extraction of the pigment of *S. nigra* GH12

In fact, the highest yield green pigment extract is produced by using ethanol as an extracting solvent via liquid-liquid extraction method that was found to be different from the reports by Abraham and Chauhan, 2018 [50]. Thereafter, the recovered extract was vacuum dried at a lower pressure (Fig. 17).



Figure 17. (a). Pigment extracted by ethanol solvent; (b). Dried pigment after ethanol extraction

3.18. Effect of heat treatment on green pigment stability

Thermal stability of extracted pigment was studied by heating the extracts at 40, 50, 60, 70, 80, 90 and 100 °C for 60 min (Fig. 18). Based on the absorbance values measured for green extracts before and after heat treatments, retentions of pigment as related to heating temperature were calculated. The stability of green pigment was markedly influenced by heat treatment. At 40, 50 and 60°C, no significant loss occurred in pigment content. When the heating temperature was elevated up to 70, 80 and 90°C, retention of pigment was still as high as 97.4, 93 and 86%, respectively after 60 min. Even when heat treatment was carried out at 100 °C for 60 min, green pigment retained more than 79% of its pigment content.




Figure 18. Effect of heat treatment on the retention of green pigment produced by *S. nigra* GH12 heated for 1 hr at different temperatures

3.19 Effect of pH on the stability of green pigment produced by *S. nigra* GH12

Color stability of pigment extracts was determined at a wide range of pH values between 4 and 10 at a temperature of 37°C by measuring the absorbance values of green pigment at λ_{\max} of 340 nm after 1hr. The results obtained are illustrated in Fig. 19. Data showed that green pigment were almost stable under aqueous conditions. It was observed that the green color of the extracts retained its high stability at pH 8.5.




Figure 19. Effect different pH on retention of green pigment produced by *S. nigra* GH12

3.20. UV-absorption spectrum of extracted green pigment produced by *S. nigra* GH12

The UV absorption spectrum of green pigment extract produced by *S. nigra* GH12 is shown in Fig. 20. Data indicated that absorption maximum peak of green pigment extract was range from 300 to 360nm, with λ_{\max} at 340 nm. However, Kazi *et al.*, 2022 [28] has reported a λ_{\max} 247 for the produced pigment from *Streptomyces* species on full UV-scanning. On the other hand, Selim *et al.*, 2008 [51] has demonstrated a λ_{\max} 520 for roselle pigments.




Figure 20. The UV absorption spectrum of extracted green pigment produced by *S. nigra* GH12

3.21 GC/MS characterization of green pigment produced by *Streptomyces nigra* GH12

Different physicochemical techniques were used for characterization of many *Streptomyces* derived bioactive pigments such as UV-, IR-, Raman- and/ or NMR spectroscopy together with one or more of MS-spectrometric ionization tools[51,52,53,54,-55]. GC/MS hyphenated technique is one of the most powerful analytical profiling tools for a fast and precise chemical characterization of the natural extracts, and microbial bio-pigments, as well. Chen *et al.*, 2018

used both GC/ and LC/hyphenated chromatographs with MS-spectrometry for identification of > 500 and 38 compounds, respectively in a potent antitumor active extract from *Streptomyces nigra* sp. nov. Sinapyl alcohol, phloroglucinol, azelaic acid, hydroxyurea, rosteron, shikimic acid, spermidine, 2-deoxy-D-glucose, and dehydroepiandrosterone have been detected among major identified metabolites. In the current study GC/MS analysis identified 57 compounds in the form of their silylated derivatives (Table 1) from the green pigment of the investigated new *S. nigra* GH12 on the basis of matching the corresponding MS-spectra and other output parameters with those of WILEY 09 and NIST 14 mass spectral library database [29]. Fig. 21 displayed the corresponding total ion current (TIC), that showed how much crowded the target pigment with different types of natural organic metabolites. It is worth mentioning that nine major silyl derivatives recorded for Lactic acid (**3**, 6.18 min, 19.72%), 3-Methylbutanoic acid (**5**, 8.73 min, 11.82%), Carbamic acid (**2**, 6.03 min, 7.80%), Ethylene (**13**, 14.78 min, 6.35%), *meso*-Erythritol (**23**, 17.97 min, 4.98%), 2-Methylpropanoic acid (**25**, 19.02 min, 4.54%), Palmitic acid (**42**, 28.12 min, 2.54%), 4-Hydroxy-*N*-valeric acid (**15**, 16.20 min, 2.20%) and 1-Monopalmitin (**51**, 37.29 min, 2.11%). In addition, twelve metabolites, i.e. **6**, **9**, **17-19**, **21**, **24**, **26**, **28**, **34**, **35** and **49** demonstrated more than 1% relative concentration. Their MS spectra were presented in figure 1S.

Table 1. Identified silylated derivatives of the constitutive metabolites in the green pigment of *S. nigra* GH12 isolate

P.No.	R _t , min	RC, %	MF	MW	Chemical name
1	4.02	0.38	C ₈ H ₁₉ F ₃ N ₂ Si ₂	256	<i>N,N'</i> -Bis(trimethylsilyl)trifluoroacetamine
2	6.03	7.80	C ₇ H ₁₉ NO ₂ Si ₂	205	Carbamic acid, <i>N</i> -(trimethylsilyl)-, trimethylsilyl ester
3	6.18	19.72	C ₉ H ₂₂ O ₃ Si ₂	234	Lactic acid, bis(trimethylsilyl) derivative
4	8.55	0.61	C ₁₀ H ₂₄ O ₃ Si ₂	248	Propanoic acid,2-methyl-3-[(trimethylsilyl)oxy]-, trimethylsilyl ester
5	8.73	11.82	C ₁₁ H ₂₆ O ₃ Si ₂	262	Butanoic acid,3-methyl-2-[(trimethylsilyl)oxy]-, trimethylsilyl ester
6	10.62	1.39	C ₁₂ H ₂₈ O ₃ Si ₂	276	2-Hydroxyisocaproic acid, bis(trimethylsilyl) derivative
7	10.73	0.95	C ₁₂ H ₂₈ O ₃ Si ₂	276	Pentanoic acid 3-methyl-2-[(trimethylsilyl)oxy]-, trimethylsilylester
8	11.83	0.88	C ₁₂ H ₃₂ O ₃ Si ₃	308	Glycerol, tris(trimethylsilyl) derivative
9	11.90	1.02	C ₈ H ₁₆ O ₃ Si	188	Levulinic acid, trimethylsilylester
10	12.45	0.31	C ₁₃ H ₃₄ O ₃ Si ₃	322	1,2,3-Butanetriol, tris(trimethylsilyl) derivative
11	14.28	0.57	C ₁₁ H ₂₄ O ₄ Si ₂	276	3,4-Dihydroxy-5-methyl-dihydrofuran,-2-one,(D)-, bis(trimethylsilyl) derivative
12	14.38	0.76	C ₁₇ H ₄₂ O ₅ Si ₄	438	<i>Erythro</i> -pentonic acid, 2-deoxy-3,4,5-tris-O-(trimethylsilyl)- trimethylsilyl ester
13	14.78	6.35	C ₁₁ H ₂₈ O ₃ Si ₃	292	Tris(trimethylsiloxy)ethylene
14	15.68	0.63	C ₁₃ H ₃₂ O ₄ Si ₃	336	(R,S)-3,4-dihydroxybutanoicacid trimethylsilylester
15	16.20	2.20	C ₁₁ H ₂₆ O ₃ Si ₂	262	4-Hydroxy- <i>N</i> -valeric acid bis(trimethylsilyl)
16	16.35	0.38	C ₁₃ H ₃₄ O ₃ Si ₃	322	1,2,3-Butanetriol, tris(trimethylsilyl) derivative
17	16.45	1.18	C ₁₈ H ₄₆ O ₄ Si ₄	438	3,8-Dioxa-2,9-disiladecane,2,2,9,9-tetramethyl-5,6-

18	16.63	1.47	C ₁₁ H ₂₈ O ₃ Si ₃	292	bis[[trimethylsilyloxy]methyl]- Tris(trimethylsilyloxy)ethylene
19	16.81	1.93	C ₁₀ H ₂₄ O ₃ Si ₂	248	Butanoic acid, 4-[[trimethylsilyloxy]-, trimethylsilylester
20	17.43	0.40	C ₁₄ H ₃₄ O ₄ Si ₃	350	D-Erythro-pentofuranose, 2-deoxy-1,3,5-tris-O- (trimethylsilyl)-
21	17.59	1.67	C ₁₀ H ₂₄ O ₂ Si ₂	232	2-Butene-1,4-diol, bis(trimethylsilyl) derivative
22	17.75	0.90	C ₁₆ H ₄₂ O ₄ Si ₄	410	L-Threitol, tetrakis(trimethylsilyl) derivative
23	17.97	4.98	C ₁₆ H ₄₂ O ₄ Si ₄	410	<i>Meso</i> -Erythritol, tetrakis(trimethylsilyl) derivative
24	18.63	1.29	C ₁₆ H ₄₀ O ₅ Si ₄	424	L-Threonic acid, tris(trimethylsilyl ether, trimethylsilylester
25	19.02	4.54	C ₁₃ H ₃₂ O ₄ Si ₃	336	Propanoic acid, 2-methyl-2,3-bis[[trimethyl silyloxy]-,trimethylsilylester
26	20.23	1.42	C ₁₇ H ₄₄ O ₄ Si ₄	424	Pentitol 3-desoxy-tetrakis(trimethylsilyl)
27	21.20	0.32	C ₁₇ H ₄₂ O ₅ Si ₄	438	Ribonic acid, 2-desoxy-tetrakis-O-(trimethylsilyl)-
28	21.36	1.27	C ₁₄ H ₃₁ BO ₆ Si ₂	362	α -D-Galactopyranoside,methyl 2,6-bis-O- (trimethylsilyl)-,cyclicmethylboronate
29	21.52	0.34	C ₁₈ H ₄₄ O ₅ Si ₄	452	1,5-Anhydrohexitol, tetrakis-O-(trimethylsilyl)- derivative
30	21.94	0.52	C ₁₄ H ₃₆ O ₃ Si ₃	336	3,9-Dioxa-2,10-disilaundecane,2,2,10,10- tetramethyl-5-[[trimethylsilyloxy]-
31	22.03	0.65	C ₁₃ H ₃₂ O ₄ Si ₃	336	((4,5-bis[[trimethylsilyloxy]tris-O-(trimethylsilyl)- 2,2,10,10-tetramethyl-5-[[trimethylsilyloxy]-3,9- dioxa-2,10-disilaundecane
32	22.14	0.26	C ₁₄ H ₃₆ O ₃ Si ₃	336	
33	22.42	0.58	C ₂₀ H ₅₂ O ₅ Si ₅	512	Xylitol, pentakis(trimethylsilyl) derivative
34	23.02	1.47	C ₂₀ H ₅₂ O ₅ Si ₅	512	Adonitol, pentakis(trimethylsilyl) derivative
35	23.39	1.96	C ₁₆ H ₃₇ BO ₆ Si ₃	420	α -D-Galactopyranose,1,2,3-tris-O-(trimethylsilyl)-, cyclic methylboronate
36	25.03	0.54	C ₁₇ H ₄₂ O ₅ Si ₄	438	α -Lyxopyranose, tetrakis-O-(trimethylsilyl) derivative
37	25.11	0.48	C ₂₂ H ₅₄ O ₆ Si ₅	554	D-Pinitol, pentakis(trimethylsilyl)ether
38	26.14	0.59	C ₁₇ H ₄₂ O ₅ Si ₄	438	α -D-Xylopyranose, tetrakis(trimethylsilyl) derivative
39	26.51	0.36	C ₂₁ H ₅₂ O ₆ Si ₅	540	α -D-Allopyranose, pentakis(trimethylsilyl) derivative
40	27.00	0.43	C ₁₈ H ₄₄ O ₅ Si ₄	452	1,5-Anhydrohexitol, tetrakis(trimethylsilyl) derivative
41	27.67	0.64	C ₂₁ H ₅₂ O ₆ Si ₅	540	D-Allofuranose,pentakis(trimethylsilyl)ether
42	28.12	2.54	C ₁₉ H ₄₀ O ₂ Si	328	Palmitic acid, trimethylsilyl derivative
43	31.05	0.85	C ₂₁ H ₄₂ O ₂ Si	354	11-Octadecenoicacid, (E)-, trimethylsilyl derivative
44	31.59	0.38	C ₂₁ H ₄₄ O ₂ Si	356	Stearic acid, trimethylsilyl derivative
45	32.35	0.64	C ₁₅ H ₃₄ O ₅ Si ₃	378	Levoglucosan, tris-(trimethylsilyl) derivative
46	32.82	0.35	C ₂₇ H ₆₆ O ₈ Si ₆	686	Glyceryl-glycoside trimethylsilyl ether
47	32.92	0.50	C ₁₈ H ₄₄ O ₅ Si ₄	452	1,5-Anhydrohexitol, tetrakis(trimethylsilyl)- derivative
48	33.01	0.39	C ₃₆ H ₈₆ O ₁₁ Si ₈	918	2- α -Mannobiose, octakis(trimethylsilyl) ether
49	33.53	1.81	C ₁₉ H ₃₈ N ₂ O ₆ Si ₃	474	5-Methyluridine, tris(trimethylsilyl)- derivative

50	36.81	0.42	C ₂₅ H ₅₄ O ₄ Si ₂	474	2-Palmitoylglycerol, bis(trimethylsilyl) derivative
51	37.29	2.11	C ₂₅ H ₅₄ O ₄ Si ₂	474	1-Monopalmitin, bis(trimethylsilyl) derivative
52	38.28	0.46	C ₃₆ H ₈₆ O ₁₁ Si ₈	918	Maltose, octakis(trimethylsilyl) derivative, isomer 1
53	38.81	0.83	C ₃₆ H ₈₆ O ₁₁ Si ₈	918	α -D-Lactose, (isomer 2), octakis(trimethylsilyl)
54	40.01	0.79	C ₂₆ H ₃₆ O ₅ Si	456	(+)--(6-Endo-7-exo)-7-[4-[(<i>tert</i> -butyldimethylsilyl)oxy]-3-methoxyphenyl]-3-methoxy-6-methyl-5-(2-propenyl)bicyclo[3.2.1]oct-3-ene-2,8-dione
55	40.19	0.31	C ₃₅ H ₇₄ N ₄ O ₁₃ Si ₅	898	L-Asparagine, N ϵ -[2-(acetylamino)-4-O-[2-(acetylamino)-2-deoxy-3,4,6-tris-O-(trimethylsilyl)- α -D-glucopyranosyl]-2-deoxy-3,6-bis-O-(trimethylsilyl)- α -D-glucopyranosyl]- α -D-glucopyranosiduronic acid, 3-(5-ethylhexahydro-2,4,6-trioxo-5-pyrimidinyl)-1,1-dimethylpropyl 2,3,4-tris-O-(trimethylsilyl)-, methylester
56	41.02	0.49	C ₂₇ H ₅₂ N ₂ O ₁₀ Si ₃	648	α -D-Glucopyranosiduronic acid, 3-(5-ethylhexahydro-2,4,6-trioxo-5-pyrimidinyl)-1,1-dimethylpropyl 2,3,4-tris-O-(trimethylsilyl)-, methylester
57	41.83	0.68	C ₁₇ H ₄₂ O ₅ Si ₄	438	α -Arabinopyranose, tetrakis(trimethylsilyl) derivative




Figure 21. TIC chromatogram for the total green pigment of *S. nigra* GH12 isolate

Conclusion

This research directed to find optimized production conditions and characterization of a green pigment biosynthesized by a new local *Streptomyces* isolate, namely *S. nigra* strain GH12. the chemical composition diversity of the pigment extract was further investigated by GC/MS analysis that revealed the presence of 57 metabolites in their silylated form.

REFERENCES

1. Rapp G. Pigments and colorants archaeomineralogy. Springer, Berlin. 2009; 201–221.
2. Lyu X, Lyu Y, Yu, H, Chen W, Ye L, Yang R. Biotechnological advances for improving natural pigment production: a state-of-the-art review. *Bioresources and Bioprocessing*. 2022; 9:8. doi.org/10.1186/s40643-022-00497-4
3. Valle-Veg, P, Lucas-Florentino, B. Toxicología de Alimentos. Instituto Nacional de Salud Pública, Centro Nacional de Salud Ambiental: Cuernavaca, México. 2000; ISBN 92 75 37004
4. Aixa A. Sarmiento-Tovar, Laura Silva, Jeysson Sánchez-Suárez and Luis Diaz. Streptomyces-Derived Bioactive Pigments: Ecofriendly Source of Bioactive Compounds. *Coatings*. 2022; 12: 1858 32 -35.
5. Fakruddin M. Biosurfactant: Production and Application. *J. Pet. Environ. Biotechnol*. 2012; 3:124.
6. Pailliè-Jiménez ME, Stincone P, Brandelli A. Natural Pigments of Microbial Origin. *Front. Sustain. Food Syst*. 2020; 4: 590439.
7. Morales-Oyervides L, Oliveir, J, Sousa-Gallagher M, Méndez-Zaval A, Montañez JC. Assessment of the Dyeing Properties of the Pigments Produced by *Talaromyces* spp. *J. Fungi*. 2017; 3: 38-43.
8. Paul T, Bandyopadhyay TK, Mondal A, Tiwari ON, Muthuraj M, Bhuni B. A Comprehensive Review on Recent Trends in Production, Purification, and Applications of Prodigiosin. *Biomass Convers. Biorefinery*. 2020; 12:1409–1431.
9. Müller MM, Hausmann R. Regulatory and Metabolic Network of Rhamnolipid Biosynthesis: Traditional and Advanced Engineering towards Biotechnological Production. *Appl. Microbiol. Biotechnol*. 2011; 91: 251–264.
10. Sen T, Barro, CJ. Deshmukh, S.K. Microbial Pigments in the Food Industry—Challenges and the Way Forward. *Front. Nutr*. 2019; 6& 7-13
11. Juric´ S, Juric´ M, Król-Kilin´ ska Z, Vlahovic´ek-Kahlina K, Vincekovic´ M, Dragovic´-Uzelac V, Donsi F. Sources, Stability, Encapsulation and Application of Natural Pigments in Foods. *Food Rev. Int*. 2020; 38: 1735–1790.
12. Rodriguez-Amaya DB. Natural Food Pigments and Colorants. *Curr. Opin. Food Sci*. 2016; 7: 20–26.
13. Kumar S, Kumar V, Ambika AA, Nag D, Kumar V, Darnal S, Thakur V, Patial V, Singh D. Microbial Pigments: Learning from Himalayan Perspective to Industrial Applications. *J. Ind. Microbiol. Biotechnol*. 2022; 49: kuac 017.

14. Metwally RA, El Sikaily A, El-Sersy NA, Ghozlan HA, Sabry SA. Antimicrobial Activity of Textile Fabrics Dyed with Prodigiosin Pigment Extracted from Marine *Serratia rubidaea* RAM_Alex Bacteria. Egypt. J. Aquat. Res. 2021; 47: 301–305.
15. Tuli HS, Chaudhary P, Beniwal V, Sharma AK. Microbial Pigments as Natural Color Sources: Current Trends and Future Perspectives. J. Food Sci. Technol. 2015; 52: 4669–4678.
16. Dharmaraj S. Marine Streptomyces as a Novel Source of Bioactive Substances. World J. Microbiol. Biotechnol. 2010; 26: 2123–2139.
17. Berdi J. Bioactive Microbial Metabolites. J. Antibiot. 2005; 58: 1–26.
18. Harir M, Bendif, H, Bellahcene M, Fortas Z, Pogni R. Streptomyces Secondary Metabolites. In Basic Biology and Applications of Actinobacteria; Intech Open: London, UK, 2018; 99–121.
19. Siddharth S, Vittal RR. Evaluation of Antimicrobial, Enzyme Inhibitory, Antioxidant and Cytotoxic Activities of Partially Purified Volatile Metabolites of Marine *Streptomyces sp.* S2A. Microorganisms .2018; 6:72-82.
20. Azman AS, Mawang CI, Abubakar S. Bacterial Pigments: The Bioactivities and as an Alternative for Therapeutic Applications. Nat. Prod. Commun. 2018; 13: 1747–1754.
21. Narsing Rao MP, Xiao M, Li W. Fungal and Bacterial Pigments: Secondary Metabolites with Wide Applications. Front. Microbiol. 2017; 8: 1113.
22. Elattaapy AM, Selim MAE. Factors Affecting Red Pigment Production by Local Fungal Isolate of *Penicillium sp.* J. of Agricultural Chemistry and Biotechnology. Mansoura Univ. 2020; 11 (9):255-261.
23. El-Sayed GM, Abosereh NA, Ibrahim SA, Abd-Elrazik AB, Hammad MA, Hafez FM. Identification of gene encoding organophosphorus hydrolase (OPH) enzyme in potent organophosphorus degrading bacterial isolates. J Environ Sci Technol. 2018; 11:175–189.
24. Edgar RC. (2004): MUSCLE a multiple sequence alignment method with reduced time and space complexity. BMC Bioinformatics 2018; 5:113. doi: 10.1186/1471-2105-5-113.
25. Tamura K, Stecher G., and Kumar S. MEGA 11: Molecular Evolutionary Genetics Analysis Version 11. Molecular Biology and Evolution. 2021; doi.org/10.1093/molbev/msab120.
26. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol Biol Evol. 1987; 4(4): 406–425.
27. Parmar R S, Singh C. A comprehensive study of ecofriendly natural pigment and its applications,” Biochemistry and Biophysics Reports. 2018; 13: 22–26.

28. Kazi, Z. et al. Production, characterization, and antimicrobial activity of pigment from *Streptomyces* species. *Journal of Nanomaterials*. 2022; 3962301.
29. Abd El-Kareem MSM; Rabbih MAE, Selim ETM, Elsherbiny EAE, El-Khatee AY. Application of GC/EIMS in Combination with Semi-Empirical Calculations for Identification and investigation of some volatile components in Basil essential oil. *International Journal of Analytical Mass Spectrometry and Chromatography* 04:14-25. DOI: 10.4236/ijamsc.2016;41002.
30. de Castro RJS. Sato HH. Biologically active peptides: Processes for their generation, purification and identification and applications as natural additives in the food and pharmaceutical industries. *Food Research International*. 2015; 74:185–198.
31. Morales-Oyervides L, Ruiz-Sanchez JP, Oliveira JC, Sousa-Gallagher MJ, Mendez-Zavala A, Giuffrida D, Dufosse L, Montanez J. Biotechnological approaches for the production of natural colorants by *Talaromyces/penicillium*: a review. *Biotechnol Adv*. <https://doi.org/10.1016/j.biotechadv.2020.107601>.
32. Sethi BK, Parida P, Sahoo SL, Behera BC. Extracellular production and characterization of red pigment from *Penicillium purpurogenum* BKS9. *Algerian Journal of Natural Products*, 2016 ;4(3): 379-392.
33. Afshari M, Shahidi F., Mortazavi SA, Tabatabai F, Es'haghi Z. Investigating the influence of pH, temperature and agitation speed on yellow pigment production by *Penicillium aculeatum* ATCC 10409. *Natural product research*. 2015; 29(14): 1300-1306.
34. Mendez A, Pérez C, Montañéz JC, Martínez G, Aguilar CN. Red pigment production by *Penicillium purpurogenum* GH2 is influenced by pH and temperature. *Journal of Zhejiang University Science B*. 2011; 12(12): 961-968.
35. Santos-Ebinuma VC, Roberto IC, Simas Teixeira M F, Pessoa Jr A. Improving of red colorants production by a new *Penicillium purpurogenum* strain in submerged culture and the effect of different parameters in their stability. *Biotechnology progress*. 2013; 29(3): 778-785.
36. Cho YJ, Park JP, Hwan HJ, Kim SW, Choi JW, Yu JW. Production of red pigment by submerged culture of *Paecilomyces sinclairii*. *Letters in Applied Microbiology*. 2002; 35(3): 195-202.
37. Gunasekaran S, Poorniammal R. Optimization of fermentation conditions for red pigment production from *Penicillium* sp. under submerged cultivation. *African journal of Biotechnology*. 2008; 7(12): 1894-1898.
38. Salim RG, Fadel M, Youssef YA, Taie HAA, Abosereh NA, El-Sayed GM, Marzouk M. A local *Talaromyces atroseus* TRP-NRC isolate: isolation, improvement, and biotechnological approach combined with LC/HRESI-MS characterization, skin safety, and

- wool fabric dyeing ability of the produced red pigment mixture *Journal of Genetic Engineering and Biotechnology*. 2022; 20:62.
39. Sankhyayan M, Walia A, Putatunda C. Production of red pigment from fungal isolate DMMS-1. *Int. J. Curr. Microbiol. App. Sci.* 2019; 8(4): 2839-2846.
 40. Sashidharan P, Raja R, Karthik C, Ranandakumar S, Indra Arulselvi, P Isolation and characterization of yellow pigment producing *Exiguobacterium sps.* *Journal of Biochemical Technology*. 2013; 4:632- 63.
 41. Joshi VK, Attri D, Rana NS. Optimization of apple pomace based medium and fermentation conditions for pigment production by *Sarcina sp.* *Indian J. Natural Product. Res.* 2011; 2: 421-427.
 42. El-Banna AA, Abd El-Razek AM, El-Mahdy AR. Some Factors Affecting the Production of Carotenoids by *Rhodotorula glutinis var. glutinis*. *Food and Nutrition Sciences*. 2012; 3(1): 17082. [10.4236/fns.2012.31011](https://doi.org/10.4236/fns.2012.31011)
 43. Babitha S, Soccol C R, Pandey A. Jackfruit seed-A novel substrate for the production of *Monascus* pigments through solid-state fermentation. *Food Technology and Biotechnology*. 2006; 44 (4): 465–471.
 44. Ji H, Jiang D, Cao L. “Optimization of fermentation parameters on T-DNA inserted *Monascus purpureus* mutant MT24 with high pigment production capacity,” *Res. J. Biotechnol.* 2012; 7: 9-14.
 45. Babitha S. Soccol R, Pandey A. “Solid-state fermentation for the production of *Monascus pigments* from jackfruit seed,” *Biores. Technol.* 2007; 98: 1554-1560.
 46. Arora P K, Sharma A, Mehta R, Shenoy BD, Srivastava A, Sing VP. “Metabolism of 4-chloro-2-nitrophenol in a Gram positive bacterium, *Exiguo bacterium sp.* PMA,” *Microbial. Cell Fact.* 2012; 11: 150-160.
 47. Zahan KA, Ismail N S, Leong C R, Ab Rashid S, Tong WY. Monascorubin production by *Penicillium minioluteum* ED24 in a solid-state fermentation using sesame seed cake as substrate. *Materials Today: Proceedings*. 2020; 31: 127–135.
 48. Chadni Z, Rahaman M H, Jerin I, Hoque KMF, Reza M A. (Extraction and optimisation of red pigment production as secondary metabolites from *Talaromyces verruculosus* and its potential use in textile industries. *Mycology*. 2017; 8(1): 48-57.
 49. Silbir, Goksungur Y. Natural red pigment production by *monascus purpureus* in submerged fermentation systems using a food industry waste: Brewer’s spent grain. *Foods*. 2019; 8(5):161.
 50. Abraham J, Chauhan R. Profiling of red pigment produced by *Streptomyces sp.* JAR6 and its bioactivity *Biotech.* 2018; 8(1): 22. doi: [10.1007/s13205-017-1044-7](https://doi.org/10.1007/s13205-017-1044-7).

51. Selim K A , Khalil K E, Abdel-Bary M S, Abdel-Azeim NA .Extraction, Encapsulation and Utilization of Red Pigments from Roselle (*Hibiscus sabdariffa* L.) as Natural Food Colourants Alex. J. Fd. Sci. & Technol. Special Volume Conference, 2008; 7-20.
52. Chen C, Ye Y, Wang R, Zhang Y, Wu C, Debnath SC, Ma Z, Wang J, Wu M . *Streptomyces nigra* sp. nov. is a novel Actinobacterium isolated from mangrove soil and exerts a potent antitumor activity *in vitro*. Front. Microbiol. 2018; 9:1587. doi: 10.3389/fmicb.2018.01587
53. Polapally R, Mansani M, Rajkumar K, Burgula S, Hameeda B, Alhazmi A, et al. Melanin pigment of *Streptomyces puniceus* RHPR9 exhibits antibacterial, antioxidant and anticancer activities. PLoS ONE 2022; 17(4): e0266676. doi.org/10.1371/journal.pone.0266676.
54. Ibrahim WM, Olama ZA, Abou- elela G M, Ramadan HS, Hegazy G E, El Badan DEl S. Exploring the antimicrobial, antiviral, antioxidant, and antitumor potentials of marine *Streptomyces tunisiensis* W4MT573222 pigment isolated from Abu- Qir sediments, Egypt. Microbial Cell Factories. 2023; 22:94.
55. Zhang S, Wu J, Jiang Z, Zhang L, Song T, Liu X, Yin C and Zhang Y .Pigments of aminophenoxazinones and viridomycins produced by termite-associated *Streptomyces tanashiensis* BYF-112. Front. Microbiol. 2023; 13:1110811. doi:10.3389/fmicb. 2022. 1110811.