

## Original Research Article

# Evaluation of Serum Leptin Levels and Some Anthropometric Parameters among Women of Reproductive Age in Port Harcourt, Nigeria.

### ABSTRACT

**Aim:** To evaluate and correlate the levels of leptin and some anthropometric parameters among women of reproductive age in Port Harcourt.

**Study Design:** The population of women of reproductive age in Port Harcourt Local Government Area was sampled. The samples were taken from Creek Road Market, Borokiri, Port Harcourt within the period November, 2022 to March, 2023.

Comment [m1]: Sampling tecqnique?

Comment [m2]: Delete.

**Methodology:** The study involved 150 subjects, who were within the age bracket of 20 and 60 years, who had given their informed consent to participate in this study, and were without chronic condition or undergoing any medical treatment or procedure. The anthropometric parameters, such as body mass index (BMI), Waist-to-Hip ratio (WHR) were obtained using appropriate techniques. Three milliliters (3ml) of venous blood was collected into plain bottles for the assay of leptin (using ELISA technique).

**Results:** The mean values of the BMI and WHR were  $28.72 \pm 10.31 \text{ kg/m}^2$  and  $0.83 \pm 0.22$  respectively. These were above the reference ranges. Leptin ( $3.56 \pm 0.93 \text{ ng/ml}$ ) was within the reference range. The results also show that the BMI and WHR correlated positively with the leptin levels. Also, BMI, WHR and leptin values increased with age.

**Conclusion:** There was positive correlation between leptin levels and all the anthropometric parameters showing that there might be a tendency of the population under study to be over weighted and obese. It is recommended that these parameters be monitored in women of reproductive age as part of the assessment of their health status.

**Keywords:** Anthropometric measurements, Leptin levels, reproductive age women, Port Harcourt.

## 1. INTRODUCTION

Anthropometric parameters serve as simple parameters that can provide valuable health information. Anthropometric indices are important clinical parameters which can be easily assessed at a low cost [1]. The proven link between anthropometric indices and leptin has immense public health implications. Anthropometric measurements are non-invasive, easy to apply, and inexpensive techniques which help to evaluate body composition of all ages [2]. Measurements of some anthropometric indices will be guidance for the growth and development of the future adult life.

However, leptin, a polypeptide hormone, secreted in concentrations proportional to body fat mass, plays an important role in several physiological functions like energy homeostasis, immunity, and reproduction, with possible implication in

other conditions (such as hepatic steatosis, depression etc.) [3, 4, 5]. Leptin is required to maintain normal body weight, as it lowers food intake and increases energy expenditure [3].

Obese individuals, however, might express high serum leptin concentrations but fail to properly control food intake and regulate the body's energy reserve, thus exhibiting leptin resistance. Moreover, high leptin levels that fail to regulate insulin secretion might suggest leptin resistance at the pancreatic  $\beta$ -cell level [5].

Anthropometric measurements and indices such as waist circumference (WC), hip circumference (HC), mid arm circumference (MAC), body mass index (BMI), waist to hip ratio (WHR) and waist to height ratio (WHtR) are used as simple standard measures to assess obesity and body fat levels worldwide [6,7,8]. Both serum leptin and anthropometric parameters are used to assess obesity. Thus, it would be of value if both these parameters were compared as parameters to assess obesity and related disorders.

There have been several studies on anthropometric parameters and leptin in different parts of the world. This includes study by Chigbu & Aja [9] in Southeast Nigeria proposing prevalence with BMI in 1st trimester alone as **Comment [m3]:** expansiate  
However not much has been established with regards anthropometric indices and leptin in Women of reproductive age living in Port Harcourt. For instance, values like waist circumference in African populations have not been properly defined due to lack of appropriate data, and therefore, it has been recommended that the cut-off points derived from European population groups are used for African subjects [10]. In fact, there may be obvious differences, **Comment [m4]:** Correct spelling  
the interpretation of Waist Circumference value as there seems to be a lack of universally accepted site for measuring Waist Circumference and the large variation of Waist Circumference optimal cut-off also affected by age, sex, race, and ethnicity; and so also with other anthropometric parameters as observed in even BMI which is noted not to be sensitive to Women [11,12] Again, in Africa, most work on this is limited by its design (retrospective) and small sample size which was recorded in Benin to be 323 [13]. **Comment [m5]:** Can you cite the results of few studies that looked at reproductive age women and leptin levels and what they got?

Hence evaluating the serum leptin levels and some anthropometric parameters in different age group of reproductive women living in Port Harcourt, Nigeria. **Comment [m6]:** Rephrase

## 2. MATERIALS AND METHODS

**2.1 Study Population:** The population of women of reproductive age in Port Harcourt Local Government Area is 281,120 [14]. Using the Yamani formula to calculate the sample size, we have:

$$n = \frac{N}{1 + N e^2}$$

where n= sample size, N= population of women of reproductive age in Port Harcourt = 281, 120 and e= error margin= 0.05. From the formula, n= 40 women. However, this study used a sample size of 150 subjects. **Comment [m7]:** Recheck this sample size calculation. n=399

**2.2 Study Area:** The samples were taken from Creek Road Market, Borokiri, Port Harcourt Local Government Area of Rivers State.

**2.3 Population Size:** A sample size of 150 apparently healthy women of reproductive age subjects between the ages of 20-60 years were used. A well-structured questionnaire was used to gather relevant information (such as age, Weight, Height, BMI, Waist Circumference, Hip Circumference, WHR, etc) from each subject after a written informed consent had been obtained from them.

**2.4 Data Collection:** Demographic data and clinical history were obtained using structured questionnaires. Body weight, height, waist circumference (WC), hip circumference (HC) were measured while body mass index (BMI) and waist to hip ratio (WHR) were calculated in all study subjects.

**2.5 Criteria for Sample collection:** All non-diabetic obese subjects had initial fasting plasma glucose (FPG) screening test done with Accu-check active glucometer to aid in the selection of subjects with normoglycaemia alone.

**2.6 Sample Collection:** 3 mls of venous blood sample under aseptic procedure from the cubital fossa of each subject after an overnight fast of at least 8 hours was taken into a plain specimen bottle for serum leptin analysis using an

enzyme linked immunosorbent assay (ELISA) method. The test was carried out and interpreted according to the manufacturer's instructions.

**Reagents:** Human LEP (Leptin) ELISA kit manufactured by ELK biotech, Denver, USA. Lot: 20365084741, Expiry Date: 30.11.2023, Mindray MR96A.

**2.7 Anthropometric Measurements:** Measurement of Height (H) as described by Ononamadu *et al.* [15] was carried out. Measurement of Weight (W) as described Lee *et al.* [16] was carried out on the subjects. Calculation of Body Mass Index (BMI) as described by Ononamadu *et al.* [15] was done. Measurement of Waist Circumference (WC) as described by WHO [17] (2011) was carried out and measurement of Hip Circumference (HC) as described by WHO [17] was carried out on the subjects. Then, calculation of Waist to Hip Ratio (WHR) as described by Lee *et al.* [16] was done.

**Comment [m8]:** Can you expaciate how all these measurements were actually done apart from just giving the references

**2.8 Statistical Analysis:** The data generated from this study were analyzed using Statistical Package for Sciences (SPSS) version 25 (SPSS Inc. Chicago Illinois). Results were expressed as mean  $\pm$  standard deviation (SD) and frequency (%), and presented in tables and charts as appropriate. Comparison of parameters with p-values less than or equal to 0.05, was considered statistically significant.

### 3. RESULTS

#### 3.1 Demographic Characteristics of Subjects

The demographic data collated indicated a significant difference between the age interval of subjects (P=0.131). There was significant difference in tribe, with Ijaw tribe having the highest frequency of 85. There was significant difference in the level of education of subjects as seen in the Table 1 as very few subjects have attained tertiary education. There was significant difference in the income status of subjects (P<0.001). There was significant difference in marital status of subjects (P= 0.033) and their parity as shown in Table 1.

**Table 1: Demographic Characteristics of Subjects**

Subjects	Frequency	Prevalence	p-value	X <sup>2</sup> -value
<b>Age Interval</b>				
a. 20 - 29	18	10	0.131	5.311
b. 30 – 39	32	17		
c. 40 – 49	60	51		
d. 50 - 59	40	22		
<b>Tribe</b>				
a. Ijaw	85	57	0.033	8.522
b. Ikwerre	42	28		
c. Ogoni/Elemé	05	03		
d. Igbo	18	12		
<b>Occupation</b>				
a. Business/Traders	150	100	-	-
<b>Level of Education</b>				

a. Primary	78	52	0.013	5.443
b. Secondary	55	37		
c. Tertiary	10	07		
d. None	07	04		
<b>Religion</b>				
a. Christianity	145	97	<0.001	10.153
b. Islam	05	03		
<b>Income Status</b>				
a. Low	140	94	<0.001	9.903
b. Middle	10	06		
c. Upper	0	-		
<b>Marital Status</b>				
a. Single	20	30	0.033	8.522
b. Married	87	58		
c. Divorced	25	17		
d. Widowed	18	12		
<b>Parity</b>				
a. Nullipara	21	14	0.001	5.443
b. Multipara	92	61		
c. Primipara	37	25		

### 3.2 Results of Anthropometric Parameters among the Subjects

The number of subjects with BMI of 18.5-24.9 (normal weight), 25-29.9 (overweight) and 30 & above (obese) were 39, 50 and 61. This showed no significant difference in the number of subjects in the different categories (P=0.089). The number of subjects with waist-to-hip ratio of "<0.81" and "0.81 and above" were 55 and 95 respectively showing a significant difference in the number of subjects (P=0.001). The waist circumference of subject with "<88cm" and ">88cm" results were 63 and 87 respectively showing a significant difference in number of subject (P=0.021) while the hip circumference of subjects with "<98cm" and ">98cm" were 5 and 145 respectively, also showing a significant difference in the number of subjects (P=0.001). This is reflected in Table 2.

**Table 2: Comparison of Anthropometric Parameters among the Subjects**

Anthropometric Parameter	Number	Prevalence	p-value	X <sup>2</sup> -value	Remark
Body Mass Index (kg/m <sup>2</sup> )					
<18.5	-				
18.5-24.9	39	26	0.089	4.840	NS
25-29.9	50	33			

**Comment [m9]:** Give a footnote on the meaning of these abbreviations in the remark column

30 & above	61	41			
Waist Circumference					
≤ 88	63	42	0.021	4.443	S
> 88	87	58			
Hip Circumference					
≤ 98	5	3	0.001	6.931	S
> 98	145	97			
Waist-to-hip ratio					
<0.81	55	37	0.001	10.667	S
0.81 & above	95	63			

### 3.3 Results of the mean values of anthropometric measurements with Leptin values among Subjects

The mean values of Body Mass Index, Waist-to-Hip Ratio, Leptin, Waist Circumference and Hip Circumference among the subjects were  $28.72 \pm 10.31 \text{ Kg/m}^2$ ,  $0.83 \pm 0.22$ ,  $3.56 \pm 0.93 \text{ ng/ml}$ ,  $94.40 \pm 8.05 \text{ cm}$  and  $113.96 \pm 5.38 \text{ cm}$  respectively. This is shown in Table 3.

**Table 3: Mean ± SD of Anthropometric Parameters /Leptin levels in the Subjects**

Parameter	Mean ± SD	Reference Values
Body Mass Index (kg/m <sup>2</sup> )	28.72 ± 10.31	18.5-24.9
Waist-to-hip ratio	0.83 ± 0.22	<0.81
Leptin (ng/ml)	3.56 ± 0.93	0.5-15.5
Waist Circumference	94.40 ± 8.05	≤88cm
Hip Circumference	113.96 ± 5.38	≤48cm

### 3.4 Comparison of Parameters according to Age Groups

The BMI for subjects within age ranges (years) of 20-29, 30-39, 40-49, 50-59, were 24.41±3.52, 27.52±3.46, 33.29±5.31, 33.75±2.71 respectively. There was a significant difference in BMI (P<0.001) among the different age ranges. The WHR for the age ranges were 0.82±0.03, 0.82±0.03, 0.85±0.03, 0.85±0.02 respectively showing a significant difference in WHR (P<0.001). The leptin levels (ng/ml) for subjects in the different age ranges were 2.19 ± 0.96, 2.84 ± 1.00, 6.03 ± 5.15, and 4.81 ± 2.67 respectively. There was a significant difference in the leptin levels (P<0.001) as shown in Table 4.

There was a positive correlation of leptin levels and the anthropometric parameters as shown on Figure 1 and Figure 2.

**Table 4: Comparison of Anthropometric Parameters according to Age Groups**

	BMI (kg/m <sup>2</sup> )	WHR	Leptin (ng/ml)
A (20-29)	24.41 ± 3.52 <sup>a</sup>	0.82 ± 0.03 <sup>a</sup>	2.19 ± 0.96 <sup>a</sup>
B (30-39)	27.52 ± 3.46 <sup>b</sup>	0.82 ± 0.03 <sup>a</sup>	2.84 ± 1.00 <sup>a</sup>
C (40-49)	33.29 ± 5.31 <sup>c</sup>	0.85 ± 0.03 <sup>b</sup>	4.03 ± 5.15 <sup>b</sup>
D (50-59)	33.75 ± 2.71 <sup>c</sup>	0.85 ± 0.02 <sup>b</sup>	4.81 ± 2.67 <sup>b</sup>
p-value	<0.001	<0.001	<0.001
F-value	24.169	8.415	8.989
Remarks	S	S	S

Values with different superscripts are significantly different from each other (P≤0.05)





Figure 1: Correlation Plot of Leptin versus BMI



**Figure 2: Correlation Plot of Leptin versus WHR**

#### **4. DISCUSSION**

This work assessed the levels of leptin and some anthropometric parameters among female residents of reproductive age in Port Harcourt. The study involved 150 female subjects within the ages of 20 and 60 years. From the results, 33% of the subjects were overweight. This finding is similar to the findings reported by Ghose, [18] in the South-East, Nigeria. This may be due to the fact that obesity has been reported to be higher among urban dwellers than in rural dwellers, occasioned by the introduction of processed foods and its increased consumption[18].

The results also indicates that there was no statistical difference in the number of subjects in the different classes of BMI ( $P=0.089$ ). Again, the mean BMI for the subjects in this study was higher than the WHO reference range for healthy classification indicating that the subjects in this study are regarded as obese. This may be due to changes in the dietary pattern among the subjects, which has been reported to be an important underlying factor for the increasing prevalence of overweight/obesity and associated complications, especially among women[18]. Sedentary lifestyle has also been reported to be a significant factor for obesity and associated complications [19]. This is seen in the lifestyle of these women in the market. This finding agrees with the work of Onyeji & Sanusi [20], who reported a similar finding among women of reproductive age in South-East, Nigeria.

From the study, the number of subjects who had WHR above the reference range was significantly more than the number of subjects whose WHR were below the reference range ( $P=0.001$ ). The WHR has been reported to be higher in females, as well as higher among urban dwellers. This is linked to the dietary and lifestyle changes in our society. Our findings indicates that truncal obesity is common among the study subjects. This finding agrees with the work of Tagbo *et al.*[21].

The results from this study indicate that the mean BMI and WHR of the subjects were above the reference limits. However, the mean values of leptin levels were within the reference range. These findings indicate that the increased anthropometric parameters are not as a result of hyperglycaemia because the subjects were initially screened for normal fasting blood sugar level and none recruited for this study are diabetic. However, our observation indicates that majority of the subjects are obese which might be due to unhealthy eating habits as seen among market women and a high level of sedentary lifestyle lacking exercises as reported in the Niger Delta region of Nigeria [22]. Though, some studies shows that women in developing countries largely experience raised BMI as a sign of beauty and affluence, which contributes to obesity [23].

The mean values of BMI, WHR and leptin levels significantly increases with age as observed in this study. This may be due to physiological weight gain that result from increase in body fat [24]. This finding agrees with the report of Low *et al.*[23], which stated that body weight of women increases with age, and this weight gain might contribute to obesity and other complicated factors. The study also revealed a positive correlation between leptin and BMI, WHR implying that these anthropometric parameters increases with the levels of leptin. Hence, these parameters be assessed in women of reproductive age, in order to provide some useful health information for women's health.

#### **5. CONCLUSION**

This study evaluated the levels of leptin and some anthropometric parameters in women of reproductive age in Port Harcourt. The mean levels of BMI and WHR of the subjects were above the reference range while the leptin levels were within reference ranges. These results indicate a prevalent obesity among the subjects, but not necessarily as a results of raised blood sugar. The results also show that the BMI and WHR all correlated positively with the leptin level, indicating that leptin may affect these anthropometric parameters. Our findings also show that BMI, WHR and Leptin values increased with age. Hence, the need to be assessed in women of reproductive age, as they may provide some useful health information for maintaining a better health status of women within this age brackets.

#### **ETHICAL APPROVAL**

Ethical approval was obtained from the Health Research and Ethics Committee, Rivers State Hospitals Management Board, Rivers State referenced RSHMB/RSHREC/2022/030.

## REFERENCES

Comment [m10]: Choose one style of referencing.

1. Egbe EO, Asuquo OA, Ekwere EO, Olufemi F, & Ohwovoriole AE. Assessment of anthropometric indices among residents of Calabar, South-East Nigeria. *Indian Journal of Endocrinology and Metabolism*. 2014; 18(3): 386-389.
2. Karakaş S, Tellioglu A, Dişçigil G, Karul AB & Türkmen M . Relationships between anthropometric measurements, leptin and IGF-1 levels in Turkish healthy newborns. *Journal of Clinical and Experimental Investigations*. 2015; 6(3): 214-219.
3. Ramos-Lobo AM & Donato JJ. The role of leptin in health and disease. *Temperature (Austin)*. 2017; 4(3): 258-291.
4. Cernea S, Both E, Huţanu A, Şular FL & Roiban AL. Correlations of serum leptin and leptin resistance with depression and anxiety in patients with type 2 diabetes. *Psychiatry Clinical Neuroscience*. 2019; 73(12): 745-753.
5. Cernea S, Both E. & Fodor A. The association of anthropometric parameters with markers of insulin and leptin secretion and resistance in type 2 diabetes mellitus. *Revista Romana de Medicina de Laborator*. 2020; 28(3): 299-314.
6. Casadei K & Kiel J. Anthropometric Measurement. In: *StatPearls [Internet]*. Treasure Island (FL): StatPearls Publishing. 2022. PMID: 30726000.
7. Akpınar E, Bashan I, Bozdemir N & Saatci E. Which is the best anthropometric technique to identify obesity: body mass index, waist circumference or waist-hip ratio? *Coll Antropol*. 2007;31(2):387-93. PMID: 17847914.
8. Lee BJ & Yim MH. Comparison of anthropometric and body composition indices in the identification of metabolic risk factors. *Sci Rep* . 2021;11: 9931-9934.
9. Chigbu CO & Aja LO. Obesity in Pregnancy in south east Nigeria. *Annual Medical Health Science Research*. 2011; 2(1): 1 - 6.
10. Crowther NJ & Norris SA. The Current Waist Circumference Cut Point Used for the Diagnosis of Metabolic Syndrome in Sub-Saharan African Women Is Not Appropriate. *PLoS ONE*. 2012; 7(11): e48883. doi:10.1371/journal.pone.0048883
11. Ross R, Neeland IJ, Yamashita S, Shai I, Seidell J, Magni P, Santos RD, Arsenault B, Cuevas A, Hu FB, Griffin BA, Zambon A, Barter P, Fruchart JC, Eckel RH, Matsuzawa Y, Després JP. Waist circumference as a vital sign in clinical practice: a Consensus Statement from the IAS and ICCR Working Group on Visceral Obesity. *Nat Rev Endocrinol*. 2020;16(3):177-189. doi: 10.1038/s41574-019-0310-7.
12. Guerra RS, Amaral TF, Marques EA, Mota J, Restivo MT. Anatomical location for waist circumference measurement in older adults: a preliminary study. *Nutr Hosp*. 2012; 27(5):1554-61. doi: 10.3305/nh.2012.27.5.5922. PMID: 23478705.
13. Djiolo F, Megnigbeto OA, De Souza J, Takpara I, Santos P & Alihonou E. Influence of Maternal weight on Pregnancy outcome in Cotonou (Benin). *Journal of Gynecology, Obstetrics Biological Reproduction*. 2002; 31(3): 243-247.
14. Kelechi AM. Perceived health implications of infertility among women of reproductive age in Port Harcourt metropolis of Rivers State. *International Journal of Innovative Medicine & Medicinal Plants Research*. 2022; 10(1), 18-26.
15. Ononamadu CJ, Ezekwesili CN, Onyeukwu OF, Umeogaju UF, Ezeigwe OC, & Ihegboro GO. Comparative analysis of anthropometric indices of obesity as correlates and potential predictors of risk for hypertension and pre hypertension in a population in Nigeria. *Cardiovascular Journal of Africa*. 2017; 28(2): 92–99.

16. Lee JW, Lim NK, Back TH, Park SH & Park HY. Anthropometric indices as predictors of hypertension among men and women aged 40–69 years in the Korean population: The Korean genome and epidemiology study. *British Medical College of Public Health*, 2015;15: 140–146.
17. World Health Organization (WHO). Waist circumference and waist–hip ratio: report of a WHO expert consultation, Geneva: World Health Organization.2011.
18. Ghose B. Nutritional transition in South Asia: The emergence of non-communicable chronic diseases. *F1000 Research*. 2015; 4(8): 28-48.
19. Okoh M. Socio-Demographic correlates of overweight and obesity among women of reproductive age in Nigeria. *African Journal of Reproductive Health*.2013; 17 (4): 66-76.
20. Onyeji GN. & Sanusi RA. Prevalence of overweight and obesity among women of reproductive age in south-east Nigeria. *Nigerian Journal of Nutritional Sciences*. 2018; 39(2): 73-81.
21. Tagbo SO, Abebe D & Oguoma VM. Overweight and obesity among non-pregnant women of reproductive age in Nigeria: Findings from the 2008-2018 Nigerian Demographic and Health Survey. *Public Health*. 2021; 198: 348-357.
22. Alikor CA & Nwafor CE. The prevalence and Predictors of generalised obesity in a rural farming community in the Niger Delta Region of Nigeria. *The Nigerian Health Journal*, 2016; 16(3): 1-10.
23. Low S, Chin MC & Deurenberg-Yap M. Review on epidemic of obesity. *Annals Academy of Medicine Singapore*, 2009; 38(1): 57-59.
24. Rezende FAC, Ribeiro AQ, Priore SE & Franceschini SCC. Anthropometric differences related to genders and age in the elderly. *Nutricion Hospitalaria*, 2015; 32(2), 757-764.

UNDER PEER REVIEW

