

Blue Carbon Ecosystem Changes and Ethnobotany Values of Indian Sundarbans; in Anthropocene

ABSTRACT

People in the inter-tidal zones of the Ganga-Brahmaputra-Meghanna Delta regions are reliant on mangroves for food, medicine, protection, and livelihood. The present Anthropocene epoch has a deteriorated impact on people's societal, fiscal, and demography due to the climate change loss of agriculture and land use. The mangrove vegetation of Indian Sundarbans is declining in the South 24 Parganna district, West Bengal, the world's largest estuarine forest, and the UNESCO Heritage, site.

Remote sensing (RS) and geographic information systems (GIS) techniques were used to download data of the Landsat 8 OLI from USGS/GLOVIS of SOUTH 24 Parganas, West Bengal and analysed by ERDAS IMAGINE applications for speculation of the land use/land cover dynamics changes from the year 2015 to 2020. Ground truthing was conducted by site visit and medicinal use of plants is collected from local people or from various literatures to gain knowledge of the sustenance of SDG-3 and SDG-15 of life in the Sundarbans.

The distribution of various mangroves was identified in the core, buffer and manipulation zones of the Sundarbans Biosphere Reserve, their stratification along the cross-section, zoning, and use for the stakeholders and as folk medicines. The mangrove vegetation is dwindling along the coasts of the South 24 Parganas. There is subsidence, emergence and submergence of land in the archipelago of Islands. The mangrove loss has been transformed into settlements or agricultural land. Recommendations are made to sustain the inter-tidal vegetation.

Keywords: GBM Delta, Sundarbans, Mangroves, GIS, South-24-Parganas, Ethnobotany

1.0 Introduction:

The east coast of India with large deltas is undergoing sinking, shrinking and subsidence of deltas, sea level rise (SLR), sedimentation, coastal erosion, and tidal inundations. Mangroves, the halophytic grow in the transitional interface between sea and terrestrial. They are the tidal influencing areas comprising standalone flora (epiphytes, shrubs, ground ferns, grasses and trees) and typical brackish water fauna. This unique ecological assemblage of halophytes is found in Sundarbans in South 24 Parganas of West Bengal, housed in the southern zone of the Ganga-Brahmaputra-Meghanna (GBM) delta. The Sundarbans act as a transition zone between the land and the ocean. It was a tiger reserve in 1973 declared as a wild sanctuary in 1977 further a national park in 1984. This mangrove ecosystem is of late Cretaceous origin [146 - 100 Mya]. Its expanse is ever-changing with the transgression and retrogression of the Bay of Bengal coast. The evergreen blue carbon ecosystem is under threat due to Anthropocene stress, climate change and dwindling fast and not compatible with Sustainable development goals, SDG 3 and SDG-15, (Rahman et al., 2020^[1], Bera et al., 2022^[2], Banerjee et al., 2023^[3]).

1.1 Anthropocene: Homo sapiens are governing the geo-bio-hydrosphere including the space beyond Earth's atmosphere. Population growth has extra-fragmented the land, from states to

districts grounded on nature's conformation in the Anthropocene (the human epoch) probably from 1950 and a larger scale from its golden spike period (1980). Presently, sustaining human life has urged for an incremental menace to humans, biodiversity, ecology, and the environment due to population density, modernization, urbanization, Industrialisation, and land use strategies, (Cruzen et al., 2000^[4], Zalasiewicz et al., 2016^[5], Mishra et al, 2020^[6], Waters et al., 2023^[7]).

1.2 Sustainable Development Goals: The SDGs 1-SDG 17 were accepted by India as one of the members out of 193 members of the United States on 15th Sept. 2015. SDG 3 demands healthy survival and encourages well-being for all genders from birth to death whereas SDG 15 promises to guard, reestablish and endorse sustainable usage of terrestrial ecosystems. The importance is emphasized sustainably dealing forests, combating desertification, and halting with setbacks of degraded biodiversity loss, (Gera et al., 2018^[8], Hassani et al, 2021^[9], Mishra et al., 2023^[10]).

1.3 Mangroves and their associates: Mangroves are halophytic plants that grow in the transition zones in coastal low-lying intertidal zones comprising aerophytes, shrubs, coastal ferns, grasses and trees. The Indian coast covers 4992 sq. km of mangroves out of 713789 km² global figure. <https://geographyhost.com/forest-cover-in-india-important-statistics/>. The Sundarbans (India and Bangladesh), the archipelago of islands, are housed in the Ganga Brahmaputra Meghna delta (GBMD). The largest mangrove delta is housed in about 10200 km² (4200 km² in India and the rest in Bangladesh) out of which about 2112.11 km² of India's lower Gangetic delta (FSI, 2019^[11] and <https://sundarbanresidency.com/history.html>). Deep Mangrove covers 257.10 sq. km in the South 24 Parganas. The well-diversified vegetation has developed certain morphological, biological, physiological, and ecological common adaptability to thrive along tropical coasts and creeks having climatological, biological, ecological, and economic importance. Sundarbans Wetland has an International Status, ranking 27th (Ramsar Site no. 2370). (Kuenzer et al. 2011^[12], Chakrabarti, 2015^[13], Das et al. 2023^[14]).

1.4 Blue carbon: The coastal system consisting of salt marshes, plankton, and seagrass ecosystems, efficiently stores carbon. They maintain the carbon balance and are called blue carbon. The blue carbon systems shield the coast from erosion, augment accretion, and regulate oceanic water standards, marine life and forestry. Presently this ecosystem is in jeopardy due to its resource over-exploitation, pollution, and anthropogenic intervention in its geomorphology. To sustain the role and maintain the concentration of CO₂ as per SDG-13. It warrants devoted insistence to restore the coastal blue carbon ecosystem to convert as a lasting carbon sink and to reduce carbon sequestrations. iucn.org/resources/issues-brief/blue-carbon. MNG and MNA ecosystems can store and sequester huge quantities of blue carbon (Corg) but are area-specific. The CO₂ emissions through coastal vegetation systems bear only 0.2% of total global CO₂ emissions but about 18% of CO₂ emissions from the tropical coastal ocean (Alongi 2020^[40])

1.5 STUDY AREA: The Population of 24 Parganas (S) was 1.38 million in 1951 whereas the present population has risen to 9.022 million only in south 24 Parganas (North 24 Parganas not taken), (Mandal 2020^[42], Census India, 2011^[43]). During the Meghalayan period (≈4.1k YBP) of the Holocene epoch, there were the Sundarbans mangroves. The Sundarbans, the largest mangrove, house between 88°55' E to 89° E and 21°30' N to 23°30' N, i.e., between the

Hooghly River (Kolkata) to Baleswar River (Khulna). The Indian mangroves have an area of about 3800 km². Sundarbans Biosphere Reserve (SBR) of India is located within 21°32'–22°40'N and 88°05'–89°51'E in districts of south 24- Parganas. The present study area is within 21°32'–22°14'N and 88°10'–89°05'E. Indian SBR has an area of about 2,400 km².




Fig 1: The Index map of the South 24 Parganas district in West Bengal part of GBM delta




Fig 2 (a): Sundari plant in Hooghly estuary (b): Knee roots in Passur or Pitamari (*Xylocarpus mekongensis*) mangroves (c) The Hental in Matla River in S-24-Parganas (d) Hental forests

The 'Sundarbans' are named after the main mangrove species 'Sundari' (*Heritiera fomes*) which is endangered due to cutting, and top-dying diseases. The other attraction of Sundarbans is the Royal Bengal tiger (*Panthera Tigris* ssp. *Tigris*), which is endangered now, (Dutta et al., 2017^[15]). The irrigation demand in Sundarbans is about 2784 MCum. The waterbodies (small or large) are 70000 in number and supplemented by 8,000 shallow tube wells that cannot cater for the demand, (Das et al., 2015^[16]). Food security is at stake for 87.5% of the low-income people in Sundarbans. Present changes in climatic, unsafe food security warrant an increase in the irrigation network, (Fig 2 a,b,c,d).

2.0.0 Review of Literature:

The 24-Parganas was established on 15th July 1757 by Mir Jafar, the first Nawab of Bengal. The South 24 Parganas was formed on 1st March 1986. Which was later bifurcated into subdivisions- (Alipore and Diamond Harbour) and divided into 30 blocks (GoWB and South 24 Parganas website^[47]).

The Multidimensional Rural Poverty Index Score of the South 24 Parganas as per NITI Ayog, 2023 that in the year 2015-16 and 2019-21 were 0.116 and 0.064 respectively and in urban areas were 0.053 and 0.022 respectively. (Where Multidimensional Poverty Index (MPI) = of India was 0.117 in 2015-16 and 0.066 in 2019-2021. (NITI Ayog, TATA-2023^[18])

Mangroves are developed in ≈ 150000 sq. km globally., (Kumar et al., 2022^[17]). India shares 4975 km², (0.15% of the globe) along the country's east, west, and AN Islands. Coasts. The largest mangroves in West Bengal cover 42.45%, followed by Gujarat (23.66%) (FSI, 2019^[11]). Indian Sundarbans have enumerated 27 pure mangrove species out of 46 true global and 40 mangrove species, while the east coast of India has 27 species, while the west coast has 27 species (Sreelekshmi et al. 2020,^[19] Nandi et al., 2021^[20], Chanda et al., 2023^[21]). MSL is a proxy for mangrove area delineation), (Payo et al. 2016^[22], Ghosh et al., 2018^[23])

Geomorphologic study of Sundarbans and south Parganas is difficult for ground survey due to inaccessibility and tough terrain. Remote sensing (RS) is one of the proper tools for mapping and monitoring inaccessible topography, and natural resources (Nandy et al., 2011^[24]; Sahana et al., 2015^[25]; Kushwaha et al., 2018^[26]; 2020, Thakur et al., 2021^[27], Mishra et al., 2023^[28]). The on-ground survey is hard, time-consuming, and laborious in mangrove tidal swamps, Remote sensing (RS) methodology can be conducted for the mapping and monitoring of the ground truthing and up-to-date information, (Giri et al., 2014^[29], Mishra et al., 2021^[30], Chanda et al., 2023^[21]).

RS techniques have the potential to recognize, map, and monitor changes in the mangrove and its associates in coastal areas, using hyper-spectral satellite images and information for digital image processing to classify mangroves bowls, (Heenkenda et al., 2014^[31], Datta et al., 2022^[32], Doodee et al. 2023^[33], Sudirman et al., 2023^[34]). Landsat TM, SPOT XS, IRS LISS-III, CARTOSAT-1 (2.5m), and Sentinel-2, are common sensors used to map mangroves. Mangrove species identification is helpful in high-resolution multispectral images rather than in coarser-resolution imageries (Satyanarayana et al., 2011^[35], Jia et al., 2016^[36], Nandi et al., 2021^[20]).

The effective management of a holistic blue carbon ecosystem can refurbish the strategy to bridge the inmates to protect stakeholders' risk from deterioration of ecology, community, lifeline, society, and economy. There is a lag between coordination between its stakeholders,

achieving suitable restoration goals and prescribing SOP (standard operating procedures), which are essential for the restoration of degraded mangrove areas.

2.1 Objectives:

Since 2016, there have been few literature studies against land use change estimation literature available. The objectives of the present geospatial study to investigate:

1. To access the multidimensional ethnobotanical folk drugs of the aboriginals of the area.
2. To speculate the functions of Sundarbans ecology, the current changes in the land use (LU) of South 24 Parganas in the last eight years (2015-2022).
3. The impact of the mangrove ecosystem on its stakeholders in the last two decades.

3.1 Ganga-Brahmaputra Meghana (GBM) Delta:

The Ganga Brahmaputra Meghana (GBM) deltaic canopy covers 1.7 million km² and covers two nations, India and Bangladesh covering an area of about 100000sq km. GBM basin covers six states, i.e., India, Bangladesh, Bhutan, China, Myanmar and Nepal. The GBM delta is the world's largest and wildest budding delta of Asia, which is formed by the tectonic rift and drift of the Indian Plate with the Eurasian Plate. The 355km long delta runs parallel to the Bay of Bengal (BoB) coast. It is shaped by the course of the conjoint deposition of sediment through GBM rivers (~1 BT/annum). The Ganga R. only carries about 150 to 590 MT/Y, the Brahmaputra 135 to 615 MT/y and the Meghana (6-12 MT/annum) have a decreasing trend (data 1960-2008). The ephemeral rivers and the longshore drift of about 260km long swampy coast have an average annual subsidence of 3-7mm/year (Steckler 2021, NASA [37]). The delta is fertile and has agricultural potential for India and Bangladesh, (BD) Rahman et al., 2018^[38], Lovelock 2019^[39], Alongi 2020^[40], Mishra et al., 2017^[41]).




Fig 3(a): Demographic study of South 24 Parganas Fig 3(b): The decadal population map of South 24 Parganas (source modified: <https://s24pgs.gov.in/assets/webdoc/mediauplo>)

3.2 South 24 Pargana (West Bengal)

The south 24 Parganas is a district (formed March 1st 1986) adjacent to the Cosmopolis, the Kolkata city of area 9960 km², comprising 8.153 million people (2011 Census). The district falls under Survey of India Topo Sheet No. 79B/2 to 6, 79B/10 to 12, 79B/15 and 16, 79C/1 to 6, 79C/9, and C/10. The lower part is housed in the Sundarbans, the largest mangroves of the globe. The connectivity is lacking due to a highly anastomosed drainage network, hilly land, and mangroves. The district is fed by the rivers Matla (wide estuary), the Raj

Mangal (tidal estuary), the Saptamukhi (tidal estuarine), the Hooghly (260km long, treated as holy Ganga), the Vidyadhari (estuarine), the Thakuran (called Jamira, tidal estuarine) and the Piyali (a connector between Bidyadhari and Matla).




Fig 4(a): GISTerrain map 24 Parganas (S), Fig 4(b): The soil map of 24 Parganas (S) Rivers (Source: GoWB; Dist. Survey Report -2018) Fig (c): Zoning of 24 Parganas (S) (Source: The World Bank, the IWA modified)

The district is trifurcated as (i) The marshy Sundarbans (Zone I) (ii) The Non-Sundarbans (countryside) (Zone II), and (iii) The Urban. (Zone III). Zone III are the urban and sub-urban in NW comprising deltaic plains of the Ganges starting from Farakka. Zone II (NE and Mid-west) lower delta is composed of anastomosis of 1st-order drainages emerging the 2nd-order distributaries by bifurcation within both mangrove swamps, Levees, and marshes with mangrove associates. The Zone I areas are the coastal belt consisting of creeks and Sundarbans with pure MNG. Irrespective of zones, the entire land is highly fertile and has high yields of rice, timber, fish, honey, sugarcane, beetle nuts etc, (Bagchi 2017[44]). Indian Sundarbans in the phytosociological study had five zones but presently transformed into four zones (loss of a few species of mangroves). It is due to the impact of Tides (Low and High),

Regional Sea level changes (RSLR), the impact of cyclones and human interventions, between 2017-2021(Chowdhury et al, 2023[45]). The Meghalayan prehistoric livelihood and habitation pattern has changed during the Anthropocene.

The south 24 Parganas is disaster-wise very sensitive. Cyclones with a return period of 4.62 years/cyclone, the flood of 22 events, average lightning days/ year 48 days, cold wave days of 20 days/year that confronted between 1969 to 2019. The av. rainfall of 1750 mm/year, and the maximum temperature of 48°C.(Edmonds et al., 2022^[46], GoWB Survey Report 2018^[47]).

Sundarbans; The world's largest mangroves:

Sundarbans have a coastal length of 260 km along the shoreline with *Rhizophora* mangrove species (largest Sundari), mangrove associates, and saltwater swamps formed at the lower Ganges Delta, https://s24pgs.gov.in/historical_background. Sundarbans halophytes are under threat/endangered and are included under the Red List of Ecosystems (Sievers et al., 2020^[48]). The evergreen canopy of the Sundarbans accommodates halophytes like the Sundari, Kakra, Keora, Garan, Hental, Golpata, Dhundul, Passur, Garjan varieties, etc. The distributaries small or large like Malta, the Hatania and Doania, the Go Saba, the Piyali, the Raiman gal, the saptamukhi, and the Thakuran, and the muddy islands such as Sagar, Ghoramara, Henery, and the Locha Achara Islands situated in the district (Chakrabarti et al, 2015^[13]). Species like *rotundifolia*, *Aegialitis*, *Heritiera fomes*, *S. Griffith*, and *Sonneratia apetala* have turned up endemic due to anthropogenic stress, westerly tectonic uplift, subsidence (3-7 mm/y). The mangroves protect the inmate's livelihood and its ecosystem from devastating floods, tides and cyclonic storms. In the 21st century, the cyclones had an impact on the south 24 Parganas but moderated by the mangrove ecosystem. They are Gonu (2007); Sidr (2007); Phailin (2013), Hudhud (2014), Mora (2017), Fani (2019), Amphan (2020); YAAS (2021); Sitrang (2022), and Mocha (2023)(Mishra et al, 2022^[49])

Indian Sundarbans:

The Indian Sundarbans are part of the GBM subaqueous delta covering 19 blocks out of 30 blocks. The Indian fabric of mangroves, a complex ecoregion had 6402 sq. km. in 2001 which has shrunk to 6385 sq. km in 2009. It was having erosion of 64.16 sq km and accretion of 20.12 sq km resulting in a balancing erosion of 44.04 sq km in a decade, (Ghosh et al., 2015^[49]). Near the coast, the forest enters the mangrove swamp; which shelters many desolate animals, brackish water species, and amphibians like crocodiles. It is the last zoonotic sanctuary of the Royal Bengal tigers (≈ 400 in number) and 30000 spotted deer. The rare bovines existing are sheep (Garole breeds), Muscovy ducks (China Hans), etc.

Ecosystem Services:

The fiscal and livelihood services obtained from Sundar Ban mangroves are added by fishing nurseries, aquaculture, fuel for the poor, honey, traditional medicines, etc. In addition, mutation of the forest to aquaculture and agriculture and, the construction of jetties and harbours to accommodate the increasing population. South 24 Parganas is enrolled as "The UNESCO World Heritage" list and named "The Sundarbans and Sundarbans National Park" respectively.

Sundarbans recorded various medicinal plants such as 17 types of MNG (true mangroves), 24 types of MNA (Mangrove Associates), 57 varieties of NMG (Non-Mangrove terrestrial

species), 13 varieties of Fabaceae, 5 species of Lamiaceae and Compositae (Islum et al, 2022^[50]), *Mangrove Species in Sundarbans: dominant is Heritiera (fomes or littoral, Sundari tree), Avicennia, Xylocarpus, Sonneratia, Bruguiera, Cereops, Aegicera, and Rhizophora. Nipa, a palm adapted to estuaries and mangroves, line the channels).*

Sundarbans Biosphere Reserve (SBR):

People in the SNR area are financially backward and are directly reliant on bit-resources like fish, crabs, prawns, timber, honey, and fuel wood from SBR. This eco-region has high yields, from land, rivers and diverse biodiversity. The weather remains humid and the Indian summer monsoon is active from June to Sept. with an average annual rainfall of 1750mm, av. Maximum Temperature is 35 °C and water bodies are allochthonous types, tide-dominated mangrove wetlands. The mangrove is in an estuarine, funnel-shaped linear tidal mud flats with anastomosed channels falling in BoB. Tidal level in high tides also varies from 3 to 6.5 m seasonally and water pH from 7.2 to 7.9. 24 numbers. The heart of the Sundarbans biosphere reserve has dwindled due to erosion, submergence and human over-exploitation of land, vegetation, and water resources.

The core area of SBR is restricted to human access for wood, honey, fishing and maintaining a tranquil environment for the endangered flora, fauna, avifauna and aquafauna. The buffer areas are admissible for honey, folk medicine collection, fishing, and limited dry wood cutting. Poaching and theft of forest products are prohibited from being protected by armed forest guards on patrol or in motorised boats. There is no nocturnal trespass within the river/forest. Forest bit houses and camps are set at vulnerable positions within the park. Location-wise the distribution of flora is in **Table 1**.

Table 1: The dominating MNG, MNA and NMG species found in Sundarbans

#	Location of the place	Local name	Botanical Name
1	<i>Estuarine zone, Tidal zone, banks, River mouths; swamps new sedimentation; sub-aqueous soil (Highly saline tolerant)</i>	Jatbaen; Pairabaen; Kalabaen Kripa Garia Tora	Avicennia officinalis Avicennia alba Avicennia marina Lumnitzera racemosa Kandelia candel Aegialitis rotundifolia
2	<i>Low salinity, Tidal current passing; channels in highland, Creeks in Mid Estuarine zone</i>	Garjan Goran Math Goran Bakul Keora	Rhizophora apiculata Ceriops Decandra Ceriops Tagal Bruguiera cylindrica Sonneratia petala
3	<i>Inner estuarine Zone: river inflow, less saline; elevated inland, Compact soil</i>	Genwa Kankara Khalsi Ora Hental	Excoecaria agallocha Bruguiera gymnorhiza Aegiceras corniculatum Sonneratia caseolaris Phoenix paludosa

4	Rare(R), endemic (E), Sporadic (S) restricted to specific salinity	Amur (E) Dhundul (E) Passur (E) Sundari(R) Golpata (R) Palm-swamps (S)	Agaliadomestica Xylocarpus granatum Xyclocarpusmekongensis Heritiera fomes Nypafruticans Palm-swamps
5	Freshwater Mangroves, dense and discontinuous, bushy and dwarf, even bushy	Hargoza Ban Lebu LataSundari	Acanthus ilicifolius Merope angulates Brownlowiaterasa

Ethno-medicinal utility of Sundarbans mangroves:

Studies reveal the regular use of proper doses of pneumatophores, twigs, leaves, and barks are taken for medicinal use against diabetes, hepatitis, gastrointestinal disorders, goitre, piles, and skin ailments, etc. as naturopathic drugs by traditional Vaidya (Doctors). Various reports say the common and popularly available MNG and MNA mangrove varieties available in Indian Sundarbans possess antiviral, antibacterial, antifeedant antioxidant, antileukemia, antidiarrheal, and antitumor pharmacological properties and also behave excellently as antioxidants, anti-microbial, antihyperglycemic, antinociceptive, and anti-malignant. Phytochemical studies revealed they contain chemical ingredients i.e., saponins, glycosides, alkaloids, steroid tannins, flavonoids, gums, phytosterols, and antidiabetic, (Islam et al., 2022^[51]) [Table 2].

Table 2: Major mangrove species used as ethnobotany

Local name	Botanical name	Part used	Mode of attempt	Disease attempted /uses	Sources
Sundari	<i>Heritiera fomes</i> :	Leaves, seeds,	Extraction by decoction	Diarrhoea, dysentery, colic, acidity, indigestion, constipation, stomach ache, bloating, lack of appetite and Gastrointestinal disorders	
		woods,	Powder	Piles, Laxatives	
		Stem bark	Paste	Eczema, abscess, boils, aches, rash, infections, scabies, itch, dermatitis, sores, scar, and warts	Mehmud et al, 2014 ^[53]
		Bark	decoction	Diabetes & Goitre	
		Twig	Toothbrush	Toothache; oral infection	
Indian mangrove 8 types	<i>Avicennia (Officinalis L. or Marina)</i>	Leaves roots & barks	Paste or decoction	Injuries or inflammation (alkaloids); cancer, diabetes, malaria, rheumatism, asthma, smallpox and ulcer	Patra et al., 2011 ^[54] ; Namaji et al, 2013 ^[55]
Acanthaceae, JATbaen (such 8 varieties)	<i>Avicennia officinalis; Avicennia marina</i>	Leaves roots, flowers bark, twigs	Diabetic and diuretic, immunity developer	rheumatism, asthma, tumour, dyspepsia, paralysis	Thatoi et al, 2016 ^[56]
Hental	Phoenix paludosa; Arecaceae	Upper soft part	Bark	To Control Cough	
Dhundul (tree)	<i>Xylocarpus granatum</i>	Bark	paste	dysentery, diarrhoea and other abdominal troubles, Febrifuges.	

		Seed ash	seed ash+sulfur&coconut oil	Allergy; Itching	Alamgir et al, 2007 ^[57]
		Fruit	Direct or maceration	Antiinflammatory,antidiarrhoeal,	
Caw Phal (Climber)	Sarcolobusglobosus	Fruit, leaves	Extract or maceration	antimicrobial activity, rheumatism, dengue	Alamgir et al, 2007 ^[57]
Math Garan	Ceriops tagal	Bark and wood	Extract and treat wood	Antioxidant and anticorrosive, Construction material, dying and tanning	Shamsuzzaman et al., 2021 ^[58]
Genwa (blind your eye mangrove)	Excoecariaagallocha	Leaves, flowers, twigs	Latex is poisonous so	antioxidant, antimicrobial, anti-inflammatory, analgesic, antiulcer, anticancer, anti-reverse,anti-filarial, transcripts, antihistamine-release, DNA damage protective, antidiabetic, antitumor	Mondal et al, 2016 ^[59]
Garjan	Rhizophora mucronata	Leaves, flowers twigs	Extract or maceration	Antidiabetic and antioxidant activity; antifungal activity	Adhikari et al, 2016 ^[60]
Keora	Sonneratia apetala	Fruit, leaves, seed	Seed power/ extraction	Antibacterial, Anti-Diarrhoeal, Analgesic,Antioxidant, Cytotoxic Activities; asthmaulcers, febrifuge, swellings, sprains, bleeding, haemorrhages, piles	Hossain et al., 2017 ^[61]
Kankra	Bruguieragymnorhiza	bark; leaves	Extract	abortifacient;Liver disorders	Sur et al, 2016 ^[62]
Khalsi (Odiya-Teluni)	Aegicerascorniculatum	Leaves, bark	decoction	Antidiabetic, , analgesic, asthma, Diarohea, and rheumatism.	Gurudeeban et al, 2012 ^[63]
Pasur	Xylocarpusmekongensis (3species)	seeds, fruits, stem bark, leaf, twigs	Seeds power/ extraction, decoction	Anti-oxidant, anticancer, anti-diabetic, antimicrobial, anti-malarial, antifeedant, neuro-protective diarrhoea, cholera, dysentery, fever, viral infections	Nabeelah Bibi et al, 2019 ^[64] ; Dey et al., 2021 ^[65]
Jele Goran	Ceriopsdecandra	Leaves	Extract	Antidiabetic, antimicrobial; inflammation, and cancer.	Mahmud et al., 2018 ^[66]
Tora	Ding Hou & Tora Senna Tora	Leaves	Extract	Antioxidant, antibacterial, anti-inflammatory; bronchitis, itches, rheumatism, ringworm, leprosy, dyspepsia, liver & heart disorders.	Rahman et al., 2023 ^[67]
Bhola	Hibiscus tiliaceus	Flowers	extract	Antibiotic;bronchitis, fevers, antihistaminic; coughs, ear infections, abscesses, postpartum disorders, skin diseases	Abdul Awal et al., 2016 ^[68]
Golpata	Nypafruticans	Fruit	Extract	land scaping, antidiabetics and antioxidant, less toxic;healing gout, seeds creamy flavour	Yossuf et al., 2015 ^[69]

The tree Sundari in Sundarbans, a major medicinal plant is gradually becoming “endangered” on the IUCN Red List.

Methods and methodology:

The present study has focused on investigating the changes in the mangrove forest area due to natural causes, and the hydrological regime due to substantial variations in topography,

salinity, erosion, tidal current inundations, and species species-specific plantation to sustain the MNG and MNA. Baseline information is collected for the pre-requisite for identifying, mapping, and estimation of mangroves by experienced survey staff, hydrology, geomorphology, the amplitude of tides, inundation area, and the existence of predators. Due to their inaccessibility and inundation, it is a herculean task to conduct field surveys within the tidal swampy mangrove forests with tigers, crocodiles etc.. So, a species-specific afforestation program is necessary for Sundarbans, (Nandy et al. 2011^[24]).

GIS Methodology to detect LULC changes:

The remote sensing (RS) technique has increasingly emerged as an emerging innovative tool for mapping and timely change of land use (mangroves), though downloaded RS data cannot replace the field survey data, the classification and the change detection. The RS claims the advantage of synoptic coverage, free/ low-cost, time-saving, accurate and repetitive coverage, during watershed, mangrove, coastal, and disaster management, (Crowley et al., 2020^[71], Kumar et al., 2021^[17] Mishra et al., 2022^[72]).


The LULC mapping plays a vital role in planning, managing and monitoring land utilization aspects against territorial developmental programs, forest policy-making and their enactment and monitoring aspects.

Data-Processing:


The Landsat 8 OLI data for continuous data for the year 2015 to 2022 of SOUTH 24 Parganas (downloaded either 2nd quarter of Nov. or the first quarter of Dec.) of the year. After completing the transfer from USGS/GLOVIS was extracted in a zip file and was opened in TIFF files of diverse bands in ERDAS IMAGINE. The chronological processes followed are layer stacking, subset imaging and supervised classification. Further, the steps are setting preferences (input and output directory), adding layers, inserting geometry, and identifying particular land features manually to have appropriate pixel accuracy. Then select the raster menu and select all signatures or classes. The Satellite image found five features- (i) Water bodies (ii) Agricultural land (iii) Vegetation (iv) Built and (v) Marshy land to have supervised classification and find the error data. To get an accurate supervised image decision rules can be tried one after other.

Present setup


MNG occurred in the study area from the years 2015-16, 2016-17, 2017-18, 2018-19, and 2019-20 (for five years). The cloud cover pictures were not considered and the error in processing within <5% is accepted. The classification method is applied for data analysis, the classification matrix is used to analyze urban growth. The classified images are interpreted to obtain several pixels for calculating area. Temporal analysis helps in studying the change detection for analyzing the decrease in the number of water and vegetation image pixels. In the process of five years of data analysis, we can be able to find out the major changes in terms of land surface /land cover, vegetation and many more then we can easily interpret the analysis process with appropriate calculations and graphs that how much area gradually changed in percentage (%) is given below.




a)




b)




c)



d)



e)



f)




Fig 5 (a) to (J): The year-wise areas for land use for mangroves, mangrove associates and inland taxa in Sundarbans, S-24-Parganas.

The result reveals that the Sundarbans area is populous. People have exploited the edges of the land for either settlements or agriculture at the cost of mangroves or mangrove Swamps, (Fig 5((a-f) and (Table 3)

Table 3: The land use and land cover of theclassified study area in % of South 24 Parganas

Class	2015-16	2016-17	2017-18	2018-19	2019-20
Classified	% of Area	% of Area	% of Area	% of Area	% of Area
Shallow water	2.4	2.2	1.6	1.0	2.1
Mangrove Forest	2.9	2.5	2.6	3.6	3.4
Mangrove swamp	10.2	9.9	9.6	9.1	9.4
Settlement Area	11.6	13.8	10.8	14.2	10.6
Waterlogged area	7.1	7.3	3.1	6.5	6.0
River bed	8.1	14.6	13.1	9.4	12.2
Land and scrub	16.1	12.5	17.5	11.9	17.7
Water (deep)	12.1	8.2	10.9	14.8	09.2
Not Classified	29.5	29	28.8	29.5	29.4

Considering the mangrove forest, The LULC map for Sundarbans in south 24 Parganas was constructed and the corresponding land utilisation changes were calculated.




Figure 6 a-d: Land use area changes in % for various features between the year 2015 and 2022

It is inferred from the satellite data study that one per cent of the study area has been converted from mangrove forests to mangrove swamps. The causes may be attributed to either increase in settlement area 5 (a to d).

Table 4: land use changes in % for various features in Sundarbans in South 24 Parganas, WB.

Class	2015	2022	% change (+/-)
Classified	% of Area	% of Area	
Mangrove Forest	4	3.0	(-) 1.0
Mangrove swamp	4	5	(+) 1
Settlement Area	7	24	(+) 17
Waterlogged area	7.1	6.0	(-) 1.1
Agricultural land	15	10.0	(-) 5
Land and scrub	7	7	0
Waterbody	16	2	(-)14
Not Classified	40	43	--

The structure of the result of the analysis indicates that within seven years the built-up area and the transformed water bodies (ponds) to agricultural land. have surged by (17%), the cause being a rise

in population. Due to the slamming of the extremely severe cyclonic storm Amphan 2020, there was great devastation to Sundarbans.




Fig 7: The various islands existing (1930) within Indian Sundarbans are downsized in 2022

The topographical maps of 1930 of the Survey of India have been compared with the 2022 present map. Some larger islands within the Indian Sundarbans have been submerged totally or partly like Sagar sands. This confirms that the Bay of Bengal water level is rising and retrograding the WB coast. The record reveals that 54 islands in the Indian Sundarbans are either emerging, vanishing or vulnerable to submergence (https://www.Sundarban-affairs-wb.in/home/page/islands_of_sundarban).

Table 5: The status of various Islands under the Indian Sundarbans, India

#	Status of Islands	Name of the Islands
1	Islands disappeared from the present map	Lohachara, Bedford, Kabasgadi and Suparibhanga etc.
2	Islands under slow disappearance (vulnerable islands) to erosion are	Sagar, Ghoramara, Dakshin Surendranagar, Mousuni, Namkhana, Sagar sands, Dhanchi, Dalhousie, Bulchery, Bhangaduni, Mousini Lothian, Baliakhal Lothian, and Jambudwip
3	Islands emerged (in 1970 and later submerged in 2010).	South Talpatti after the Bholia (1970) cyclone, on the territory line and uninhabitable.
4	Enlargement	Islands enlarging in eastern and northern margins of Indian Sundarbans Islands

Source: Mandal 2015^[73], www.sundarbanaffairswb.in/home/page/islands_of_sundarban

Spatial changes in the territory and outskirts are noticed in Sundarbans. The archipelago of islands mostly submerged and rarely emerged depending upon RSLR, littoral drift, and sediment deposition at the mouth of the estuaries. The area of the Islands was estimated from old SOI maps from the British Period and stated that Indian Sundarbans mangroves inclusive of the waterbodies and drainage channels in the area were 1773 (10064 Km²), in 1873 (7894 Km²), 1973 (4277 km²), and 2016 3576 Km² @ 5.63% excluding the area of the riverine channels. (Unikrishnan et al, 2007, Mishra et al, 2017, Hussain et al 2017).

Discussion

Mangroves grow along the sea-terrestrial interface between land and sea. The mangrove forests can be built up where diurnaltidal inundationby tidal waves occurs. The gigantic Indian Sundarbans formed over a subaerial, subaqueous and tidal GBM delta. They survive within the elevation of the alluvial fans, paleo deltas and the Pleistocene terraces. The unique heritage is housed in the fag end of the littoral drift or offshore current along the East Coast. The Himalayan rivers are ephemeral and carry huge amounts of sediment. The littoral sediment and the inland fluvial sediment get deposited along the flood plains, which act as a catalyst for the sustenance ofMNG and MNA.

Mangroves generally thrive in the most traumatic and unreceptive settings of highly saline, large tidal fluctuations, raised surface air temperature (SAT) and high humidity, strong gusty winds and silty anaerobic soil. These extremely tolerant plants and shrubs can protect stakeholders from erosion, adverse climate and cyclones, and there is insistent dwindling in the mangrove population. The various human services provided by these unique lifelinefloral kingdomurge for its sustainable existence from nature and human cruelty. Its degradationis far higher in Asia than in other tropical and subtropical regions.

Human-based challenges are overfishing, tourism, food, medicine, fuel, shelter, deforestation, and construction materials are mainly human-based. Nature-based forces are subsidence, submergence, climate change, meteorological extremes, and ownership of land holdings of the swamps and mangroves. The main redressal actions are people's awareness, education, economic growth, clear landownership, distinct plans and proposals, the PPP mode development, participatory forest management,

The present study aims to detect theLU changesfrom 1773 to 2016 and especially (2015–2020) of Indianmangrove zones using the RStechique,The diminishing trend in Indian mangroves and D/Cs is seen. The changes perceived are in mangrove swamps, and fallow lands, at the cost of waterbodies, mangrove swamps and lands covered with MNG and MNA.It is high time to protect our coastalwealtheither by legal interventions, people's participation, community involvement or PPP mode as one step to complywith SDG-15 i.e., life and land.

Regularly forming nurseries, provision for backwaterintermediate Inundation and species-specific afforestation are to be adopted to keep the mangrove forest sustainable. Baseline information about the rejuvenation of the existing mangroves is a criterion to identify, map, and estimatemangroves' status. The constraint duringthe sustenance of maintenance of mangroves is strenuous and inaccessibleto conduct field surveys amidstcoastal creeks and swamps mangrove forests.

The Sundarbansin South 24 Parganashavetried replantation programmes but are futilefor adequate knowledge, communication, and documentation forthe restoration of mangrove ecology.The national policy of food security (SDG-2) in changing climate, surged irrigation potential, and harnessing more water resources are essential. This is possible with morerainwater harvesting structures, rejuvenation and renovation of defunct and depleted drainage channels and recharging of UG aquifers are warranted.

SDG-15 can not be ignored as the protection, restoration and promotion of forests demand its sustainable usage of telluric ecosystems. Actions are tomanage the forest's sustainability,

fight against desertification, stop and reverse gear the degradation and cease biodiversity loss to adhere to SDG-5.

Mangroves, the ensembles as significant asylum to coastal/estuarine bio-diversity. They act as the bio-shields to extreme onshore events like gusty winds of cyclones, High tides, coastal erosion, high floods, and tsunami waves. These bio-shields never betray their ecosystem, zoonotic, sheltering avifauna, and biomass-dependant stakeholders and even add to the blue-carbon ecosystem, (ISFR 2021)

CONCLUSION:

Human activities have in recent years become recognised as a major force in shaping the biosphere. The land use/land cover pattern of a region is an outcome of natural and socioeconomic factors and their utilization by man in time and space. The land is becoming a scarce resource due to immense agricultural and demographic pressure. Remote sensing (RS) and geographic information systems (GIS) techniques provide effective tools for analyzing the land use dynamics of the region as well as for monitoring, mapping and management of natural resources. The objective of the present study is to analyze the land use/land cover dynamics from the year 2015 to 2020. This project is mainly to show the satellite image and how it has changed in five years.

In the last 5 years duration (2015–2020) mangrove coverage has declined significantly. The study also depicts that the total mangrove forest area has decreased in Sundarbans with an increase in water area. The study reveals that the S-24-Parganas have lands in swamps and creeks available for mangrove plantations. Urbanization and agriculture when linked to other land use/land cover features, have a high rate of conversion against other land usages.

The apocalyptic sea level rises in the study area, sediment-rich (instead of sediment paucity) and live delta under erratic climate change, global warming under changing sun earth geometry and anthropogenic attack to nature has already deteriorated ($\approx 75\%$) and cannot recoup. The present South 24 Parganas milieu cannot recuperate like the 19th-century biome but only can be maintained as it is today.

The district of South 24 Parganas is known to be the active delta of the Ganges, where the formation of the delta is still an ongoing process. The Sundarbans and its mangroves and their sustainability by adopting community contribution and spending social capital is the only successful technique to preserve UNESCO's World Heritage, the most active Delta and its vast bio-diversity of coastal lands, brackish water flora, fauna, avifauna, and aqua fauna.

Reference:

1. Rahman, MM, Ghosh T., Salehin M., Ghosh, A., Haque A., Hossain, MA., Das, S., Hazra S.. et al. (2020). Ganges-Brahmaputra-Meghna Delta, Bangladesh and India: A Transnational Mega-Delta. In: Nicholls, R., Adger, W., Hutton, C., Hanson, S. (eds) *Deltas in the Anthropocene*. Palgrave Macmillan, Cham. https://doi.org/10.1007/978-3-030-23517-8_2
2. Bera, B., Bhattacharjee, S., Sengupta, N. et al. Significant reduction of carbon stocks and changes in ecosystem service valuation of Indian Sundarbans. *Sci Rep* 12, 7809 (2022). <https://doi.org/10.1038/s41598-022-11716-5>
3. Banerjee S, Chanda A, Ghosh T, Cremin E, Renaud FG. A Qualitative Assessment of Natural and Anthropogenic Drivers of Risk to Sustainable Livelihoods in the Indian Sundarbans. *Sustainability*. 2023; 15(7):6146. <https://doi.org/10.3390/su15076146>

4. Crutzen, Paul J., and Eugene F. Stoermer. 2000. "The Anthropocene." *Global Change Newsletter* 41: 17–18.
5. Zalasiewicz J, Williams M, Waters CN, Barnosky AD, Palmesino J, et al. 2016. Scale and diversity of the physical technosphere: a geological perspective. *Anthr. Rev.* 4:9–22
6. Mishra S. P., July 2020; Human Evolution/Extinction up to Present Anthropocene:India; Journal of Shanghai Jiaotong University; JSJ.U-2222.14-F (1).pdf; ISSN:1007-1172; 16 (7); 115-133
7. Waters, C. N., Turner, S. D., Zalasiewicz, J., & Head, M. J. (2023). Candidate sites and other reference sections for the Global boundary Stratotype Section and Point of the Anthropocene series. *The Anthropocene Review*, 10(1), 3–24. <https://doi.org/10.1177/20530196221136422>
8. Gera R, Narwal R, Jain M, Taneja G, Gupta S. Sustainable Development Goals: Leveraging the Global Agenda for Driving Health Policy Reforms and Achieving Universal Health Coverage in India. *Indian J Community Med.* 2018 Oct-Dec;43(4):255-259.
9. Hassani H, Huang X, MacFeely S, Entezarian MR. Big Data and the United Nations Sustainable Development Goals (UN SDGs) at a Glance. *Big Data and Cognitive Computing.* 2021; 5(3):28. <https://doi.org/10.3390/bdcc5030028>
10. Mishra SP, Mohapatra SK., Sethi KC. (2023) The Values and Blue Carbon Ecosystem of the Chilika Lagoon through Ages, India. *Environ Sci Arch* 2(2):164-184
11. FSI 2019. Mangrove resources of the country. India State of Forest Report 2019. Forest Survey of India, Ministry of Environment, Forest and Climate Change, Government of India, Dehradun
12. Kuenzer C, Bluemel A, Gebhardt S, Quoc TV, Dech S. Remote Sensing of Mangrove Ecosystems: A Review. *Remote Sensing.* 2011; 3(5):878-928. <https://doi.org/10.3390/rs3050878>
13. Chakrabarti, P., Nag, S. 2015. Rivers of West Bengal: Changing Scenario. Geoinformatics and Remote Sensing Cell, Govt. of West Bengal, Kolkata: 265p.
14. Das I, Chanda A, Akhand A, Hazra S. Carbon Biogeochemistry of the Estuaries Adjoining the Indian Sundarbans Mangrove Ecosystem: A Review. *Life (Basel).* 2023;13(4):863. doi: 10.3390/life13040863
15. Dutta MK, Bianchi TS., Mukhopadhyay SK (2017) MangroveMethane Biogeochemistry in the Indian Sundarbans: [Frontiers in Marine Science](https://doi.org/10.3389/fmars.2017.00187) 4, DOI: [10.3389/fmars.2017.00187](https://doi.org/10.3389/fmars.2017.00187)
16. Das, S.; Bhadra, T.; Hazra, S., Water for Agriculture in a Vulnerable Delta: A Case Study of Indian Sundarbans, NASA Astrophysics Data System (ADS), 2015-12-01
17. Kumar D, Dhaloiya A, Nain AS, Sharma MP, Singh A. Prioritization of Watershed Using Remote Sensing and Geographic Information System. *Sustainability.* 2021; 13(16):9456. <https://doi.org/10.3390/su13169456>
18. NITI Ayog, (2023). National multidimensional poverty index India, a progress review 2023.
19. Sreelekshmi, S., Nandan, S.B., Kaimal, S.V., Radhakrishnan, C.K. and Suresh, V.R., 2020. Mangrove species diversity, stand structure and zonation pattern concerning environmental factors—A case study at Sundarbans delta, east coast of India. *Regional Studies in Marine Science*, 35, p. 101111
20. Nandy, S., Tamang M., Kushwaha SPS., (2021), Plant Diversity Assessment in Indian Sundarbans Mangroves: A Geoinformatics Approach. Ch-5, In book: Sundarbans Mangrove Systems, DOI: 10.1201/9781003083573-5
21. Chanda A, Akhand A. Challenges towards the Sustainability and Enhancement of the Indian Sundarban Mangrove's Blue Carbon Stock. *Life.* 2023; 13(8):1787. <https://doi.org/10.3390/life13081787>
22. Payo, A., Mukhopadhyay, A., Hazra, S. Ghosh, S. Ghosh, S. Brown, RJ, Nicholls, L. Bricheno, J. Wolf, S. Kay, A.N. Lázár, and A. Haque. 2016. Projected changes in the area of the Sundarbans mangrove forest in Bangladesh due to SLR by 2100. *Climatic Change:* 1–13. <https://doi.org/10.1007/s10584-016-1769-z>
23. Ghosh T., 2018. Dynamics of the Sundarbans Mangroves in Bangladesh Under Climate Change. Ch-26, from the book: *Ecosystem Services for Well-Being in Deltas*; R. J. Nicholls et al. (eds.), https://doi.org/10.1007/978-3-319-71093-8_26
24. Nandy, S., Kushwaha, S.P.S., Dadhwal, V.K., (2011). Forest degradation assessment in the upper catchment of the river Tons using remote sensing and GIS. *Ecological Indicators*, 11(2), pp. 509–513
25. Sahana, M., Sajjad, H., Ahmed, R. (2015). Assessing spatio-temporal health of forest cover using forest canopy density model and forest fragmentation approach in Sundarbans reserve forest, India. *Model. Earth Syst. Environ.* 1, 49, <https://doi.org/10.1007/s40808-015-0043-0>
26. Kushwaha, SPS, Nandy, S., Shah, M.A., Agarwal, R. and Mukhopadhyay, S., 2018. Forest cover monitoring and prediction in a Lesser Himalayan elephant landscape, *Current Sci.*, 115(3), 510–516

27. Thakur, S., Mondal, I., Bar, S., Nandi, S., Ghosh, PB., Das P., De, TK, (2021). Shoreline changes and their impact on the mangrove ecosystems of some islands of Indian Sundarbans, North-East coast of India, *Journal of Cleaner Production*, 284, 124764, <https://doi.org/10.1016/j.jclepro.2020.124764>.
28. Mishra, S.P., Mishra, S., Bera, D.K. (2023). Anthropocene Lithology and Hydropower search through North Eastern Ghats Hills, Odisha. *Water productivity journal*, 3(1): 75-104. DOI: 10.22034/wpj.2023.170350
29. Giri, S., Mukhopadhyay, A., Hazra, S., Mukherjee, S., Roy, D., Ghosh, S., Ghosh, T., Mitra, D., (2014). A study on abundance and distribution of mangrove species in Indian Sundarbans using RS technique. *J. of Coastal Cons.*, 18(4), 359–367
30. Mishra, SP., Barik, KK., Pattanaik, SK., (2021). The Vulnerability and Management to the Blue Carbon Ecosystem: Coastal Odisha, *International Journal of Lakes and Rivers.*, ISSN 0973-457014(1); 43-70.
31. Heenkenda MK, Joyce KE, Maier SW, Bartolo R. Mangrove Species Identification: Comparing WorldView-2 with Aerial Photographs. *Remote Sensing*. 2014; 6(7):6064-6088. <https://doi.org/10.3390/rs6076064>
32. Datta, D., Bairagi, M., Dey, M. et al. Spatially explicit estimation of soil organic carbon stock of an estuarine mangrove wetland of eastern India using elemental analysis and very-fine resolution satellite data. *Ecol Process* 11, 30 (2022). <https://doi.org/10.1186/s13717-022-00370-4>
33. Doodee MDD, Rughooputh SDDV, Jawaheer S., (2023). Remote sensing monitoring of mangrove growth rate at selected planted sites in Mauritius. *S Afr J Sci.*;119(1/2), DOI: 10.17159/sajs.2023/13716
34. Sudirman N., Salim, HL., Rustam, A, Ati1 RNA, Hermina R., Marini Y, Suryono1et DD, (2023). IOP Conf. Ser.: Earth Environ. Sci. 1148 012041, doi:10.1088/1755-1315/1148/1/012041
35. Behera S.,Mohamad K. A., Idris I., Husain M., Dahdouh F., Assessment of mangrove vegetation based on remote sensing and ground-truth measurements at Tumpat, Kelantan Delta, East Coast of Peninsular Malaysia, *Int. J. of RS, Sensing*,2011,32, 1635 – 1650,
36. Jia M, Liu M, Wang Z, Mao D, Ren C, Cui H. Evaluating the Effectiveness of Conservation on Mangroves: A Remote Sensing-Based Comparison for Two Adjacent Protected Areas in Shenzhen and Hong Kong, China. *Remote Sensing*. 2016; 8(8):627. <https://doi.org/10.3390/rs8080627>
37. Steckler, MS., Bar O.; t Jaman, H. ; Mondal, DR. ; Grall, C., et al., 2021. Recent measurements of subsidence in the Ganges-Brahmaputra Delta, Bangladesh. <https://ui.adsabs.harvard.edu/abs/2021EGUGA.23.6562S/abstract>
38. Rahman M., Dustegir M., Karim R., 10 authors, Akter M., (2018). Recent sediment flux to the Ganges-Brahmaputra-Meghna delta system. *The Science of The Total Environment* 643(1-2):1054-1064, DOI: 10.1016/j.scitotenv.2018.06.147
39. Lovelock CE. Duarte CM. 2019 Dimensions of Blue Carbon and Emerging Perspectives *Biol. Lett.*152018078120180781; <http://doi.org/10.1098/rsbl.2018.0781>
40. Alongi DM. Global Significance of Mangrove Blue Carbon in Climate Change Mitigation. *Sci*. 2020; 2(3):67. <https://doi.org/10.3390/sci2030067>
41. Mishra SP., (2017). Management of the sediment transported by the south Mahanadi Deltaic rivers to the Chilika lagoon. *Int. J. of Adv. Res.* 5 (Jun). 1005-1020]. DOI:Article DOI: 10.21474/IJAR01/4503
42. Mandal JK., 2020. The scenario of population growth South 24 Parganas, Dist West Bengal, India. *International Journal of Research in Social Sciences*, 10(4), 10-26
43. Census of India 2011- National Population Register & Socio-Economic and Caste Census.ppt.
44. Bagchi E., 2017. Development of Basic Infrastructure: An Analysis of South 24 Parganas District in WB, India. *Bulletin of Geogr. Socio-economic Series / No. 36* (2017): 33–60, DOI: 10.1515/bog-2017-0013
45. Chowdhury A, Naz A, Sharma SB, Dasgupta R. Changes in Salinity, Mangrove Community Ecology, and Organic Blue Carbon Stock in Response to Cyclones at Indian Sundarbans. *Life (Basel)*. 2023 Jul 11;13(7):1539. doi: 10.3390/life13071539.
46. Edmonds, C., Mehhta, M., Noy, I., Banik, P. (2021). The Climate-(Ir)Resilient Society of the Indian Sundarbans. In: Brears, R.C. (eds) *The Palgrave Handbook of Climate Resilient Societies*. Palgrave Macmillan, Cham. https://doi.org/10.1007/978-3-030-42462-6_95
47. Government of West Bengal, 2018. District survey report South 24-Parganas District; Sathi Planners Private Limited; Prepared As Per Ministry of Environment, Forest And Climate Change Notification S. O. 3611 (E),25th July 2018

48. Sievers, M., Chowdhury, MR., Adame, MF., Bhadury, P., Bhargava, R., Buelow, C., et al., 2020. Indian Sundarbans mangrove forest is considered endangered under the Red List of Ecosystems, but there is cause for optimism. *Biological Conser.*, 251, 108751
49. Mishra SP., Ojha AC, MishraS., Sahu DK., 2022. Cyclogenesis and Odisha Coast, the Hotbed. *Agriculture Association of Textile Chemical and Critical Reviews Journal* (2022) 71-85, DOI://doi.org/10.58321/AATCCReview.2022.10.01.85
50. Ghosh, A., Schmidt, S., Fickert, T., & Nusser, M. (2015). The Indian Sundarbans mangrove forests: History, utilization, conservation strategies and local perception. *Diversity*, 7, 149–169.
51. Islam ATMR, Hasan MM, Islam MT, Tanaka N. Ethnobotanical study of plants used by the Munda ethnic group living around the Sundarbans, the world's largest mangrove forest in southwestern Bangladesh. *J Ethnopharmacol.* 2022 Mar 1; 285:114853. doi: 10.1016/j.jep.2021.114853.
52. Mondal M., Paul S., Bhattacharya S., Biswas A., 2020. Micro-level Assessment of Rural Societal Vulnerability of Coastal Regions: An Insight into Sagar Island, West Bengal, India. *Sage, Asia-Pacific Journal of Rural Development* 30(1–2) 55–88, 2020, DOI: 10.1177/1018529120946230
53. Mahmud I, Islam MK, Saha S, Barman AK, Rahman Mmet.al., Pharmacological and Ethnomedicinal Overview of Heritiera fomes: Future Prospects. *Int Sch Res Notices.* 2014;2014:938543. doi: 10.1155/2014/938543
54. Patra, K.J., Dhal, K.N. and Thatoi, N.H. (2011) In Vitro Bioactivity and Phytochemical Screening of Suaeda maritima (Dumort): A Mangrove Associate from Bhitarkanika, India. *Asian Pacific Journal of Tropical Medicine*, 4, 727-734. [http://dx.doi.org/10.1016/S1995-7645\(11\)60182-X](http://dx.doi.org/10.1016/S1995-7645(11)60182-X)
55. Namazi R, Zabihollahi R, Behbahani M, Rezaei A. Inhibitory Activity of Avicennia marina, a Medicinal Plant in Persian Folk Medicine, against HIV and HSV. *Iran J Pharm Res.* 2013 Spring;12(2):435-43.
56. Thatoi, H., Samantaray, D., Das, SK., (2016). The genus Avicennia, a pioneer group of dominant mangrove plant species with potential medicinal values: a review, *Frontiers in Life Science*, DOI: 10.1080/21553769.2016.1235619
57. Alamgir M., Rob, DC Kundu, JHK, Joy, MM. 2007. Bioactivity of two medicinal plants Xylocarpus granatum Koen. (Meliaceae) and Sarcobolus globosus Wall. (Asclepiadaceae) of Sundarbans mangrove forest M, *Sarder Oriental Pharmacy and Experimental Medicine* 20077(4), 379-384 1
58. Shamsuzzaman M, Kalaiselvi K, Prabakaran M. Evaluation of Antioxidant and Anticorrosive Activities of Ceriops tagal Plant Extract. *Applied Sci.* 2021; 11(21):10150. <https://doi.org/10.3390/app112110150>
59. Mondal S, Ghosh D, Ramakrishna K. A Complete Profile on Blind-your-eye Mangrove Excoecaria Agallocha L. (Euphorbiaceae): Ethnobotany, Phytochemistry, and Pharmacological Aspects. *Pharmacogn Rev.* 2016.10(20):123-138, doi: 10.4103/0973-7847.194049.
60. Adhikari A, Ray M, Das AK, Sur TK. Antidiabetic and antioxidant activity of Rhizophora mucronata leaves (Indian Sundarbans mangrove): An in vitro and in vivo study. *Ayu.* 2016 Jan-Mar;37(1):76-81. doi: 10.4103/ayu.AYU_182_15.
61. Hossain SJ, Islam MR, Pervin T, Iftekharuzzaman M, Hamdi OAA, Mubassara S, Saifuzzaman M, Shilpi JA. Antibacterial, Anti-Diarrhoeal, Analgesic, Cytotoxic Activities, and GC-MS Profiling of Sonneratia apetala (Buch. -Ham.) Seed. *Prev Nutr Food Sci.* 2017 Sep;22(3):157-165. doi: 10.3746/pnf.2017.22.3.157.
62. Sur TK, Hazra A, Hazra AK, Bhattacharyya D. Antioxidant and hepatoprotective properties of Indian Sunderban mangrove Bruguiera gymnorrhiza L. leave. *J Basic Clin Pharm.* 2016 Jun;7(3):75-9. doi: 10.4103/0976-0105.183262
63. Gurudeeban S, Satyavani K, Ramanathan T, Balasubramanian T. Antidiabetic effect of a black mangrove species Aegicerascorniculatum in alloxan-induced diabetic rats. *J Adv Pharm Technol Res.* 2012 Jan;3(1):52-6. doi: 10.4103/2231-4040.93560
64. Nabeelah Bibi S, Fawzi MM, Gokhan Z, Rajesh J, Nadeem N, R.R. RK, R.D.D.G. A, Pandian SK. Ethnopharmacology, Phytochemistry, and Global Distribution of Mangroves—A Comprehensive Review. *Marine Drugs.* 2019; 17(4):231. <https://doi.org/10.3390/md17040231>
65. Dey, D, Quispe, C., Hossain,R., Jain, D., Khan, RA., Janmeda, P., Islam MT., et al., 2021. Ethnomedicinal Use, Phytochemistry, and Pharmacology of Xylocarpus
66. granatum J. Koenig", *Evidence-Based Complementary and Alternative Medicine*, 2021, 1- 16, <https://doi.org/10.1155/2021/8922196>

67. Mahmud I, Shahria N, Yeasmin S, Iqbal A, Mukul EH, Gain S, Shilpi JA, Islam MK. Ethnomedicinal, phytochemical and pharmacological profile of a mangrove plant *CeriopsDecandraGriffDin* Hou. *J Complement Integr Med.* 2018 Jun 22;16(1). doi: 10.1515/jcim-2017-0129.
68. Rahman MM, Al Noman MA, Khatun S, Alam R, Shetu MMH, Talukder EK, Imon RR, Biswas MY, Anis-UI-Haque KM, Uddin MJ, Akhter S. Evaluation of *Senna tora* (L.) Roxb. leaves as a source of bioactive molecules with antioxidant, anti-inflammatory and antibacterial potential. *Heliyon.* 2023 19;9(1): e12855. doi:10.1016/j.heliyon. 2023.e12855.
69. Abdul-Awal SM, Nazmir S, Nasrin S, Nurunnabi TR, Uddin SJ. Evaluation of the pharmacological activity of *Hibiscus tiliaceus*. *Springerplus.* 2016 Jul 29;5(1):1209. doi: 10.1186/s40064-016-2891-0.
70. Yusoff NA, Yam MF, Beh HK, Abdul Razak KN, Widyawati T, Mahmud R, Ahmad M, Asmawi MZ. Antidiabetic and antioxidant activities of *Nypafruticans* Wurm. vinegar sample from Malaysia. *Asian Pac J Trop Med.* 2015 Aug;8(8):595-605. doi: 10.1016/j.apjtm.2015.07.015.
71. Crowley, M.A., Cardille, J.A. Remote Sensing's Recent and Future Contributions to Landscape Ecology. *Curr Landscape Ecol Rep* 5, 45–57 (2020). [https://doi.org/ 10.1007/s40823-020-00054-9](https://doi.org/10.1007/s40823-020-00054-9)
72. Kumar AP., Gnanappazham, L., Selvam,V., Ramasubramanian R., Kar CS., (2015). Developing a spectral library of mangrove species of Indian east coast using field spectroscopy, *Geocarto Inter*, 30:5, 580-599. DOI: 10.1080/10106049.2014.985743
73. Mishra, S. P., Mishra, A., Kumar, C., Sahu, D. K., & Mishra, S. (2022). Distressed Lives and Livelihood in Biosphere Reserves during Anthropocene; Similipal Forest Blaze -2021. *Current Journal of Applied Science and Technology*, 41(25), 17-27. <https://doi.org/10.9734/cjast/2022/v41i2531772>
74. Mondal BK., 2015. Nature of Propensity of Indian Sundarbans. *International Journal of Applied Research and Studies (I JARS)*, 4(1)
75. India State of Forest Report (ISFR), 2011 to 2021, Forest Survey of India, Dehradun, <https://fsi.nic.in>
76. Mishra S. P., 2017, Stochastic Modelling of Flow and Sediment of the Rivers at Deltahead, East Coast of India, *American Journal of Operation Research, Scientific Research*, Vol. 7 (6), PP. 331-347, DOI: 10.4236/ajor.2017.76025
77. Hussain, N., Khanam, R., Khan E., Two and Half Century's Changes of World Largest Mangrove Forest: A Geo-informatics Based Study on Sundarbans Mangrove Forest, Bangladesh, India. *Modern Envir. Sci. and Eng.* 2017, 3(6), 419-423DOI: 10.15341/mese (2333-2581)/06.03.2017/007
78. Mishra S. P., 2017, Stochastic Modelling of Flow and Sediment of the Rivers at Delta head, East Coast of India, *American Journal of Operation Research, Scientific Research*, 7 (6), 331-347, DOI: 10.4236/ajor.2017.76025
79. Mahamud U, Takewaka S. Shoreline Change around a River Delta on the Cox's Bazar Coast of Bangladesh. *Journal of Marine Science and Eng.* 2018; 6(3):80. <https://doi.org/10.3390/jmse6030080>