

Original Research Article

Effect of Different Pulsing Solutions on Postharvest Life of Tuberose (*Polianthes tuberosa* L.) cv. Prajwal

Abstract

Tuberose flower (*Polianthes tuberosa* L.) is a perennial and bulbous flowering plant, related to the Amaryllidaceae family and now it is placed in family Agavaceae. The investigation entitled "Effect of different pulsing solutions on postharvest life of tuberose (*Polianthes tuberosa* L.) cv. Prajwal" was carried out at the experimental farm and laboratory of Department of Floriculture and Landscape Architecture, College of Horticulture, SVPUAT, Meerut during the year of 2021& 2022. Experiment was laid out in Completely Randomized Design with 10 treatments and 3 replications. It was observed that the overall best treatment for different parameters for improving the vase life of tuberose were recorded under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) followed by the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) and T₈ (Sucrose 15% + 8-HQC 200 ppm) resulted in most of the postharvest parameters such as amount of pulsing solution consumed (13.16, 15.82 ml), days taken to opening of floret (2.53, 2.54 days), floret diameter (3.38, 3.40 cm), vase life (10.78, 10.80 days), weight change/loss in cut spike (13.29 %, 14.52 %), fungal infection, fragrance score (2.95, 2.96), freshness index (4.97, 4.98) and moisture content (89.89, 85.60, 83.21%).

Keywords: Tuberose, Pulsing, Sucrose, 8-HQC, Pulsing solution, distilled water and Vase life.

1. Introduction

Tuberose flower (*Polianthes tuberosa* L.) is most important bulbous flower plant related to a member of the Amaryllidaceae family and now it is placed in the family Agavaceae. It originated from Mexico and was grown before its conquest in 1522. It is

Comment [AR1]: The title does not need to be rewritten, just write the purpose of the study

Comment [AR2]: Delete, just 5 words

Comment [AR3]: Use the most recent literature for at least the last 10 years

one of the most popular and fragrant cut flowers grown in India as well as worldwide (Singh and Shankar 2011). For single-stemmed tuberose, the basic chromosome number is $2n=60$, whereas the chromosome number of double-flowered species have ranged from $2n=50$ to $2n=54$, 60 and 120 (Lin and Shen 2004), (Karihaloo, 2019).

Single-flowered cultivars that have one row of corolla sections are mostly used to extract essential oils and loose flowers etc., while double-type of cultivars of tuberose that have more than two rows of corolla sections are used for cut flowers and displays of garden. The 'single type' of tuberose are more fragrant than the 'double type' and concentration ranges from 0.08-0.14 per cent (Singh and Uma 1995).

Sucrose plays a most important role in the quality of cut flowers because it acts as a source of food or respiratory substrate and delays the protein degradation and improves the balance of water in cut flowers (Asrar, 2012). The addition of sucrose to the solution increased the mechanical rigidity of the stem inducing thickening of the cell wall and lignifications of vascular tissues (Steinitz, 1982). The addition of sucrose replaces the depletion of carbohydrates from cut stems and maintains the respiratory pool thereby prolonging vase life (Marousky, 1971). 8-hydroxyquinoline citrate (8-HQC) plays an important role in decreasing the physiological blockage in the sterile tissue (Rogers, 1973) and (Elhindi, 2012). It acts as an antimicrobial agent for its activity thus it helps in maintaining the flower head diameter of flowers (Nowak and Rudnicki 1979). 8-hydroxyquinoline citrate (8-HQC) has been better known to possess property like antimicrobial that inhibits bacterial and physiological vascular blockage and enhances uptake of water in cut flowers (Larsen and Cromarty 1967).

Pulsing is a process in which freshly harvested flowers are kept in a solution for few seconds to hours to extend their vase life (Sao and Verma 2020). It has been reported that pulsing treatments prevent infections of vascular and ethylene production inhibition and results in a prolonged period of storage and flowers with high quality and increased vase life (Vidhya and Bhattacharjee 2002). The main challenge of tuberose is that being a cut flower, it is a short-lived and is prone to rapid deterioration. The other important challenges and symptoms that shorten the vase life are abscission of buds, lack of bud opening and leaf yellowing due to the sensitivity of cut tuberose as referred

by Reid and Jiang (2012). In India, pulsing helps enhance the vase life of cut Tuberose, there is a wide scope for experimentation, standardization and improvement of different techniques in tuberose. Therefore the present study was carried out with the objective to evaluate the effect of different Pulsing solutions on the Postharvest Life of Tuberose (*Polianthes tuberosa* L.) cv. Prajwal.

2. Materials and Methods

The present investigation was carried out in the Laboratory of the Department of Floriculture and Landscape Architecture, College of Horticulture, SVPUAT, Meerut during the year of 2021&2022. The experimental location, Meerut comes under the semi-arid region and Agro-climatic plain zone of Uttar Pradesh state and lies at North West Plain Zone, India, 28.99° N Latitude and 77.7° E Longitude with an altitude of 220 m above the mean sea level. The general climate in Meerut region of Western Uttar Pradesh is dry sub-humid type with annual rainfall varying from 900-1000 mm approximately every year.

Comment [AR4]: Pay attention to correct writing

2.1 Plant material

Tuberose (*Polianthes tuberosa* L.) cultivar 'Prajwal' uniform, healthy and disease free bulbs were obtained from the previous year tuberose crop grown in the field of Horticulture Research Centre (HRC), Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut. Planting was done in beds at a spacing of 30x30 cm. The cut spikes of uniform size were harvested in the morning hours (7-8 am) with the help of sharp secateurs.

2.2 Treatments and observations

The experiments were repeated twice for confirmation of the results, with ambient temperature of 15-18°C, 60±5 Relative humidity and average radiation around 5000 Lux for a period of 8±2 h/day. After harvesting, cut spikes were taken to the laboratory for conducting the experiment. When the first 1-2 florets start opening, these spikes were immediately placed in a bucket containing distilled water. After transporting to our laboratory, a slanting cut was given at the base of each cut spikes in such a

manner that the final length of cut spike remained 50-55 cm. Before placed cut spikes in different pulsing solutions, firstly their fresh weights were recorded. Thereafter, these cut spikes were placed in 500 ml of glass bottles containing 250 ml of pulsing solution at different levels for 12 hours at room temperature. The neck of each bottle was covered with the help of aluminum foil to check evaporation of the solution or distilled water. The experiment were subjected to 10 treatments of pulsing and holding solutions with 3 replications, arranged in a completely randomized design.

Treatment details of holding solutions used in the experiment consists of : T₀: Control (Distilled water), T₁: Sucrose 5% + 8-HQC (100 ppm), T₂: Sucrose 5% + 8-HQC (200 ppm), T₃: Sucrose 5% + 8-HQC (300 ppm), T₄: Sucrose 10% + 8-HQC (100 ppm), T₅: Sucrose 10% + 8-HQC (200 ppm), T₆: Sucrose 10% + 8-HQC (300 ppm), T₇: Sucrose 15% + 8-HQC (100 ppm), T₈: Sucrose 15% + 8-HQC (200 ppm), T₉: Sucrose 15% + 8-HQC (300 ppm). The observations were recorded on different post-harvest parameters such as, amount of pulsing solution consumed, days taken to opening of floret, floret diameter, vase life, weight change/loss in cut spike, fungal infection, fragrance score, freshness index and moisture content.

2.3 Statistical analysis

The recorded data were statistically analysis (ANOVA) using the software Graph pad prism, USA. Source of variation were pulsing agents viz., Sucrose and 8-HQC. Mean data were also compared by using Duncan's new multiple range test whereas the effect were significant at 5% level of significance.

3. Results and discussion

3.1 Amount of pulsing solution consumed: (ml)

It is evident from Table 1 that the maximum pulsing solution consumed (13.16 ml) during the first season 2021-22 were recorded under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) at par (12.90 ml and 12.89 ml, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₆ (Sucrose 10% + 8-HQC 300 ppm) while, minimum pulsing solution consumed (5.11 ml) was recorded under T₀ (Control). During

Comment [AR5]: Delete

the second season of 2022-23, maximum pulsing solution consumed (15.82 ml) was recorded under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) followed by (14.75 ml and 13.25 ml, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₆ (Sucrose 10% + 8-HQC 300 ppm) While, minimum pulsing solution consumed (6.24 ml) was recorded under T₀ (Control). In the present experiment, pulsing solution consumed was maximum under these treatments, it might be due to the fact that 8-HQC is known to enhance uptake of solution (Larsen and Cromarty, 1967) in cut flower. These treatments improved longevity of flower by decreasing microbial growth in bundles of vascular and increased tendency of cut flowers to absorb more solution. Similar findings were reported by (Marousky, 1969 in roses, Nowak and Rudnicki, 1990 in cut flowers, Murry, 2014 in tuberose and Thakur *et al.*, 2019 in iris).

Comment [AR6]: Use the latest literature

Comment [AR7]: Use the latest literature

Table 1: Effect of different pulsing solutions on pulsing solution consumed (ml) by *Polianthes tuberosa* L. cv. Prajwal

Treatment	Pulsing Solution	Pulsing Solution Consumed (ml)	
		2021-22	2022-23
T ₀	Control (Distilled Water)	5.11	6.24
T ₁	Sucrose 5% + 8-HQC (100ppm)	8.28	10.32
T ₂	Sucrose 5% + 8-HQC (200ppm)	9.31	10.78
T ₃	Sucrose 5% + 8-HQC (300ppm)	8.92	9.77
T ₄	Sucrose 10% + 8-HQC (100ppm)	10.45	11.56
T ₅	Sucrose 10% + 8-HQC (200ppm)	10.14	11.40
T ₆	Sucrose 10% + 8-HQC (300ppm)	12.89	13.25
T ₇	Sucrose 15% + 8-HQC (100ppm)	11.51	12.42
T ₈	Sucrose 15% + 8-HQC (200ppm)	12.90	14.75
T ₉	Sucrose 15% + 8-HQC (300ppm)	13.16	15.82
	SEm ±	0.34	0.39
	C.D.(p=0.05)	1.03	1.19

3.2 Days taken to opening of floret: (days)

Comment [AR8]: Delete

It is clear from Table 2 that solution of pulsing improved days taken to opening of floret of cut spikes of tuberose. The minimum days taken to opening of floret (2.53 days) during the first season 2021-22 were recorded under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) at par (2.65 days and 2.74 days, respectively) under the treatment

T₉(Sucrose 15% + 8-HQC 300 ppm) and T₆ (Sucrose 10% + 8-HQC 300 ppm) while, maximum days taken to opening of floret (4.13 days) was recorded under T₀ (control). During the second season of 2022-23, minimum days taken to opening of floret (2.54 days) were recorded under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) at par (2.66 days and 2.75 days, respectively) under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) and T₆ (Sucrose 10% + 8-HQC 300 ppm) while, maximum days taken to opening of floret (4.16 days) was recorded under T₀ (control). In the present experiment, minimum days taken to floret opening under these treatments, it might be due to opening of floret is dependent on levels of carbohydrate in the petal (Doorn *et al.*, 1991 in carnation) and growth of petal is associated with opening of flower bud which results from expansion of cell and this requires the influx of water and carbohydrates into cell of petal (Evan and Reid 1988 in bedding plants). The solutions of sugars and biocide are effective for bud opening of cut flowers like carnation, chrysanthemum and gladiolus (Bhattacharjee, 1999 in cut flowers). The absence of any pulsing solution (control) did not induced earliness for floret opening as nutrients supply was inhibited. Similar findings were reported by (Sigma *et al.*, 2018 in tuberose, Thakur *et al.*, 2019 in iris and Baidya and Chakrabarty 2020 in tuberose).

Comment [AR9]: Use the latest literature

Comment [AR10]: Use the latest literature

Table 2: Effect of different pulsing solutions on days taken to opening of floret (days) in *Polianthes tuberosa* L. cv. Prajwal

Treatment	Pulsing Solution	Days taken to opening of floret (days)	
		2021-22	2022-23
T ₀	Control (Distilled Water)	4.13	4.16
T ₁	Sucrose 5% + 8-HQC (100ppm)	3.36	3.38
T ₂	Sucrose 5% + 8-HQC (200ppm)	3.18	3.21
T ₃	Sucrose 5% + 8-HQC (300ppm)	3.56	3.57
T ₄	Sucrose 10% + 8-HQC (100ppm)	2.96	2.98
T ₅	Sucrose 10% + 8-HQC (200ppm)	3.10	3.12
T ₆	Sucrose 10% + 8-HQC (300ppm)	2.74	2.75
T ₇	Sucrose 15% + 8-HQC (100ppm)	2.87	2.89
T ₈	Sucrose 15% + 8-HQC (200ppm)	2.53	2.54
T ₉	Sucrose 15% + 8-HQC (300ppm)	2.65	2.66
	SEm ±	0.10	0.11
	C.D.(p=0.05)	0.32	0.33

3.3 Floret diameter: (cm)

Data presented in Table 3 revealed that solution of pulsing improved diameter of floret. The maximum diameter of floret (3.38 cm) during the first season 2021-22 were recorded under the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) at par (3.25 cm and 3.15 cm, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₉ (Sucrose 15% + 8-HQC 300 ppm) While, minimum diameter of floret (2.35 cm) was recorded under T₀ (Control). During the second season of 2022-23, maximum diameter of floret (3.40 cm) was recorded under the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) at par (3.27 cm and 3.18 cm, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₉ (Sucrose 15% + 8-HQC 300 ppm) While, minimum diameter of floret (2.37 cm) was recorded under T₀ (Control). In the present experiment, diameter of floret was maximum under these treatments; it might be due to blockage avoidance of xylem tissues by 8-HQC and sucrose utilization as source of energy by the cut flowers. The carbohydrates accumulation and more water uptake had direct effect on increase in the volume of cell. The cell of petal of untreated flowers with poor water uptake might have been flaccid and deplasmolysed, leading to reduced cell size and that of petals. Our findings were corroborated with (Kumar and Gupta 2014 in *Gladiolus*, Jain *et al.*, 2014 in *Chrysanthemum*, Jadeja *et al.*, 2017 in *Tuberose*, Thakur *et al.*, 2019 in *Iris* and Happy *et al.*, 2021 in *Tuberose*).

Table 3: Effect of different pulsing solutions on floret diameter (cm) of *Polianthes tuberosa* L. cv. Prajwal

Treatment	Pulsing Solution	Floret diameter (cm)	
		2021-22	2022-23
T ₀	Control (Distilled Water)	2.35	2.37
T ₁	Sucrose 5% + 8-HQC (100ppm)	2.60	2.63
T ₂	Sucrose 5% + 8-HQC (200ppm)	2.80	2.82
T ₃	Sucrose 5% + 8-HQC (300ppm)	2.71	2.74
T ₄	Sucrose 10% + 8-HQC (100ppm)	2.99	3.01
T ₅	Sucrose 10% + 8-HQC (200ppm)	2.88	2.89
T ₆	Sucrose 10% + 8-HQC (300ppm)	3.38	3.40
T ₇	Sucrose 15% + 8-HQC (100ppm)	3.07	3.08

T ₈	Sucrose 15% + 8-HQC (200ppm)	3.25	3.27
T ₉	Sucrose 15% + 8-HQC (300ppm)	3.15	3.18
	SEm ±	0.10	0.10
	C.D.(p=0.05)	0.31	0.31

3.4 Vase life: (days)

Comment [AR12]: Delete

The data as per Table 4 reveals that there is significant effect of different pulsing solution on vase life of cut flower of tuberose. The maximum vase life (10.78 days) during the first season 2021-22 were observed under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) which was *at par* (10.12 days and 9.48 days, respectively) under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) and T₆ (Sucrose 10% + 8-HQC 300 ppm) while, minimum vase life (5.06 days) was recorded under T₀ (Control). During the second season of 2022-23, maximum vase life (10.80 days) were observed under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) *at par* (10.14 days and 9.50 days, respectively) under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) and T₆ (Sucrose 10% + 8-HQC 300 ppm) while, minimum vase life (5.07 days) was recorded under T₀ (Control). In the present experiment, vase life was observed maximum under these treatments, it might be due to the fact that once blockage of vascular is avoided by 8-HQC with sucrose that facilitated the higher water intake and accumulation of total soluble sugars in the cells of petal probably by enhancing the osmotic driving force for the uptake of solution, resulting in longer vase life (Nagarjuna *et al.*, 2002) in tuberose. The present study is in line with the previous findings made by (Jain *et al.*, 2014 in chrysanthemum, Naznin *et al.*, 2015 in tuberose, Sigma *et al.*, 2018 in tuberose, Thakur *et al.*, 2019 in iris and Baidya and Chakrabarty 2020 in tuberose).

Table 4: Effect of different pulsing solutions on vase life (days) of *Polianthes tuberosa* L. cv. Prajwal

Treatment	Pulsing Solution	Vase life (days)	
		2021-22	2022-23
T ₀	Control (Distilled Water)	5.06	5.077
T ₁	Sucrose 5% + 8-HQC (100ppm)	6.68	6.70
T ₂	Sucrose 5% + 8-HQC (200ppm)	7.76	7.77
T ₃	Sucrose 5% + 8-HQC (300ppm)	7.05	7.08
T ₄	Sucrose 10% + 8-HQC (100ppm)	8.71	8.73

T ₅	Sucrose 10% + 8-HQC (200ppm)	8.25	8.32
T ₆	Sucrose 10% + 8-HQC (300ppm)	9.48	9.50
T ₇	Sucrose 15% + 8-HQC (100ppm)	9.04	9.07
T ₈	Sucrose 15% + 8-HQC (200ppm)	10.78	10.80
T ₉	Sucrose 15% + 8-HQC (300ppm)	10.12	10.14
	SEm ±	0.30	0.29
	C.D.(p=0.05)	0.92	0.89

3.5 Weight change/loss in cut spike: (%)

Comment [AR13]: Delete

A perusal of data presented in Table 5 revealed that the significant effect of pulsing solution on weight change. The minimum weight change/loss in cut spike (13.29 %) during the first season 2021-22 were recorded under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) followed by (15.45 % and 17.28 %, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₇ (Sucrose 15% + 8-HQC 100 ppm) while, Maximum weight change/loss in cut spike (45.12 %) was recorded under T₀ (control). During the second season of 2022-23, minimum weight change/loss in cut spike (14.52 %) during the first season 2021-22 were recorded under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) followed by (16.79 % and 18.56 %, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₇ (Sucrose 15% + 8-HQC 100 ppm) while, Maximum weight change/loss in cut spike (47.15 %) was recorded under T₀ (control). In the present experiment, weight change/loss in cut spike was minimum under these treatments, it might be due to sucrose is used as nutrition (Marousky, 1972) in cut flowers, while 8-HQC prevents physiological and microbial vascular blockage as well as stimulate closure of stomata (Larsen and Frolich 1969) in carnation that ultimately leads to minimum weight change/loss in cut spike in tuberose. Similar findings were reported by Jain *et al.*, 2014 in chrysanthemum, Murry, 2014 in tuberose, Kumar and Gupta 2014 in Gladiolus, and Thakur *et al.*, 2019 in iris).

Table 5: Effect of different pulsing solutions on weight change/loss after ending of vase life in cut spike (%) of *Polianthes tuberosa* L. cv. Prajwal

Treatment	Pulsing Solution	Weight change in cut spike (%)	
		2021-22	2022-23
T ₀	Control (Distilled Water)	45.12	47.15
T ₁	Sucrose 5% + 8-HQC (100ppm)	36.53	37.32

T ₂	Sucrose 5% + 8-HQC (200ppm)	28.09	30.25
T ₃	Sucrose 5% + 8-HQC (300ppm)	32.78	33.68
T ₄	Sucrose 10% + 8-HQC (100ppm)	25.63	26.47
T ₅	Sucrose 10% + 8-HQC (200ppm)	23.87	25.13
T ₆	Sucrose 10% + 8-HQC (300ppm)	20.59	22.40
T ₇	Sucrose 15% + 8-HQC (100ppm)	17.28	18.56
T ₈	Sucrose 15% + 8-HQC (200ppm)	15.45	16.79
T ₉	Sucrose 15% + 8-HQC (300ppm)	13.29	14.52
	SEm ±	0.91	0.96
	C.D.(p=0.05)	2.74	2.87

3.6 Fungal infection

A perusal of symbols on fungal infection are presented in Table 6 revealed that during both season (2021-22 and 2022-23), showed that T₀, T₁, T₃ and T₇ treatment solutions were infected by fungus and rest of the solutions were significantly not infected by fungal infection. Adding a suitable germicide in vase solution can prevent the growth of microbes and increase the tendency for uptake of water (Anjum *et al.*, 2001) in tuberose.

Table 6:Effect of different pulsing solutions on fungal infection in *Polianthes tuberosa* L. cv. Prajwal

Treatment	Pulsing Solution	Fungal infection	
		2021-22	2022-23
T ₀	Control (Distilled Water)	+	+
T ₁	Sucrose 5% + 8-HQC (100ppm)	+	+
T ₂	Sucrose 5% + 8-HQC (200ppm)	-	-
T ₃	Sucrose 5% + 8-HQC (300ppm)	+	+
T ₄	Sucrose 10% + 8-HQC (100ppm)	-	-
T ₅	Sucrose 10% + 8-HQC (200ppm)	-	-
T ₆	Sucrose 10% + 8-HQC (300ppm)	-	-
T ₇	Sucrose 15% + 8-HQC (100ppm)	+	+
T ₈	Sucrose 15% + 8-HQC (200ppm)	-	-
T ₉	Sucrose 15% + 8-HQC (300ppm)	-	-
	SEm ±		
	C.D.(p=0.05)		

'+' represent infected by fungus

'-' represent not infected by fungus

3.7 Fragrance score

The data as presented in Table 7 for Fragrance score revealed that the strong fragrance score (2.95) during the first season 2021-22 were noticed under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) at par (2.80 and 2.63, respectively) under the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) and T₈ (Sucrose 15% + 8-HQC 200 ppm) while, undesirable and least fragrance score (1.13) was noticed under T₀ (Control). During the second season of 2022-23, strong fragrance score (2.96) were noticed under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) at par (2.81 and 2.65, respectively) under the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) and T₈ (Sucrose 15% + 8-HQC 200 ppm) while, undesirable and least fragrance score (1.14) was noticed under T₀ (Control). In the present experiment Tuberose flower pre cooled at room temperature which retained better freshness. The endogenous and exogenous ethylene leads to loss of fragrance (Sexton *et al.*, 2005) in sweet pea. This ethylene action could be delayed by effect of sucrose (Pun *et al.*, 2016). Production of many flowers, fragrances displays a circadian cycle that is they are stronger at certain times during the day or night where as tuberose are more fragrant during evening as they are blooming in night in nature. In this context the present findings were in line with the findings made by (Reethu *et al.*, 2022 in tuberose).

Table 7: Effect of different pulsing solutions on fragrance score in *Polianthes tuberosa* L. cv. Prajwal

Treatment	Pulsing Solution	Fragrance score	
		2021-22	2022-23
T ₀	Control (Distilled Water)	1.13	1.14
T ₁	Sucrose 5% + 8-HQC (100ppm)	1.27	1.29
T ₂	Sucrose 5% + 8-HQC (200ppm)	2.13	2.15
T ₃	Sucrose 5% + 8-HQC (300ppm)	1.55	1.56
T ₄	Sucrose 10% + 8-HQC (100ppm)	2.13	2.15
T ₅	Sucrose 10% + 8-HQC (200ppm)	2.30	2.32
T ₆	Sucrose 10% + 8-HQC (300ppm)	2.80	2.81
T ₇	Sucrose 15% + 8-HQC (100ppm)	2.30	2.32
T ₈	Sucrose 15% + 8-HQC (200ppm)	2.63	2.65
T ₉	Sucrose 15% + 8-HQC (300ppm)	2.95	2.96
	SEm ±	0.08	0.08

	C.D.($p=0.05$)	0.24	0.24
--	----------------------------------	------	------

3.8 Freshness index

A perusal of data on freshness index are presented in Table 8 revealed that the original brighter colour of flowers have freshness index (4.97) during the first season 2021-22 were noticed under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm), flowers with faded from original colour at par (4.63 and 4.30, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₆ (Sucrose 10% + 8-HQC 300 ppm) while, flowers severely faded from original colour have freshness index (0.98) was noticed under T₀ (Control). During the second season of 2022-23, original brighter colour of flowers have freshness index (4.98) were noticed under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm), flowers with faded from original colour at par (4.65 and 4.31, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₆ (Sucrose 10% + 8-HQC 300 ppm) while, flowers severely faded from original colour have freshness index (0.92) was noticed under T₀ (Control). Tuberose flower pre cooled at room temperature retained better freshness. it might be due to pre-cooling is a technique used for the removal of heat of field which leads to reduction of both the rate of metabolism and prevention of loss of water which delays freshness is reported by. Similar findings were reported by (Reethu *et al.*, 2022 in tuberose).

Table 8: Effect of different pulsing solutions on freshness index in *Polianthes tuberosa* L. cv. Prajwal

Treatment	Pulsing Solution	Freshness index	
		2021-22	2022-23
T ₀	Control (Distilled Water)	0.98	0.92
T ₁	Sucrose 5% + 8-HQC (100ppm)	1.30	1.32
T ₂	Sucrose 5% + 8-HQC (200ppm)	2.63	2.64
T ₃	Sucrose 5% + 8-HQC (300ppm)	1.30	1.32
T ₄	Sucrose 10% + 8-HQC (100ppm)	2.63	2.64
T ₅	Sucrose 10% + 8-HQC (200ppm)	3.30	3.32
T ₆	Sucrose 10% + 8-HQC (300ppm)	4.30	4.31
T ₇	Sucrose 15% + 8-HQC (100ppm)	3.30	3.32
T ₈	Sucrose 15% + 8-HQC (200ppm)	4.63	4.65
T ₉	Sucrose 15% + 8-HQC (300ppm)	4.97	4.98

	SEm ±	0.12	0.12
	C.D.(p=0.05)	0.36	0.37

3.9 Moisture content (%)

Comment [AR14]: Delete

In the present investigation the moisture content decreased from 1st day to 5th day of storage. The data on moisture content for both the years (2021-22 and 2022-23) are presented in Table 9 revealed that at 1st day the maximum moisture content (89.89 %) were recorded under the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) followed by (88.60 % and 85.85 %, respectively) under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) and T₈ (Sucrose 15% + 8-HQC 200 ppm) while, minimum moisture content (67.91 %) was recorded under T₀ (Control). Similarly, at 3rd day maximum moisture content (85.60 %) were recorded under the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) followed by (83.69 % and 81.46 %, respectively) under the treatment T₈ (Sucrose 15% + 8-HQC 200 ppm) and T₉ (Sucrose 15% + 8-HQC 300 ppm) while, minimum moisture content (59.68 %) was recorded under T₀ (Control). Moreover, at 5th day maximum moisture content (83.21 %) were recorded under the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) followed by (79.03 % and 77.94 %, respectively) under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) and T₈ (Sucrose 15% + 8-HQC 200 ppm) while, minimum moisture content (54.95 %) was recorded under T₀ (Control). The present results reported that that the slower moisture loss from pre-cooled commodities during storage, it might be due to the fact that such conditions can easily be achieved by lowering the temperature of storage, as the environment tends to be more saturated simply by reduction in temperature which brings down the heat load thus loss of moisture is reduced (Lurie and Ben 1990). In addition to this, the sucrose maintained balance of water. The present results well corroborated the findings obtained in tuberose (Sharma *et al.*, 2008 in asiatic hybrid lily and Reethu *et al.*, 2022 in tuberose).

Table 9: Effect of different pulsing solutions on moisture content (%) in *Polianthes tuberosa* L. cv. Prajwal

Treat.	Pulsing Solution	Moisture (%) at 1 st day	Moisture (%) at 3 rd day	Moisture (%) at 5 th day
--------	------------------	-------------------------------------	-------------------------------------	-------------------------------------

		2021-22	2022-23	Pooled Mean	2021-22	2022-23	Pooled Mean	2021-22	2022-23	Pooled Mean
T ₀	Control (Distilled Water)	67.31	68.51	67.91	60.65	58.72	59.68	55.11	54.79	54.95
T ₁	Sucrose 5% + 8-HQC (100ppm)	78.56	78.37	78.46	73.51	73.87	73.69	65.60	66.84	66.22
T ₂	Sucrose 5% + 8-HQC (200ppm)	75.92	73.85	74.89	69.00	72.45	70.72	62.86	64.98	63.92
T ₃	Sucrose 5% + 8-HQC (300ppm)	78.32	76.94	77.63	72.83	79.81	76.32	66.54	67.93	67.23
T ₄	Sucrose 10% + 8-HQC (100ppm)	82.72	82.73	82.73	77.30	76.52	76.91	71.99	72.56	72.27
T ₅	Sucrose 10% + 8-HQC (200ppm)	83.35	81.31	82.33	80.45	80.44	80.45	73.29	74.41	73.85
T ₆	Sucrose 10% + 8-HQC (300ppm)	89.59	90.19	89.89	84.85	86.35	85.60	81.88	84.54	83.21
T ₇	Sucrose 15% + 8-HQC (100ppm)	80.30	83.90	82.10	80.64	81.05	80.84	73.88	75.14	74.51
T ₈	Sucrose 15% + 8-HQC (200ppm)	83.66	88.04	85.85	83.36	84.02	83.69	75.89	79.99	77.94
T ₉	Sucrose 15% + 8-HQC (300ppm)	87.62	89.58	88.60	81.27	81.65	81.46	76.49	81.58	79.03
	SEm ±	0.50	0.44		0.45	0.76		0.72	0.99	
	C.D.(p=0.05)	1.52	1.32		1.36	2.29		2.16	2.99	

Conclusion

From the above results, it can be concluded that the overall best treatment for different parameters for improving the vase life of tuberose during the season 2021&2022 were recorded under the treatment T₉ (Sucrose 15% + 8-HQC 300 ppm) followed by the treatment T₆ (Sucrose 10% + 8-HQC 300 ppm) and T₈ (Sucrose 15% + 8-HQC 200 ppm) which significantly improved the postharvest life of tuberose spikes cv 'Prajwal'. The various holding solution significantly improved the different postharvest parameters in tuberose viz., amount of pulsing solution consumed, days taken to opening of floret, floret diameter, vase life, weight change/loss in cut spike, fungal infection, fragrance score, freshness index and moisture content significantly influenced and performed better under these treatments.

Future prospects

There is a great scope of Pulsing as it improves and enhances the quality of cut flowers, by reviewing the different factors and approaches associated with the use of the solution of floral preservative under this experiment for the development of proper handling in the industry of cut flowers.

Comment [AR15]: Delete

References

- Anjum, M. A., Naveed, F., Shakeel, F. and Amin, S. (2001).** Effect of some chemicals on keeping quality and vase-life of tuberose (*Polianthes tuberosa* L.) cut flowers. *Life*, 12(1), 23-65.
- Baidya, B. K. and Chakrabarty, S. (2020).** Increasing vase life of tinted spikes of (*Polianthes tuberosa* L.) cv. 'Prajwal' by adding floral preservatives. *Int. J. Curr. Microbiology App. Sci.*, 9(8): 2736-2758.
- Bhattacharjee, S.K. (1999).** Postharvest management of cut flowers, cut foliages and post production management of potted plants. *Journal of Ornamental Horticulture*, (2):32-39.
- Datta, S.K. (2017).** Breeding of ornamentals: Tuberose (*Polianthes tuberosa* L.). *Curr. Sci.*, (113): 1255–1263. [CrossRef]
- Doorn, W.G.V., Zagory, D., Witle, Y. de. and Harkema, H. (1991).** Effects of vase water bacteria on the senescence of cut carnation flowers. *Postharvest Biology and Technology* (1):161-168.
- Evan, M.R. and Reid, S.H. (1988).** Growth of bedding plants in sphagnum peat and coir dust based substrates. *Journal of Environmental Horticulture* (14):187- 190.
- Happy, D. C., Devi, S., Malik, A. and Kumar, S. (2021).** Effect of Wrapping Materials and Storage Time on Shelf Life of Tuberose (*Polianthes tuberosa* L.) *Agricultural Science Digest*, (1-8).
- Jadeja, R., Singh, A., Ahlawat, T. R., Baghale, R. D. and Chawla, S. L. (2017).** Effect of various chemicals as vase solutions on post harvest flower quality parameters in tuberose (*Polianthes tuberosa*) cut spikes. *Progressive Horticulture*, 49(1): 79-85.
- Jain, R., Janakiram, T., Singh, K. P. and Kumawat, G. L. (2014).** Effect of different floral preservatives on reducing foliage discoloration and increasing vase life of chrysanthemum (*Dendranthemax grandiflora*) cv. 'White Reagan'. *Indian Journal of Agricultural Sciences*, 84(11), 1386-88.

Karihaloo, J.L. (2019) Cytological and hybridization studies in three cultivars of tuberose (*Polianthes tuberosa* L.). *Cytologia*, (84): 47–52. [CrossRef]

Kumar, S. and Gupta, A. K. (2014). Postharvest life of *Gladiolus grandiflorus* L. cv. 'Jessica' as influenced by pre-harvest application of gibberellic acid and kinetin. *Journal of Post-harvest Technology*, 2(3), 169-176.

Larsen, F.E. and Cromarty, R.W. (1967). Microorganism inhibition by 8-hydroxyquinoline citrate as related to cut flower senescence. *Journal of the American Society for Horticultural Science* (90):546-549.

Larsen, F.E. and Frolich, M. (1969). The influence of 8-HQC, N-dimethylamine succinamic and sucrose on respiration and water flow in Red Sim carnation in relation of flower senescence. *Journal of American Society of Horticultural Sciences* (94):289-291.

Lin, T.S. and Shen, T.M. (2004) Study on karyotype of *Polianthes tuberosa* L. *J. Agric. For. NCYU*, (1):1–12.

Lurie, S.K. and Ben, A.R. (1990). Physiological changes in diphenyl treated 'Granny Smith' apples. *Israel J Bot.* (38):199- 207.

Marousky, F.G. (1969). Vascular blockage, water absorption, stomatal opening and respiration of cut „Better Times“ roses treated with 8-hydroxyquinoline citrate and sucrose. *Journal of the American Society for Horticultural Science*, (94):223-226.

Marousky, F.G. (1972). Water relations, effect of floral preservatives on bud opening and keeping quality of cut flowers. *HortScience* (7):114-117.

Marousky, F.J. (1971). Inhibition of vascular blockage and increased moisture in cut rose induced by pH, 8- hydroxyquinoline citrate and sucrose. *Journal of American Society for Horticulture and Science.*, 96(1):38-41.

Murry, N. A. (2014). Postharvest Handling of Tuberose (*Polianthes tuberosa* Linn.) cv. 'Double'. MSc Thesis. Department of Floriculture and Landscape Architecture, Dr YS Parmar University of Horticulture and Forestry, Solan. (64p).

Nagarjuna, H.T., Narayanagowda, J.V. and Nagaraja, G.S. (2002). Effect of pulsing with sucrose on vase life of tuberose cv. 'Double'. *Crop Research Hisar*, (23):349- 353.

Naznin, A., Hossain, M. d., Kabita, A. M., Azizul, H. M., Islam, M. and Hasan, T. (2015). Effect of Different Preservatives on Vase Life of Tuberose, *5(2)*:105-113.

Nowak, J. and Rudnicki, I.R.M. (1990). Postharvest handling and storage of cut flowers, florist green and potted plants. *Timber Press, INC. Portland, Oregon*, (210p).

Nowak, J. and Rudnicki, R.M. (1979). Long term storage of cut flowers. *Acta. Hort.*, (91):123-134.

Reethu, G. R., Jhalegar, M. J., Jagadeesh, S. L., Bhuvaneshwari, G., Babu, A. G. and Pavankumar, P. (2022). Effect of pre-cooling methods on shelf life of cold stored tuberose florets. *Journal of Pharmacognosy and Phytochemistry*, 11(5): 181-185.

Sao, B. and Verma, L. S. (2020). Review on impact of different preservative solutions on vase-life of tuberose (*Polianthes tuberosa* L.) cut flowers. *Journal of Pharmacognosy and Phytochemistry*, 9(1): 1185-1188.

Sexton, R., Stopford, A.P., Moodie, W.T. and Porter, A.E.A. (2005). Aroma production from cut sweet pea flowers (*Lathyrus odoratus*): the role of ethylene. *Physiologia Plantarum*, (124):381-389.

Sharma, B.P., Beshir, M.M., Dilita, B.S. and Chaudhry, S.V.S. (2008). Effect of various wrapping materials and storage durations on post harvest life of Asiatic hybrid lily cv. 'Apeldoorn'. *Proc. Nwation. Symp, Scenario of Agriculture in changing climatic conditions*, (69-74).

Sigma, T. N., Hoque, M. A. and Hossain, M. M. (2018). Effect of Different Preservatives on Vase Life of Tuberose. *Journal of Crop Science and Technology*, 7(3): 10-15.

Singh, A. K. and Shankar, K. (2011). Effect of Plant Growth Regulators on Vegetative Growth and Flowering Behaviour of Tuberose (*Polianthes tuberosa* L.) cv. 'Double'. *Plant Arch.* (11): 123-125.

Singh, K.P. and Uma, S. (1995). Studies on ratoon crop in tuberose cv. Single and Double. *Indian Perfum.*, (39): 158–160.

Steinitz, B. (1982). Role of Sucrose in Stabilization of Cut Gerbera Flower Stalk. *Gartenbouwissenschaft*, (47):77-81.

Thakur, M., Sharma, B.P. and Verma, T. (2019). Effect of Different Pulsing Solutions on Postharvest Life of Iris (*Iris orientalis* Mill.). *Int.J.Curr.Microbiol.App.Sci*, **8(5)**: 2116-2124.

Vidhya, S. and Bhattacharjee, S. K. (2002). Floriculture Research Trend in India. *Ind Soc. Ornamental Hort. New Delhi*, pp. 83-86.

Reid, M. S. and Jiang, C. Z. (2012). Postharvest biology and technology of cut flowers and potted plants. *Horticultural reviews*, **(40)**: 1-54.

Asrar, A. W. A. (2012). Effects of some preservative solutions on vase life and keeping quality of snapdragon (*Antirrhinum majus* L.) cut flowers. *Journal of the Saudi Society of Agricultural Sciences*, **11(1)**: 29-35.

Rogers, M. N. (1973). An Historical and Critical Review of Postharvest Physiology Research on Cut Flowers¹. *HortScience*, **8(3)**: 189-194.

Elhindi, K. M. (2012). Evaluation of several holding solutions for prolonging vase-life and keeping quality of cut sweet pea flowers (*Lathyrus odoratus* L.). *Saudi journal of biological sciences*, **19(2)**: 195-202.

Pun, U. K., Yamada, T., Azuma, M., Tanase, K., Yoshioka, S., Shimizu-Yumoto, H. and Ichimura, K. (2016). Effect of sucrose on sensitivity to ethylene and enzyme activities and gene expression involved in ethylene biosynthesis in cut carnations. *Postharvest biology and technology*, **(121)**: 151-158.