

Original Research Article

Determination of crop-coefficients and estimation of evapotranspiration of field pea (*Pisum sativum* L.) using lysimeter and different reference evapotranspiration models for temperate region.

ABSTRACT:

During Rabi 2020-21, a field experiment was conducted at SKUAST-K, Shalimar, Srinagar, focusing on field Pea (*Pisum sativum* L.). The aim was to determine the water requirement and single crop coefficient (K_c) of pea using a lysimeter setup. Four reference evapotranspiration models were employed to calculate the reference evapotranspiration and were then compared with the actual crop evapotranspiration at different growth stages. The K_c values for field pea were 0.50, 0.80, 1.15, and 1.10 during the initial, development, mid-season and late season stages, respectively. The water requirement was found as 239.9 mm for the whole cropping period of the pea. Among the models, the Penman Montieth crop evapotranspiration model exhibited the closest agreement with the corresponding values obtained in the field through water balance study, yielding RMSE, RSR, and NSE values of 0.97, 9.5, and 11.6, respectively. These findings highlight the significance of using lysimeters for eco-hydrological crop evapotranspiration modeling in temperate regions.

Key words: Crop Evapotranspiration, Drainage type lysimeter, Crop coefficient, Pea.

INTRODUCTION:

The accurate assessment of water losses through evapotranspiration by crops is crucial due to limited water resources. Weather stations record climatic parameters such as air temperature, solar radiation, relative humidity, and wind speed, which are used to estimate plant water needs (Incrocci *et al.*, 2014). Reference evapotranspiration (ET_0) is determined by various mathematical models based on these parameters. The crop coefficient (K_c) is obtained by dividing actual crop evapotranspiration (ET_c) measured using lysimeters by the reference evapotranspiration (ET_0) and represents crop-specific water use. Accurate estimation of K_c is essential for determining the irrigation requirements of different crops in diverse climatic conditions (Doorenbos and Pruitt, 1977). Developing a specific crop coefficient (K_c) for field pea is vital for precise irrigation water planning. ~~By utilizing lysimeters to measure crop water use and established methods to calculate reference evapotranspiration rates, daily crop coefficients can be determined.~~ Properly scheduling irrigations based on the averaged water

Comment [H1]: Estimation of crop-coefficients and evapotranspiration of field pea (*Pisum sativum* L.) using lysimeter and empirical models under temperate climate

Comment [H2]: India

Comment [H3]: of this study was

Comment [H4]: empirical models

Comment [H5]: Penman Montieth crop evapotranspiration model

Comment [H6]: estimating

Comment [H7]: delete

requirement and correct timing is crucial to meet the crop's water demands and achieve optimal crop production (Mehta and Pandey, 2016).

Research indicates that gaining a better understanding of actual crop water requirements through modern technologies can lead to **saving** at least 50% of irrigation water (Ragab *et al.*, 2017). Numerous studies have explored various evapotranspiration models in different locations. For instance, Dehghani Sanij *et al.* (2004) assessed four ET₀ models in Karaj, Iran; Bormann (2011) investigated 18 PET models in the German climate; Nag *et al.* (2014) examined 14 models in India; Djaman *et al.* (2015) studied 16 ET₀ models in the Senegal River Valley; and Muniandy *et al.* (2016) tested 26 ET₀ models in Kluang, Malaysia.

Among the empirical models, the Food and Agricultural Organization recommends the Penman-Monteith equation (FAO-PM) as the standard method for estimating ET, requiring meteorological parameters such as temperature, humidity, wind speed, sunshine hours, and net radiation (Allen *et al.*, 1998). However, some researchers have also used simpler empirical models like Hargreaves-Samani, Turc, Blaney-Criddle, as they require fewer meteorological parameters.

In the Kashmir valley, pea is mainly cultivated as a Rabi crop, but in higher altitudes, it is grown as an off-season vegetable during summer. Peas can tolerate temperatures ranging from 7 to 30°C in higher tropical altitudes (Duke, 2012). Being a winter crop, peas can withstand relatively low temperatures, especially during the early stages of growth, but may not survive severe and prolonged frost (Prabhakara *et al.*, 2008). To estimate evapotranspiration accurately, it is essential to develop crop coefficients (K_c) for different models. Based on the above considerations, this experiment was undertaken to determine the crop coefficients (K_c) and estimate evapotranspiration of field pea using four reference evapotranspiration models.

MATERIALS AND METHODS:

During the period from November 2020 to May 2021, a field experiment was conducted on pea crops at Sher-e-Kashmir University of Science and Technology-Kashmir (SKUAST-K), located in Shalimar, Srinagar, Jammu & Kashmir, India. The geographical coordinates of the experimental site are approximately 34°1' N latitude and 74°9' E longitude, with an altitude of 1586 meters above mean sea level. The **climate at the** experimental site is characterized **as** **temperate**, experiencing moderately hot summers and **bitterly** cold winters, with the majority of precipitation occurring as snow during winter. Summer **temperatures** typically **range between** 30°C to 35°C, while winter temperatures can drop as low as -10°C.

Comment [H8]: save

Comment [H9]: and

Comment [H10]: no need

Comment [H11]: by

Comment [H12]: temperate climate

Comment [H13]: extremely

Comment [H14]: temperature

Comment [H15]: ranges from

The recorded annual rainfall was approximately 710 mm. Throughout the year 2020-21, the average relative humidity at the experimental site was 70%. During the Rabi season, the average number of sunshine hours recorded was 4 hours per day.

Comment [H16]: The rainfall was approximately 710 mm, relative humidity was 70% and average number of sunshine periods was 4 hours per day during the study period.

To explore the actual crop evapotranspiration, a field lysimeter setup was established. This method involves monitoring the water inflow and outflow within the crop root zone throughout the crop's growth period. However, certain fluxes like subsurface flow and deep percolation are challenging to accurately assess over short time frames. Consequently, the soil water balance method typically provides estimates of crop evapotranspiration (ET_c) over longer durations (Allen *et al.*, 1998).

Comment [H17]: no need

Comment [H18]: A lysimeter was setup in the experimental field to monitor the water inflow and outflow within the crop root zone throughout the growing period.

Comment [H19]: assess crop evapotranspiration (ET_c)

Comment [H20]: ET_c

Comment [H21]: were

The inflow and outflow variables required in the water balance equation are measured in the lysimeter set-up. The inflow to the field can consist of precipitation and applied irrigation water and water can leave the field through evapotranspiration, surface runoff, seepage, and vertical percolation. Changes in soil moisture storage were measured by soil moisture sampling at different depths of the root zone within lysimeter. The crop evapotranspiration is computed using the following water balance equation:

Comment [H22]: The inflow is the sum of precipitation and applied irrigation water and outflow consists of evapotranspiration, surface runoff, seepage, and vertical percolation.

Comment [H23]: was

$$\Delta S = P + I - ET - DP - HS - R \quad (1)$$

where, ΔS is the change in storage in the root zone (mm), P is precipitation amount (mm), I is irrigation water (mm), ET is actual evapotranspiration (mm), DP is vertical deep percolation (mm), HS is horizontal seepage through bunds (mm) and R is surface runoff (mm).

Comment [H24]: of

Comment [H25]: no need

As the experiments are conducted in a lysimeter, horizontal seepage (HS) is zero and as the soil is not fully filled in the lysimeter, surface runoff (R) is negligible. So, the water balance equation for the lysimeter set-up becomes:

Comment [H26]: were

Comment [H27]: was

Comment [H28]: soil was

Comment [H29]: was

Comment [H30]: Therefore

Comment [H31]: was

$$\Delta S = P + I - ET - DP \quad (2)$$

Comment [H32]: ΔS was

Change in storage (ΔS) is calculated using the initial and final moisture content readings over required time duration. Precipitation (R) data is taken from the meteorological observatory of Agromet Field Unit, SKUAST-K, Shalimar. Evapotranspiration (ET) is estimated using FAO-Penman-Monteith equation (Allen *et al.*, 1998). Irrigation (I) to the crop is measured by calibrated hosepipe.

Comment [H33]: was

Comment [H34]: ET was

Comment [H35]: I was

The evapotranspiration rate from a well-watered reference surface not short of water is referred to as the reference crop evapotranspiration or ET_0 . Table 1 provides a summary of the four most commonly used reference evapotranspiration models. To determine the

Comment [H36]: is known as

suitability of these models for specific agro-climatic conditions, crop reference evapotranspiration has been calculated using local climatic data and modified crop coefficient values. Selecting an appropriate ET_0 estimation method among the various available methods for a particular agro-climatic condition can be challenging.

Comment [H37]: was

Comment [H38]: no need

The concept of crop coefficient (K_c) was first introduced by Jensen in 1968 and further developed by other researchers such as Jensen (1968), Doorenbos and Pruitt (1975), Doorenbos and Pruitt (1977), and Jensen (2011). Determining the K_c value is essential as it represents the crop-specific water use, enabling accurate estimation of irrigation requirements. ~~Various climatological parameters that influence K_c are utilized for estimating reference evapotranspiration (ET_0).~~ To accommodate different growth stages of crops under diverse climatic conditions, Doorenbos and Pruitt (1977) suggested determining stage-wise crop coefficients.

Comment [H39]: of

Comment [H40]: about the calculation of

The Food and Agricultural Organization (FAO) provides standard crop growth stages corresponding to various crops. A numerical procedure is employed to modify the crop coefficient (K_c) values (Allen *et al.*, 1998). Therefore, the modified FAO values of crop coefficients are recommended for different ET_0 models in the agro-climatic conditions of the present study. Specifically, the crop coefficient for the initial stage is denoted as $K_{c\text{ ini}}$, while the coefficients for the mid-season and end stages are referred to as $K_{c\text{ mid}}$ and $K_{c\text{ end}}$, respectively.

Comment [H41]: please include year

Comment [H42]: were used in this study.

Comment [H43]: was

Comment [H44]: were

In order to precisely evaluate various methods, a quantitative assessment procedure has been employed, incorporating error statistics as proposed by Ambrose and Roesch in 1982. The error statistics used for this evaluation include Root Mean Square Error (RMSE), the Ratio of the Root Mean Square Error to the standard deviation of measured data (RSR), and the Nash Sutcliffe Efficiency (NSE) suggested by Moriasi *et al.* in 2007.

Comment [H45]: was

Comment [H46]: (1982)

Comment [H47]: (2007)

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (ET_{\text{obs}} - ET_{\text{cal}})^2} \quad (3)$$

$$NSE = 1 - \frac{\sum_{i=1}^n (ET_{\text{obs}} - ET_{\text{cal}})^2}{\sum_{i=1}^n (ET_{\text{obs}} - ET_{\text{mean}})^2} \quad (4)$$

$$RSR = \frac{\sqrt{\sum_{i=1}^n (ET_{\text{obs}} - ET_{\text{cal}})^2}}{\sqrt{\sum_{i=1}^n (ET_{\text{obs}} - ET_{\text{mean}})^2}} \quad (5)$$

Where, ET_{cal} = calculated ET_c by Models (mm)

ET_{obs} = observed ET_c by lysimeter method (mm)

ET_{mean} = average daily ET_c observed over the season (mm)

The FAO recommended K_e values for various growth stages (Allen *et al.*, 1998) were adjusted to suit the local climatic conditions for different stages. Crop evapotranspiration reflects the crop's water requirement, which is calculated as the product of the specific K_e value for the given period and the corresponding reference evapotranspiration (ET_0).

Table 1: Various models used for computing ET_0

Model	Formula	Reference
Priestley-Taylor	$ET_0 = \alpha + \frac{\Delta}{\Delta + \gamma} (R_n - G)$	Shuttleworth W J, 1992
Penman-Monteith	$ET_0 = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T + 273} U_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34 U_2)}$	Allen <i>et al.</i> , 1998
FAO Pan-Evaporation	$ET_0 = E_{pan} \times K_p$	Allen <i>et al.</i> , 1998
Hargreaves-Samani	$ET_0 = 0.0023 (T_{max} - T_{min}) \sqrt{(T_{mean} + 17.8) R_a}$	Hargreaves and Samani, 1985
Blaney-Criddle	$E_0 = P (0.46T + 8)$	Allen and Pruitt, 1986

Comment [H48]: (1992)

Comment [H49]: (1998)

Comment [H50]: (1998)

Comment [H51]: (1985)

Comment [H52]: (1986)

RESULTS AND DISCUSSION:

Plant moisture depletion from the root zone is primarily influenced by the daily crop evapotranspiration values. The amount of moisture taken up from the root zone corresponds to the crop evapotranspiration. The variability in the moisture content within the effective root zone is largely indicative of the crop evapotranspiration (Evans *et al.*, 1996). The estimated crop evapotranspiration is obtained through a specific model, which is the product of two factors, K_e (crop coefficient) and ET_0 (reference evapotranspiration). The model predicts the crop evapotranspiration for a given period. Figure 1 illustrates the reference evapotranspiration values calculated using different models over the course of one year.

Comment [H53]: Please describe the figure 1

Comment [H54]: study period




Figure 1. Reference evapotranspiration values calculated using different models over the course of one year.

Comment [H55]: Reference evapotranspiration values calculated using different models

Monthly climatic parameters of the study area

The monthly meteorological parameters during pea crop growing season of the experiment i.e. November 2020 to May 2021 is presented in Table 2.

Table 2. Monthly meteorological parameters

Comment [H56]: Please move it to materials and methods section

Month	Temperature (°C)		Relative humidity (%)		Sunshine durations (hrs)	Rainfall (mm)	Wind speed (Km/hr)
	Maximum	Minimum	RH ₁	RH ₂			
Nov	13.7	-1.15	83.1	64.8	2.9	6.14	0.59
Dec	9.4	-3.2	90.3	69.5	2.6	15.0	0.4
Jan	5.8	-5.87	91.5	73.4	1.6	24.6	0.2
Feb	12.6	-0.87	86.7	60.6	4.5	4.58	0.62
Mar	15.0	3.78	81.2	60.7	3.8	15.6	1.62
Apr	18.7	5.3	75.1	48.4	4.9	17.5	1.7
May	24.5	9.4	79.6	54.8	6.2	5.34	2

Crop duration

The pea seeds were sown on 20th of November 2020 and it took 180 days to reach maturity and harvesting stages. The crop duration was divided into four stages as shown in Table 3.

Comment [H59]: Please move it to materials and methods section

The values of K_c vary with different crop growth stages from 0.50 to 1.10 during initial, development, mid and end stages respectively as recommended by FAO.

Comment [H57]: insert comma

Comment [H58]: insert year

Table 3. Duration of crop growth period and crop coefficient (K_c) of pea crop

Crop stages	Crop duration	K_c value
Initial stage	35	0.50
Development stage	45	0.80
Mid-season stage	70	1.15
End-season stage	30	1.10

Evaluation of stage-wise ET_0 using different empirical methods

The stage wise mean reference evapotranspiration at different pea crop growth stages were calculated at experimental field of COAE&T, Shalimar using different empirical methods and it is presented in Table 4. The variation in reference evapotranspiration (ET_0) of same sample during each stage by different methods of determination was because of the various climatic factors that the particular procedure considers. Similar results pertaining to variation in ET_0 values by different methods was reported by Ahmad *et al.*, (2017).

Comment [H60]: The stage wise calculated mean reference evapotranspiration (ET_0) at different pea crop growth stages is presented in Table 4.

Comment [H61]: Please describe the table

Comment [H62]: Variation in ET_0 at each growth stage under different methods was because of the use of various climatic factors.

Table 4. Mean Reference evapotranspiration at different stages using different empirical methods

Stages	Penman-Monteith (mm/day)	Hargreaves (mm/day)	Blaney-Criddle (mm/day)	Open pan (mm/day)
Initial	1.43	1.19	1.04	1.24
Development	1.35	1.09	0.87	1.15
Mid-stage	2.74	2.59	2.44	2.60
End-stage	4.59	4.54	4.38	4.51

Comment [H63]: ET_0 at different growth stage

The mean Crop evapotranspiration (ET_c) at different stages of pea crop growing season was calculated by different empirical methods by multiplying the reference evapotranspiration with crop coefficients recommended in FAO. ~~Also, crop evapotranspiration was calculated by lysimeter using water balance equation at each crop growth stage.~~ The stage wise mean crop evapotranspiration of pea crop growing season is shown in Table 5. Similar results pertaining to variation in ET_c , values by different methods was reported by Ahmad *et al.*, (2017).

Comment [H64]: c

Comment [H65]: by

Comment [H66]: please describe the Table 5

Table 5. Mean Crop evapotranspiration (ET_c) by using Lysimeter and different empirical methods at different stages

Stages	Penman-Monteith (mm/day)	Hargreaves (mm/day)	Blaney-Criddle (mm/day)	Open Pan (mm/day)	Lysimeter (mm/day)
Initial	0.71	0.59	0.52	0.62	0.66
Development	1.08	0.87	0.69	0.95	1.01
Mid-stage	3.15	2.97	2.80	2.99	3.11
End-stage	5.04	4.99	4.81	4.96	4.97

Comment [H67]: Mean crop evapotranspiration (ET_c) at different growth stages

Regression analysis between Crop evapotranspiration (ET_c) by Lysimeter and by different empirical methods

To show a more clear and convincing picture of analysis a single linear regression ($y = mx + c$) or commonly called straight line equation is used for each of the estimation in all stages. In the graphical representation, X-axis was crop evapotranspiration (ET_{Obs}) by lysimeter method and Y-axis was crop evapotranspiration by a particular method that we selected.

Comment [H68]: Relationship between crop evapotranspiration (ET_c) by Lysimeter and by different empirical methods

Agreement between the calculated (ET_{Cal}) and observed evapotranspiration (ET_{Obs}) values was quantitatively evaluated using the Nash Sutcliffe efficiency (NSE), the ratio of the root mean square error to the standard deviation of measured data (RSR), and root mean square error (RMSE). Based on the above statistical evaluation the performance of the different empirical methods in estimation of crop evapotranspiration for each crop stage varied from 'Good to Very Good'. The comparison of calculated evapotranspiration (ET_{Cal}) by empirical methods and observed evapotranspiration (ET_{Obs}) values using lysimeter during each growth stage was rated graphically from Fig. 2 to Fig. 5.

Comment [H69]: no need

With the course of analysis of result, it was found that Penman-Monteith method has a close relationship with lysimeter method. The RMSE, RSR, and NSE values indicated that the Penman-Monteith method performed 'Very Good' in estimating the evapotranspiration of pea crop during each crop growth stage. With the reference from the data it was concluded that the Penman-Monteith method of determination of reference evapotranspiration would be adopted as best method. The RMSE, NSE and RSR values indicated that the Hargreaves method performed "Very Good" in estimating the evapotranspiration of pea crop during initial, development, mid and End stage. Hargreaves method can be the best substitute in similar results pertaining to the performance evaluation of different empirical methods.

Comment [H70]: The relationship between calculated evapotranspiration by empirical methods and observed evapotranspiration using lysimeter during each growth stage is presented in Figure 2 to Figure 5.

Comment [H71]: is

Comment [H72]: where are the values?

Comment [H73]: very unclear

Comment [H74]: not clear




Fig.2. ET_c Lysimeter versus ET_c by Penman-Monteith method




Fig 3. ET_c Lysimeter versus ET_c by Hargreaves method




Fig. 4. ET_c lysimeter versus ET_c by Blaney-criddle method




Fig. 5. ET_c Lysimeter versus ET_c by Open pan method

Conclusion

Spatio-temporal variability of rainfall events coupled with their irregularity and amount received during crop growing seasons is a major concern for enhancing agricultural production in temperate regions of India. Such situation requires provision of irrigation facilities for enhancing agricultural production in the region. Moreover, under irrigated conditions, efforts should be made to apply judicious amount of water which lowers the cost of production and save the valuable water resources of the country for enhancing irrigation water use efficiency. To accomplish this, proper irrigation schedules need to be devised, proclaimed and adopted by the stakeholders for enhancing the water productivity. It is observed that about 99% of the water uptake by plants from soil is lost as evapotranspiration. Therefore, determination of daily crop evapotranspiration at different crop growth stages of pea crop was undertaken in this study using non-weighing drainage type field lysimeter which is the direct method of estimating evapotranspiration. Penman-Monteith and three other models viz., Hargreaves, Blaney-Criddle and Open Pan methods were used for estimation of reference evapotranspiration (ET_0). FAO Penman-Monteith Model has been found to perform better than other reference evapotranspiration (ET_0) models in predicting crop evapotranspiration (ET_c). The total crop evapotranspiration from the lysimeter study during the crop growing period was 230.4 mm. In order to minimize the loss of water and to precisely meet the crop water demand to produce greater yields with enhanced water use efficiency crop water management practices are essential.

Comment [H75]: no need

Comment [H76]: delete

Comment [H77]: In order to minimize the loss of water and to precisely meet the crop water demand for better yields with enhanced water use efficiency, crop water management practices are essential.

References

- Ahmad L., Parvaze S., Mahdi S. S., Dekhle B. S., Parvaze S., Majid M. and Wani F.S. (2017). Comparison of Potential Evapotranspiration Models and Establishment of Potential Evapotranspiration Curves for Temperate Kashmir Valley. *Current Journal of Applied Science and Technology* **24**(3): 1-10.
- Allen R.G. and Pruitt W.O., (1986). Rational use of the FAO Blaney-Criddle formula. *Journal of Irrigation and Drainage Engineering*, *112*(2):139-155.
- Allen R. G., Pereira L. S., Raes D. and Smith M. (1998). Crop evapotranspiration: guidelines for computing crop water requirements. Irrigation and Drainage Paper **56**. UN-FAO, Rome, Italy.
- Ambrose J. R. B. and Roesch S. E. (1982). Dynamic estuary model performance. *Journal of Environment Engineering Division* **108**: 51–7.
- Bormann H. (2011). Sensitivity analysis of 18 different potential evapotranspiration models to observed climatic change at German climate stations. *Climatic Change*, *104*(3-4):729-753.
- Dehghani Sanij H., Yamamoto T. and Rasiah V. (2004). Assessment of evapotranspiration estimation models for use in semi-arid environments, *Agricultural water management* **64**: 91-106
- Djaman K., Balde A.B., Sow A., Muller B., Irmak S., N'Diaye M.K., Manneh B. Moukoubi Y.D., Futakuchi K. and Saito K., (2015). Evaluation of sixteen reference evapotranspiration methods under sahelian conditions in the Senegal River Valley. *Journal of Hydrology: regional studies*, *3*:139-159.
- Doorenbos J. and Pruitt W. O. (1975). Guidelines for predicting crop water requirements, Irrigation Drainage Paper no. 24, FAO-ONU, Rome, Italy, 168 p.
- Doorenbos J. and Pruitt W. O. (1977). Crop Water Requirements. FAO Irrigation and Drainage Paper 24. FAO, Rome, Italy, 144 p.
- Duke J. (2012). Handbook of legumes of world economic importance. Springer Science & Business Media.
- Evans R., Sneed R. E. and Cassel D. K. (1996). Irrigation scheduling to improve water- and energy- efficiencies. Pub. No., AG 452- 4, North Carolina Cooperative Extension Service. Evaporation. *Journal of Hydrology* **45**: 276–84.
- Hargreaves G. L. and Samani Z. A. (1985). Reference crop evapotranspiration from temperature. *Appl. Eng. Agric. Trans. ASAE* **1** (2): 96–9.

- Incrocci L., Incrocci G., di Vita A., Pardossi A., Bibbiani C., Marzioletti P., Balendonck J. (2014). Scheduling irrigation in heterogeneous container nursery crops. *Acta Hort.*, 1034: 193-200.
- Jensen M. E. (1968). Water consumption by agricultural plants. Chapter-1. (In). *Water Deficits and Plant Growth*, Vol. II pp 1–22 Kozlowski T T (Ed). Academic Press, New York.
- Mehta R. and Pandey V., (2016). Crop water requirement (ET_c) of different crops of middle Gujarat. *Journal of agrometeorology*, 18(1):83-87.
- Moriasi D. N., Arnold J., Van Liew M. W., Bingner R.L., Harmel R.D. and Veith T.L. (2007). Model evaluation guidelines for systematic quantification of accuracy in watershed simulations. *Transaction of the ASAE* **50**: 885-900.
- Muniandy J.M., Yusop Z. and Askari M., (2016). Evaluation of reference evapotranspiration models and determination of crop coefficient for *Momordica charantia* and *Capsicum annum*. *Agricultural Water Management*, 169:77-89.
- Nag A., Adamala S., Raghuwanshi N.S., Singh R. and Bandyopadhyay A. (2014). Estimation and ranking of reference evapotranspiration for different spatial scales in India. *J Indian Water Resour Soc*, 34(3):35-45.
- Prabhakara, C., Iacovazzi J. R., Yoo J. M., Kim K. M., and Bell T.L. (2008). A method to estimate rain rate over tropical oceans with the TRMM microwave imager radiometer. *Journal of the Meteorological Society of Japan* **86**(1): 203-212.
- Ragab R., Evans J.G., Battilani A. and Solimando D., (2017). Towards accurate estimation of crop water requirement without the crop coefficient K_c : New approach using modern technologies. *Irrigation and Drainage*, 66(4):469-477.
- Shuttleworth W. J. (1992). *Evaporation: Handbook of Hydrology*. Maidment D R, (ed.). McGraw-Hill, New York.