

Original Research Article

Impact of long-term use of fertilizers and manure on distribution of soil-inorganic phosphorus fractions in a Vertisol

Abstract

~~The present study was conducted under All India Coordinated Research Project on Long Term Fertilizer Experiment at the Research Farm Department of Soil Science and Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh, India.~~ The investigation was carried out to assess the distribution of P fractions in a Vertisol. Eight treatments were imposed with four replications in simple randomized block design. The treatments were applied in combination of different doses of fertilizer viz T₁ (50% NPK), T₂ (100% NPK), T₃ (150% NPK), T₄ (100% N), T₅ (100% NP), T₆ (100% NPK+FYM), T₇ (100% NPK-S) and T₈ (control). The various forms of phosphorus in the soil were observed in the following order: Ca-P > Fe-P > Al-P > Occluded-P > Saloid-P. The Ca-P content was highest when 100% NPK fertilizer with farmyard manure (FYM) were used together. The analysis showed that the presence of saloid-P significantly influences the availability of phosphorus in Vertisol soil. Based on the current investigation, it is recommended to use integrated nutrient management, which includes both 100% NPK fertilizer with FYM, to prevent phosphorus deficiency in soybean cultivation in Vertisol.

Keywords: *LTFE, Inorganic Fertilizer, Phosphorus fraction, Vertisol.*

Introduction

Phosphorus is an important nutrient for maintaining the safety of the world's food supply, but its availability is limited and non-renewable, which makes its efficient use even more important (Roberts and Johnston., 2015). Phosphorus in soils is present both in organic as well as in inorganic compounds with aluminium (Al), iron (Fe), and calcium (Ca). These substances have an impact on the availability of phosphorus to plants (Tan, 1998). The supply of labile inorganic P (fertilizer P) is controlled by phosphorus sorption, and the isotherm aids in forecasting how effectively crops will use that labile P. It is generally known that phosphate adsorption is influenced by a variety of soil characteristics, including clay concentration, CaCO₃, organic matter, Al and Fe content, CEC, and the sources of P that are added to the soil. The long-term fertilizer experiment was conducted with major attention to the yield performance of crops that are under balanced and imbalanced manure and fertilizer applications in the soil fertility management of a cropping sequence (Katyal 2015). In a permanent manorial experiment. The long-term fertilizer experiment provides very good opportunity to address the issue. Keeping this in view, a study of phosphorus fractions under long term use of fertilizers and manure in a Vertisol of Jabalpur, Madhya Pradesh, India.

Material and Methods

Experimental site, Climate and Soil characteristics

The All India Coordinated Research Project on Long Term Fertilizer Experiment, funded by ICAR, commenced in 1972 at the Department of Soil Science, Jawaharlal Nehru Krishi Vishwa Vidyalaya in Jabalpur, Madhya Pradesh, India. The Jabalpur region has a distinct semi-arid and subtropical climate with dry summers and cold winters. The experimental site is approximately 393 meters above mean sea level at 23° 10' N latitude and 79° 57' E longitude. The average temperature throughout the winter season, which runs from November to February, is 4°C to 33°C, with a relative humidity range of 70% to 90%. Generally speaking, the months of March to June are dry and warm. The temperature can rise as high as 45°C in the summer and fall to below 5°C or 10°C in the winter. Mid-June through mid-September is the monsoon season. The area receives 1350 mm of rain on average each year. The experimental field featured medium black soil from the Kheri series, classified as fine montmorillonitic hyperthermic Typic Hapluster, with a soil pH of 7.6, electrical conductivity of 0.18 dS m⁻¹ (measured at a 1:2.5 soil-to-water ratio), soil organic carbon content of 5.7 g kg⁻¹, available nitrogen at 193 kg ha⁻¹, available phosphorus at 7.6 kg ha⁻¹, and available potassium at 370 kg ha⁻¹. This soil had a clay texture.

Table 1. Initial soil properties of the experimental site

| Soil properties | Value | Reference |
|-----------------------------------|--------------------------|----------------------------|
| pH1:2.5 | 7.60 | Jackson (1973) |
| Electrical conductivity (EC1:2.5) | 0.18 dS m ⁻¹ | Jackson (1973) |
| Organic carbon | 5.7 g kg ⁻¹ | Walkley and Black (1934) |
| Available N | 193 kg ha ⁻¹ | Subbiah and Asija (1956) |
| Available P | 7.60 kg ha ⁻¹ | Olsen <i>et al.</i> (1954) |
| Available K | 370 kg ha ⁻¹ | Muhr <i>et al.</i> (1965) |
| Various phosphorus fractions | | |
| Saloid-P | 2.6 mg kg ⁻¹ | Peterson and Corey (1966) |
| Al-P | 26.2 mg kg ⁻¹ | |
| Fe-P | 41.5 mg kg ⁻¹ | |
| Occluded-P | 36.5 mg kg ⁻¹ | |
| Ca-P | 87.5 mg kg ⁻¹ | |

Treatments details

The experiment comprised eight different treatments:

T₁ - 50% NPK

T₂ - 100% NPK

T₃ - 150% NPK

T₄ - 100% NP

T₅ - 100% N

T₆ - 100% NPK + 5 t FYM ha⁻¹ (Farmyard Manure per hectare)

T₇ - 100% NPK – S (Sulphur-free)

T₈ – Control

Each treatment was replicated four times in a randomized block design. For soybean cultivation, the recommended fertilizer doses based on the initial soil test values were 20 kg N, 80 kg P₂O₅, and 20 kg K₂O per hectare. Urea, single superphosphate, and muriate of potash were used as the sources of nitrogen, phosphorus, and potassium, respectively. In the sulphur-free treatment (T₇), diammonium phosphate (DAP) was used instead of single superphosphate as the source of phosphorus. Farmyard manure was applied at a rate of 5 tons per hectare per year, exclusively during the kharif season for soybean crop. All recommended agricultural practices were followed to cultivate soybeans as per standard recommendations.

Phosphorus fractions in soil

The procedure of Chang and Jackson (1957) as modified by Peterson and Corey (1966) was followed for fractionation of soil phosphorus. The sequence of Saloid-P, Al-P, Fe-P, Occluded-P and Ca-P from each sample was passed through a 60-mesh sieve.

The soil extractant for various fractions in sequence were as follow: -

- ❖ Saloid-P extracted by 1 N NH_4Cl solution
- ❖ Al-P extracted by 0.5N NH_4F solution (buffered at pH 8.2)
- ❖ Fe-P extracted by 0.1 N NaOH solution
- ❖ Occluded-P extracted by 0.1N NaOH solution
- ❖ Ca-P extracted by 0.5N H_2SO_4 solution

Statistical analysis of the experimented data

The data were statistically analysed and correlations amongst various parameters were computed as per standard procedure described by (Panse and Sukhatme, 1970).

Result and Discussion

Soil pH

The pH values ranged from 7.48 to 7.58, indicating that continuous cropping and fertilizer use did not have a significant adverse effect on the soil pH. This could be attributed to the high buffering capacity of the soil, as suggested by (Dwivedi and Dwivedi, 2015). The findings align with previous research by Raghuwanshi *et al.*, (2016).

Over the course of 50th years of continuous fertilizer additions and intensive cropping, no significant alteration in soil pH was observed concerning depth. The soil's remarkable buffering capacity, as noted by Suman *et al.*, (2017), allowed it to effectively maintain its pH levels despite these agricultural practices.

Soil EC

The initial soil electrical conductivity (EC) values, ranging from 0.15 to 0.20 dS/m, remained largely unchanged throughout the study. The application of various fertilizer and manure doses did not have a significant impact on the soil's EC in a Vertisol. Even the continuous use of synthetic fertilizers over an extended period did not noticeably alter the soil's EC. The combined use of farmyard manure (FYM) and fertilizers resulted in minimal changes, likely attributed to the addition of organic manure, which enhanced the soil's buffering capacity, as reported by Gupta *et al.*, (2019).

After the 50th years of continuous fertilizer addition, the soil's electrical conductivity (EC) did not show significant changes concerning depth. These findings align with the research conducted by Panwar *et al.*, (2017), Suman *et al.*, (2018), and Khandagle *et al.*, (2019). All these studies reported similar results, indicating no considerable alterations in soil EC due to the continuous application of fertilizers.

Soil organic carbon

The data showed that the lowest organic carbon content 4.68 g kg^{-1} was noted in control. However, the organic carbon content improved significantly with proportionate increment in fertilizer addition at 50%NPK (6.28 g kg^{-1}), 100%NPK (7.27 g kg^{-1}) and 150%NPK (8.02 g kg^{-1}) doses. This finding appeared to be due to enhanced root development of crop resulting higher residues as a result of intensive farming with continuous fertilizer applications. These results are also in agreement with the finding of Parihar *et al.* (2010) and Patel *et al.*, (2018).

The application of 100% NPK+FYM, there was a noticeably higher level of soil organic carbon (6.89 g kg^{-1}) reported by Birla *et al.*, (2015) showed that organic manures (FYM) were discovered to be a practical option for improving soil organic carbon and nutrient turnover,

therebyimproving nutrient availability in the soil, maintaining soil quality, and achieving sustainable productivity. Similar results were also reported by Meshram *et al.*, (2018) and Bagde*et al.*, (2023).

UNDER PEER REVIEW

Table-2. Impact of long-term use of fertilizers and manure on soil properties in surface and sub surface soils

| Treatments | Soil pH | | EC (dSm ⁻¹) | | OC (gkg ⁻¹) | | Available N (kg ha ⁻¹) | | Available P (kg ha ⁻¹) | | Available K (kg ha ⁻¹) | |
|-----------------------|--------------|--------------|-------------------------|--------------|-------------------------|-------------|------------------------------------|--------------|------------------------------------|-------------|------------------------------------|--------------|
| | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm | 0-15cm | 15-30 cm | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm |
| 50% NPK | 7.50 | 7.62 | 0.15 | 0.16 | 6.28 | 4.86 | 226 | 197 | 19.97 | 19.37 | 260 | 250 |
| 100% NPK | 7.55 | 7.63 | 0.17 | 0.18 | 7.27 | 5.83 | 292 | 220 | 33.64 | 30.12 | 302 | 286 |
| 150% NPK | 7.58 | 7.66 | 0.20 | 0.21 | 8.02 | 6.94 | 331 | 280 | 36.72 | 34.18 | 326 | 310 |
| 100% NP | 7.53 | 7.63 | 0.16 | 0.16 | 6.74 | 5.32 | 267 | 205 | 29.71 | 27.73 | 252 | 243 |
| 100% N | 7.48 | 7.52 | 0.15 | 0.16 | 5.38 | 4.45 | 219 | 178 | 9.89 | 8.76 | 248 | 239 |
| 100% NPK + FYM | 7.54 | 7.62 | 0.16 | 0.17 | 8.94 | 7.11 | 340 | 285 | 37.53 | 35.62 | 336 | 315 |
| 100% NPK – S | 7.56 | 7.64 | 0.17 | 0.18 | 7.21 | 5.43 | 275 | 210 | 30.87 | 28.18 | 296 | 278 |
| Control | 7.51 | 7.56 | 0.15 | 0.16 | 4.68 | 4.01 | 190 | 160 | 8.91 | 8.03 | 244 | 237 |
| S.Em. ± | 0.048 | 0.036 | 0.015 | 0.014 | 0.28 | 0.24 | 8.96 | 7.59 | 0.76 | 0.69 | 8.24 | 8.68 |
| C.D. (p=0.05) | NS | NS | NS | NS | 0.85 | 0.73 | 26.4 | 22.33 | 2.24 | 2.05 | 24.25 | 25.55 |

NPK: The per cent amount of nitrogen, phosphorus and potassium applied through inorganic fertilizers.

*FYM (contains 0.53%N,0.30%P₂O₅and0.63K₂O) was applied 5 t ha⁻¹ to soybean every year 15-20 days before sowing;

#DAP was used instead of SSP as a source of P.

NS: Non-Significant

Available nitrogen in soil

The highest value of available N at 0-15 cm depth of soil was recorded in 100% NPK+FYM (340 kg ha⁻¹) treatment. However, the lowest value of available N content was observed in control (190 kg ha⁻¹) followed by 100% N (219 kg ha⁻¹). The data presented in Table 2. indicated a decrease in the available nitrogen (N) content in the soil as the depth increased. The highest available N content, measuring 285 kg ha⁻¹, was recorded in the treatment with 100% NPK + FYM at a soil depth of 15-30 cm. Observations revealed that higher nitrogen levels were observed in the surface soil, which gradually decreased with depth. The rate of decline was more pronounced from the surface to the subsurface layers, as reported by Singh *et al.*, (2013). This pattern could be explained by the presence of higher root biomass in the rhizospheric soil layer. Similar findings were also reported by Khandagleet *et al.*, (2020).

Available phosphorus in soil

The available P was changed from 8.91 kg ha⁻¹(control) to 19.97 kg ha⁻¹, 33.64 kg ha⁻¹, and 36.72 kg ha⁻¹ in the 50% NPK, 100% NPK, and 150% NPK treatments, respectively. Further, it was the highest with 100% NPK+FYM (37.53 kg ha⁻¹). while the lowest values were observed in the control plot (8.91 kg ha⁻¹), followed by the 100% N treatment (9.89 kg ha⁻¹). This indicates that higher doses of NPK fertilizers resulted in increased available P content in the 0-15 cm depth of soil. Similar results were reported by Dubey *et al.*, (2015) and Birla *et al.*, (2015). Moreover, the most elevated P content was observed in treatments involving the combined use of fertilizer and farmyard manure (FYM) at a rate of 36.72 kg ha⁻¹, highlighting the advantageous impact of FYM on P mineralization within the soil to a greater extent by Dwivedi *et al.*, (2019). The data also accentuated that P accumulation was more pronounced near the soil surface compared to deeper layers.

Available potassium in soil

The initial value in (1972) of available K was 370 kg ha⁻¹. The available potassium (K) content in the soil at the 0-15 cm depth showed a progressive increase with the successive addition of fertilizers, starting from 50% NPK (260 kg ha⁻¹), then 100% NPK (302 kg ha⁻¹), and 150% NPK (326 kg ha⁻¹) treatments, respectively. Notably, the treatment that received 100% NPK along with farmyard manure (FYM) showed the highest available K content (336 kg ha⁻¹) among all treatments. On the other hand, the control plot had the minimum available K content (244 kg ha⁻¹), followed by the 100% N treatment (248 kg ha⁻¹) at the same depth. Reported that 100% NPK+FYM treatments followed by 150% NPK treatments were related to the greatest available K status in soil. Similar results were also reported by Patidar *et al.*, (2021) and Pathariyaet *et al.*, (2022) In addition to the direct addition to the soil's accessible K fractions, the application of organic manure may have decreased K fixation and consequently raised K content due to the interaction of organic matter with clay. Inwati *et al.*, (2022) similarly reported comparable outcomes.

Phosphorus fractions in soil

Saloid-P form

The effect of continuous cropping and fertilizer applications on content of Saloid-P form revealed that (table 3) the lower content was recorded in control (4.39 mg kg⁻¹) followed by 100% N (5.27 mg kg⁻¹) which could be because of the fact that without application of fertilizer and or without P inclusion in fertilizer schedule has obviously resulted in its lower content. On the other hand, successive additions of P from 50% NPK 100% NPK and 150% NPK resulted in proportionate increase in Saloid-P in soil. This might be due to proportionate higher transformation of P from both inherent and applied sources with soil system into Saloid-P as suggested by Badrinath *et al.*, (2005). Nonetheless, the most substantial content emerged when 100% NPK was administered alongside FYM, underscoring the collaborative effect of organic manure in expediting the conversion rate into soil-resident Saloid-P, consequently rendering it available for subsequent uptake by cultivated plants.

A diminishing pattern in the Saloid-P configuration was noted as depth increased, with the greatest proportion of Saloid-P localized near the surface and gradually diminishing towards deeper soil layers. This phenomenon could be attributed to the limited mobility of Saloid-P, impeding its downward migration, as elucidated in studies by Dubey *et al.*, (2016) and Sudhakaran *et al.*, (2018).

Aluminium bound Phosphorus (Al-P)

Application of increasing doses of fertilizer increased the content of Al-P form in the soil (table 3). While it declined when P addition was committed there by lower content was noted in control (9.02 mg kg⁻¹) and 100% N (12.42 mg kg⁻¹) due to absence of addition of P. Similar results were reported by Sood and Bhardwaj, (1992). In this regard, Fe-P and Al-P were found to be predominant in the surface soil horizon, according to Ganesh *et al.*, (1993).

Al-P content exhibited a gradual decrease with increasing depth, with the most significant buildup observed at the surface. Corresponding outcomes were also documented in the study by Dubey *et al.*, (2016).

Iron bound phosphorus (Fe-P)

The data of Fe-P form was increased with successive higher P fertilizer application over without additions i.e., in control (19.17 mg kg⁻¹) and 100% N (20.08 mg kg⁻¹) which could be due to higher transformation of added fertilizer P in soil equilibrium to Fe-P retention in soil. It has also been noted that a significantly higher content of Fe-P was observed with successively higher addition of P from optimal to super optimal dose which might be due to the differential conversion of applied P in Fe-P form as a result of varying transformation (Sarkar *et al.*, 2014). In this regard, Fe-P and Al-P were found to be predominant in the surface soil horizon, according to Ganesh *et al.*, (1993).

Elevated levels of Fe-P were observed near the surface, gradually decreasing as depth increased. This phenomenon may arise from the potential movement of both phosphorus and iron ions, particularly limiting their concentration in the lower layers. Comparable findings were documented by Badrinath *et al.*, (2005).

Occluded phosphorus (Occl-P)

The data on content of Occl-P form (table 3) that intensive cropping with continuous use of fertilizer over 50th year has resulted in appreciable build-up of Occl-P form when increasing rates of fertilizer were applied over successive cropping while a declining trend was noted with the imbalance applications. Similar results have been noted by Tiwari *et al.*, (2019). It has seen that application of phosphatic fertilizers significantly increased occluded P fractions over control. Reported by Bincy *et al.*, (2023).

A decrease in the Occluded-P condition was observed as depth increased, with the greatest concentration occurring at the surface. This suggests that phosphorus remains relatively immobile within the soil's lower layers, as noted by Dubey *et al.*, (2016).

Calcium bound phosphorus (Ca-P)

The result showed that successive addition of fertilizer containing P progressively increased the content of Ca-P form over without application in an intensified cropping for the last 50th years particularly in black soils (Table 3). It has also been observed amongst all inorganic forms, the content of Ca-P ranks highest which was an indication of the fact that Ca-P form contributed to the major source of P in black soil as reported by Roy *et al.*, (2016). Lower Ca-P content was associated in the control and 100% N treatments where P fertilizer was not included. These finding are also in agreement with Sudhakaran *et al.*, (2018). The application of 100%NPK along with FYM recorded significantly higher Ca-P during the incubation period. (Bincy *et al.*, 2023)

Minor fluctuations in the Ca-P form were observed with changing depth. This occurrence may be attributed to the prevalence of calcium ions exhibiting high CEC, thereby promoting the substantial generation and development of Ca-P compounds. Comparable findings were documented in a study by Badrinath *et al.*, (2005).

Table-3. Impact of long-term use of fertilizers and manure on phosphorus fractions in surface and subsurface soils

| Treatment | Fractions of phosphorus mg kg ⁻¹ | | | | | | | | | |
|----------------|---|---------|--------|---------|--------|---------|------------|---------|--------|---------|
| | Saloid-P | | Al- P | | Fe- P | | Occluded-P | | Ca- P | |
| | 0-15cm | 15-30cm | 0-15cm | 15-30cm | 0-15cm | 15-30cm | 0-15cm | 15-30cm | 0-15cm | 15-30cm |
| 50% NPK | 12.55 | 10.40 | 23.97 | 21.27 | 31.28 | 25.94 | 8.25 | 6.46 | 110.76 | 101.94 |
| 100% NPK | 16.86 | 13.64 | 32.06 | 27.25 | 47.18 | 37.57 | 11.09 | 8.34 | 141.98 | 133.92 |
| 150% NPK | 18.20 | 15.58 | 36.48 | 31.97 | 54.39 | 44.32 | 13.67 | 10.24 | 188.06 | 179.07 |
| 100% NP | 16.17 | 12.85 | 30.12 | 25.81 | 45.49 | 38.88 | 9.78 | 7.55 | 136.52 | 128.01 |
| 100% N | 5.27 | 4.65 | 12.42 | 11.03 | 20.08 | 13.92 | 6.41 | 4.78 | 98.81 | 92.36 |
| 100% NPK + FYM | 18.87 | 16.31 | 37.12 | 32.27 | 51.55 | 45.12 | 14.36 | 11.30 | 190.12 | 181.17 |
| 100% NPK – S | 16.59 | 13.17 | 30.90 | 26.88 | 46.36 | 39.28 | 10.58 | 7.85 | 138.83 | 130.65 |
| Control | 4.39 | 3.18 | 9.02 | 7.26 | 19.17 | 11.86 | 5.02 | 3.49 | 95.06 | 91.35 |
| S.Em. ± | 0.81 | 0.76 | 0.97 | 0.51 | 2.63 | 2.63 | 0.35 | 0.38 | 7.62 | 7.04 |
| C.D. (p=0.05) | 2.40 | 2.25 | 2.86 | 1.51 | 7.76 | 7.74 | 1.04 | 1.12 | 22.43 | 20.72 |

NPK: The per cent amount of nitrogen, phosphorus and potassium applied through inorganic fertilizers.

*FYM (contains 0.53%N, 0.30%P₂O₅ and 0.63K₂O) was applied 5 t ha⁻¹ to soybean every year 15-20 days before sowing;

#DAP was used instead of SSP as a source of P.

Conclusion

The findings from a long-term experiment on soybean cultivation showed that over the past 50th years, the application of both inorganic fertilizers and organic manure (FYM) led to an increase in all forms of phosphorus (P) in both the topsoil and subsoil. The balanced use of nutrients (100% NPK) and its combination with FYM (100% NPK+FYM) had a positive effect on P content and its various forms in the soil. The study found that the relative contribution of various P forms in the soil followed this order: Ca-P > Fe-P > Al-P > Occluded P > Saloid-P in both the surface and sub-surface soils.

References

- Badrinath MS, Niranjana KS and Nandkishore TG. 2005. Distribution and fractionation of phosphorus in soils of different agro-climatic zones of Karnataka (India). *Soil Sci. Soc. Am. J.* 43: 523-528.
- Bagde V, Dwivedi BS, Dwivedi AK, Thakur R and Nagwanshi A. 2023. Effect of Long-Term Fertilizer and Farm yard Manure Application on Soil Phosphorus Fractions in a Vertisol under Soybean–Wheat Cropping Sequence. *Journal of the Indian Society of Soil Science*,71. (2): 207-216.
- Bincy, B, Ravi, MV and Latha H. 2023. Phosphorus Fractionation in Submerged Black Soil as Influenced by Different Phosphatic Fertilizers. *International Journal of Economic Plants*, 10(May, 2), 114-121.
- Birla K, Khamparia NK, Thakur RK, Dwivedi, BS and Sawarkar SD. 2015. Profile Distribution of Various Fractions of Sulphur as Influenced by Long Term Application of Fertilizers and Manure in Vertisol. *Green Farming*, 6 (4): 753-756.
- Chang SC and Jackson ML. 1957. Solubility product of iron phosphate. *Soil Science Society of America Journal*. 1957 May;21(3):265-269.
- Dubey L, Dwivedi AK and Dubey M. 2015. Long term application of fertilizer and manures on physico-chemical properties and N, P and K uptake in soybean-wheat cropping system. *Plant Archives* 15(1): 143-147.
- Dubey L, Dwivedi BS, Dwivedi AK and Thakur RK. 2016. Effect of long-term application of fertilizer and manure on profile distribution of various phosphorus fraction in Vertisol. *Green farming*, 7 (2):365-370.
- Dwivedi AK and Dwivedi BS. 2015. Impact of long-term fertilizer management for sustainable soil health and crop productivity: Issues and challenges. *JNKVV Res. J.* 49(3): 387-399.
- Dwivedi BS, Sharma A, Dwivedi AK and Thakur RK. 2019. Response of phosphorus application on productivity of wheat at farmer field. *Universal Journal of Agricultural Research* 7 (1):20-24.
- Ganesh A, Raveendran K, and Khilar KC. 1993. Influence of mineral matter on biomass pyrolysis characteristics. *Fuel*. 74(12): 1812-1822.
- Gupta A, Dwivedi AK, Nagwanshi A, Dwivedi BS, Vishwakarma AK. 2019. Impact of Long-Term Application of Inorganic Fertilizer and Farm Yard Manure on Productivity of Soybean in Vertisol. *Bulletin of Environment Pharmacology and Life Sciences*. 8(4):116-132.
- Inwati DK, Dixit BK, Dwivedi BS, Thakur R, Sharma A and Singh V. 2022. Effect of long-term fertilizer and FYM application on soil potassium fractions in a Vertisol under soybean-wheat cropping system. *The Pharma Innovation Journal*, 11(8): 2089-2093.
- Jackson ML. 1973. *Soil chemical analysis*, pentice hall of India Pvt. Ltd., New Delhi, India. 498:151-154.
- Katyal JC. 2015. Withering soil science education revival and resurgence boosters. *J Indian soc. Soil sci.* 63 ((1): 53-63.
- Khandagle A, Dwivedi BS, Aher SB, Dwivedi AK, Yashona DS and Jat D. 2019. Effect of long-term application of fertilizers and manure on soil properties. *Journal of Soils and Crops*, 29(1): 97-104.
- Khandagle A, Dwivedi, BS, Dwivedi, AK, Panwar S and Thakur RK. 2020. Nitrogen fractions under long-term fertilizer and manure applications in soybean – wheat rotation in a Vertisol. *Journal of the Indian Society of Soil Science*, 68(2): 186-193.
- Meshram MK, Dwivedi BS, Naik KR, Thakur RK and Keram KS. (2018). Impact of organic and inorganic sources of nutrients on yield, nutrient uptake, soil fertility and economic performance of rice in a Typic Haplustert. *Journal of Soils and Crops* 28 (1): 31-36.
- Muhr, GR, Datta NP, Subaramany HS, Leley VK and Dunahue RL .1965. *Soil testing*. India Asian Press. New Delhi.
- Olsen SR, Cole CV, Watanabe FS and Dean LA. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *USDA Circular 939*. U.S. Government Printing Office, Washington D.C.

- Panse VG and Sukhatme SV. 1970. Statistical method for agricultural workers. ICAR publication.
- Panwar S, Dwivedi AK, Dwivedi BS and Nagwanshi A. 2017. Distribution of zinc fractions as influenced by long-term application of fertilizers and manure in a Vertisol. *International Journal of Chemical Studies* 5(6): 1931-1934.
- Parihar, SK, Dwivedi, BS, Khan, IM and Tiwari RK. 2010. Effect of integrated nutrient management on yield and economics of little millet (*Panicum sumatrense* Roth. Ex Roem. and Schult.). *Journal of Soils and Crops* 20 (2):211-215.
- Patel G, Dwivedi BS, Dwivedi AK, Thakur R and Singh M. 2018. Long-term effect of nutrient management on soil biochemical properties in a vertisol under soybean-Wheat cropping sequence. *Journal of the Indian Society of Soil Science* 66(2):215-221.
- Pathariya P, Dwivedi BS, Dwivedi AK, Thakur RK, Singh M and Sarvade S. 2022. Potassium Balance under Soybean–wheat Cropping System in a 44-Year-Old Long Term Fertilizer Experiment on a Vertisol, *Communications in Soil Science and Plant Analysis*, 53 (2): 214-226.
- Patidar N, Dwivedi AK, Dwivedi BS, Thakur RK., Bairwa J and Sharma A. 2021. Impact of Long Term Application of Inorganic Fertilizers and Organic Manure on Soil Fertility and Crop Productivity under Soybean-Wheat Cropping System in a Vertisol. *International Journal of Environment and Climate Change*, 11(8): 24-30.
- Peterson GW and Corey RB. 1966. A modified Chang and Jackson procedure for routine fractionation of inorganic soil phosphates. *Soil Sci. Soc. Amer. Proc.* 30:563-565.
- Raghuwanshi C, Khamparia NK, Thakur R, Sahare SK, Sharma V and Dwivedi BS. 2016. Impact of long-term application of inorganic fertilizers and organic manure on yield attribute characters, yield of soybean and soil fertility in a Vertisol. *Eco.Env. & Cons.* 22(2): 745-753.
- Roberts TL and Johnston AE. 2015. Phosphorus use efficiency and management in agriculture. *Resources, conservation and recycling.* 105: 275-281.
- Roy P, Singh YV and Jat LK. 2016. Soil maturity assessment in Indo-Gangetic alluvium of Bihar using soil inorganic fractions-based weathering index: A comparative approach. *J. Indian Soc. Soil Sci.* 64(4): 333-340.
- Sarkar D, Halder A and Mandal D. 2014. Forms of Phosphorus in Relation to Soil Maturity along a Toposequence under Hot, Dry, Subhumid Agro-ecological Subregion of West Bengal. *J. Indian Soc. Soil Sci.*, 62(1): 29-37.
- Singh V, Sharma YK, Singh AK and Bordoloi LJ. 2013. Sulphur distribution in Inceptisol of Northern India and genotypic difference in sulphur uptake of rice. *African Journal of Agricultural Research* 8(10): 855-867.
- Sood RD and Bhardwaj RK. 1992. Effect of liming and phosphorus fertilization on the P fractions in an acid alfisols. *J Indian Soc. Soil Sci.* 40: 299-301.
- Subbiah BV and Asija GL. 1956. A rapid procedure for the estimation of nitrogen in soils. *Curr. Sci.*, 25: 259-260.
- Sudhakaran SV, Patil SR, Kondvilkar NB and Naik RM. 2018. Effect of 32-year long-term integrated nutrient management on soil p fractions and availability of phosphorus under sorghum-wheat cropping sequence in vertisol. *Journal of Pharmacognosy and Phytochemistry*;7(4):3410-3416.
- Suman J, Dwivedi BS, Dwivedi AK and Pandey SK. 2017. Impact of continuous addition of fertilizers and manures on soil nutrient status, uptake, protein and oil content of soybean in a Vertisol. *Research Journal of Agricultural Sci.* 8(1): 159-163.
- Suman J, Dwivedi BS, Dwivedi AK and Pandey SK. 2018. Interaction effect of phosphorus and Sulphur on yield and quality of Soybean in Vertisol. *International Journal of Current Microbial Applied Sciences* 7(3): 152-158.
- Tan KH. 1998 *Principles of soil chemistry*. 3rd ed. New York: Marcel Dekker Inc; 1998.
- Term Application of Chemical Fertilizers and Organics in Rainfed MaizeWheat Cropping System. *J. KrishiVigyan* 3(2): 48-53.
- Tiwari R, Patle T, Khaddar VK and Para P. 2019. Phosphorus Fractions in Different Soil Orders in India and their Relationship with Soil Properties. *International Journal of Current Microbiology and Applied Sciences* 8(5): 1609-1620.
- Walkley A, Black IA. 1934. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil science.* 1934 Jan 1;37(1):29-38.