

ISOLATION AND SUSCEPTIBILITY OF MICROORGANISMS RESPONSIBLE FOR ASYMPTOMATIC ANTENATAL BACTERIURIA

ABSTRACT:

Aim & objective: Determination of prevalence and bacterial profile and antibiotic sensitivity of microorganisms responsible for asymptomatic UTI during pregnancy.

Methodology: The type of study is cross-sectional, Prospective type, was conducted at the PUMHS for Women Nawabshah (SBA) after approval from the Ethical Review Committee and the duration was 2 years from 1st Jan 2021-31st Dec 2022. Samples were collected by non-probability- convenient method from all pregnant women without any clinical complaints attending the Outpatient Department of Gynecology and Obstetrics PUMHS Hospital Nawabshah. The midstream urine collected by clean catch method in sterile containers was processed for DR of urine including physical, chemical, and microscopic examination. After the urine detailed report, the samples were sent for, culture by using the following media, blood, Macconkey's, and CLED agar. Growth-positive samples were proceeding for gram staining, biochemical tests, and antibiotic sensitivity reports according to world standard methods for confirmation.

Results: Out of 477 patients 96 (20%) were positive for ASB, with the mean age of the patient being 29.32 ± 5.74 . The most common microorganisms found in the study were E.coli 67 (69.7%), followed by Klebisella in 4(4%), proteus 9(9%), pseudomonas 8(8.3%), staphylococcus aureus 2(2%), staphylococcus saprophyticus in 4(4%), streptococcus pyogen 2(2%).

There was 100% sensitivity found for all isolated microorganisms following culture examination to following antibiotics: Ciprofloxacin, Cefuroxime, Imepenem, Ceftriaxone Amikacin, Meropenem, and Piperacillin-tazobactam; however, 52% and 45% sensitivity of E.coli and 100% sensitivity of streptococci were present for amoxicillin-clavulanic acid and ampicillin respectively.

Conclusion: A rising number of silent antenatal bacteriuria was observed in younger age women with severe maternal and fetal complications in advanced stages of pregnancy and thereafter. It is therefore suggested to routinely perform urine cultures for every single suspected pregnant lady during advanced antenatal visits to diagnose and properly treat the asymptomatic bacteriuria in terms of avoiding antibiotic resistance and materno-fetal complications in the near future.

Keywords: (1) ASB (2) Antibiotic susceptibility (3) Asymptomatic bacteriuria (4) Urinary tract infection (5) Antenatal (6) Isolation

INTRODUCTION:

Prokaryotic urinary tract infections are a strong inflictor for approximately more than 150 million population per year, among them a huge number of sufferers are females during pregnancy which could be symptomatic or asymptomatic.¹ Anatomical and Physiological alteration during pregnancy enhances the potentiality for easy bacterial contamination, colonization, and proliferation of microorganisms in the urinary tract.^{2,3} Overall global literature-based evidence suggests wide variability in the prevalence of UTI from 2 -47% during pregnancy.^{4,5,6,7} If asymptomatic bacteriuria during pregnancy is left untreated the rate of development of urinary tract infection is 25%.⁸ An Ethiopian study reveals that the rate of UTI following asymptomatic bacteriuria is 70% during pregnancy.⁹ The deleterious maternal–fetal outcomes following ASB are pyelonephritis in the mother, and premature birth, low birth weight in newborns.³ Unfortunately in underdeveloped countries, there are no well-established recommended guidelines for routine antenatal screening for ASB.⁹ According to the American College of Obs: and Gynea, the screening is recommended for Asymptomatic Bacteriuria in all pregnant ladies.¹⁰ Earlier screenings and quick and particular treatment of Asymptomatic infection during pregnancy can prevent further associated disorders of urinary tract infection in pregnant women.^{11,12} The urine culture at antenatal visits is of gold standard for the diagnosis of asymptomatic bacteriuria.^{6,13} In underdeveloped states, the culture of the urine is not practiced in routine antenatal visits in clinics, due to the time factor and high cost involved (48 hrs are required usually).¹⁴ The antimicrobial drug resistance development by over and non-rational application of antibiotics during pregnancy may lead to several morbid risks to mother and child.⁵ There is a strong need to improve the diagnosis with accurate antibiotic prescription for materno-fetal well-being and it will have a positive impact on family and state health care costs.⁵ There is a wide variation in the incidence of microbial agents and antimicrobial resistance patterns globally.⁶ The E.coli or Escherichia coli during pregnancy is the most common isolate responsible for Asymptomatic Bacteriuria,^{3,7,9,15} followed by Cons¹⁵ group B Streptococci⁷, staph Aureus⁶ staph aureus³ CoNS⁹. The purpose of our study was to record the prevalence of ASB and the isolation of microorganisms responsible for asymptomatic UTI in pregnancy and the drug sensitivity patterns of those isolated microorganisms in urine samples of women attending the antenatal outpatient department at People’s Medical Hospital and Department of Pathology, People University of Medical and Health Sciences Nawabshah, Sind, Pakistan.

RATIONALE: Asymptomatic UTI during pregnancy is a silent disease and not routinely diagnosed by antenatal care providers. There is a need to be properly diagnosed based on urine culture reports from microbiological laboratories. These studies are performed in different parts of many countries, with prevalence determination to isolation of bacteria, and their antibiotic sensitivity, but the results are quite variable at every place.

MATERIAL AND METHODS: The type of study is Cross Sectional-Prospective type, by non-probability (convenient) sample technique, and was conducted at the Pathology and Outpatient Department of Obs:&Gynea: PUMHS for Women Nawabshah (SBA)). The duration of the study was January 2021-December 2022.

SAMPLE COLLECTION CRITERIA: The women attending OPD for routine antenatal visits with no any urinary tract complaints were included, the pregnant lady with symptoms of Urinary Tract Infection and currently taking antibiotic treatment were excluded.

Following an informed written consent form taken from pregnant lady a sterile container was used to take the sample of urine quantity of 10-15 ml by clean–catch method. The sample was

sent to the Pathology department for Urine Detailed Report, Culture for growth, and antibiotic sensitivity of microorganisms under strict laboratory standards by using multiple media like blood agar, an Enriched & differential media, Cystine Lactose Electrolyte Deficient agar cystine lactose electrolyte deficient which is specific for micro-organisms in urine, and Mac Conkey's agar a differential media based on lactose fermentation. The samples with positive growth cultures were forwarded for Grams staining, Bio-chemical tests of samples like Triple sugar iron/TSI test, Christensen urea agar, Simon Citrate agar test, string test, catalase test, coagulase test, and for evaluation of antibiotic sensitivity by using specific Mueller Hinton test under world standard methods. Blood agar plate: There is mammalian blood (normally horse or sheep) in blood agar plates (BAPs) generally at a concentration of 5% to 10%. Blood agar plate (BAPs). Cystine lactose electrolyte deficient (CLED) agar, Triple Sugar Iron (TSI) medium has a pH-sensitive dye the phenol red, Simmons' citrate agar is used to differentiate the gram-negative bacteria on the basis of citrate utilization, Christensen's urea medium is used for the growth and detection of bacteria produces urease, the Mueller Hinton agar media, The Catalase test: which is utilized to ascertain the catalase-positive bacteria, Coagulase test is positive when there is clot formation following incubation of bacteria with plasma. The string test is positive when Klebsiella colonies form a pink-colored string after being touched by a wire-loop and then taken back.

DATA ANALYSIS: The data collected on Performaw was designed especially and analyzed by using the 22 version of Statistical Package for the Social Sciences. The percentages of bacteria isolated were recorded and the antibiotic sensitivity was calculated. The results were presented in tables.

RESULTS

The total of 477 patients, the Mean age was 29.32 ± 5.74 years, and the recorded prevalence of ASB was 96(20%) Table no: 01. The samples of urine revealed mostly no growth 381(80%). The organisms isolated from cultures were E.coli 67 (69.7%), followed by Klebsella in 4(4%), proteus 9(9%), pseudomonas 8(8.3%), staph:Aureus 2(2%), staph:Saprophyticus in 4(4%), and the strept:pyogen 2(2%) were the least one Table no: 02.

There is 100% sensitivity to ciprofloxacin, Cefuroxime, Imepenem, Amikacin, Ceftriaxone, Meropenem, and Piperacillin-tazobactam; while E.coli is 52% sensitive to Amoxicillin, Clavulanic acid and 45% sensitivity to Ampicillin while streptococci were 100% sensitive to both antibiotics mentioned above. Table no: 03

Table no: 01 Total number of positive cases

Total number of cases		
No of patient	Positive case	Negative case
477	96(20%)	381(80%)

Table no 02: The number of microorganism positive following the culture of urine.

DISTRIBUTION OF CULTURE POSITIVE ORGANISMS							
Organisms isolated	E.coli	klebseilla	proteus	pseudomonas	staphylococcus aureus	staphylococcus saprophyticus	streptococci
Number Percentage	67(69.7%),	4(4%),	9(9%),	8(8.3%),	2(2%),	4(4%),	2(2%).

TABLE: 03 Antibiotic susceptibility of positive microorganism

antibiotics	E.coli	Klebsella	S. Saprophyticus	S.aureus	Proteus	Pseudomonas	Sterptococci
AMC	35 52%	1 25%	0	0	2 22%	0	2 100%
AMP	30 45%	0%	1 25%	0	0	0	2 100%
CXM	67 100%	4 100%	4 100%	2 100%	9 100%	4 50%	2 100%
AMK	67 100%	4 100%	4 100%	2 100%	9 100%	8 100%	2 100%
CTR	67 100%	4 100%	4 100%	2 100%	9 100%	8 100%	2 100%
CIP	67 100%	4 100%	4 100%	2 100%	9 100%	8 100%	2 100%

TZP	67 100%	4 100%	4 100%	2 100%	9 100%	8 100%	2 100%
MERO	67 100%	4 100%	4 100%	2 100%	9 100%	8 100%	2 100%
IMP	67 100%	4 100%	4 100%	2 100%	9 100%	8 100%	2 100%

DISCUSSION;

The prevalence of asymptomatic bacteriuria varies between 3.75 to 45.3% around the different parts of the globe from developed to underdeveloped countries.^{2,3,4} Our study had a 20% prevalence of asymptomatic bacteriuria, while a study conducted in year 2009 in Lahore Pakistan showed 7.3% of women developing asymptomatic bacteriuria⁶. Asymptomatic bacteriuria during pregnancy remains untreated and is related to increasing morbidity to developing acute pyelonephritis in the mother, and in neonates with increasing chances of lower gestational age, lower birth weight, and increased mortality rate in neonates. Screening of pregnant women visiting antenatal clinics including urine cultures routinely recommended to rule out asymptomatic urinary tract infection and treated with proper antibiotic prescription following culture-positive results. Julius and Celen have found an incidence of ASB to be 3.5% to 8.5%.^{3,16} while another study report from Nigeria showed a higher 86.% prevalence of asymptomatic bacteriuria¹⁷. The prevalence varies from different parts of the world, from the UK 3.75%² and 35.5 from Ghana¹⁸, 4-7% from Canada¹⁹, 4.9% from Australia,² 10% from Egypt,²⁰ 12% from Bangladesh²¹, 15.6% from Ethiopia²², 17.9% from UAE⁷, 16.7% from India¹⁵, few Indian studies showed such a huge variability in prevalence as 8%²³ to 16-17%²⁴ to 25.3%²⁵. Such a wide variation in results from different parts of the world and even from a single country may be due to different sample sizes, geographical differences, community social habits, availability of health-related facilities, and their proper consumption. A large number of studies from various areas of the globe have documented that ASB is caused by Staphylococcus hemolytic and aerobic bacilli of gram-negative group and earlier and particular antibiotic treatment was suggested. The commonest microorganism isolated in our study was E coli then Proteus, pseudomonas and Klebsiella etc. E.coli is the commonest organism causing asymptomatic UTI during pregnancy,^{7,15} but the congruent results were found at Ghana¹⁷ where gram-positive S.aureus 64% and CoNS 70% are responsible for causing asymptomatic bacteriuria in pregnancy. Proteus is the 2nd common organism in our study but Group B streptococci at UAE⁷ and CoNs are found at India¹⁵ respectively. All the common microorganisms cultured were 100% sensitive to

most antibiotics. The most commonly isolated organism E. coli was 52% sensitive to Amoxicillin-clavulonic acid and 45% to Ampicillin but was sensitive to all the other antibiotics. As Penicillin and cephalosporin are safe during pregnancy, all our patients respond well to treatment. Higher resistance was detected against Ampicillin, Ceftriaxone, Cotrimoxazole, Erythromycin, and Nitrofurantoin was detected by Ezekiel.²⁶ The higher resistance may be due to irrational use and availability of antibiotics. The study by Jain showed that E. coli was the more common causative organism isolated and was 37.6% then Enterococcus species in 21.1% on the 2nd commonest organism. Celen reported in his study that the commonest organism was E. coli 76.6% followed by 14.6% Klebsiella, but the sensitivity and specificity for fosfomycin and nitrofurantoin were greater than 99.2% and 88%. Tadesse in his study found Staphylococcus at 32.6%, most common followed by Escherichia coli at 26.1%, Staphylococcus Aureus at 13% with increased sensitivity to norfloxacin 64.7%, and 17.6% at for ampicillin²⁷.

It is therefore proposed that urine culture be practiced on antenatal visits of pregnant women at the outpatient Department before prescribing antibiotics to avoid bacterial resistance development to drugs safe in pregnancy. The results of this study suggest that after sending a urine culture of each symptomatic pregnant lady with positive urine D/R, 97% safe practices be achieved and may also decrease blind treatment burden by avoiding antibacterial drugs. In short, there is a need for more research and practices to be executed for pregnant ladies, with positive reports of urine cultures with few risk factors like previous history of Urinary Tract Infections.

CONCLUSION

The younger age pregnant ladies were increasingly affected by ASB with advancing pregnancy and parity. Therefore urine cultures should be routinely practiced during antenatal visits to diagnose the ASB and treated with prompt and proper antibiotics in order to avoid the increasing resistance of those antibiotics that can be used during pregnancy and to avoid materno-fetal complications.

On the other hand, improper use of antibiotics may cause harm to mother and child. The main reason for increasing resistance to simple antibiotics that can be used during the antenatal period to treat ASB was excessive and improper use. There is also a risk to the fetus with dispensable exposure to antibiotics. It was observed recently that exposure to antibiotics to the mother during pregnancy may lead to alterations in the differential methylation at regulatory regions of imprinted genes.⁵ By improving the diagnosis and properly prescribed antibiotics, fetal development will be smooth. All these steps will affect the future healthcare expenses. When we avoid the huge list of tests and specify them for accurate diagnosis, costs will be reduced and saved.⁵

Ethical Approval

The study was conducted at the PUMHS for Women Nawabshah (SBA) after approval from the Ethical Review Committee and the duration was 2 years from 1st Jan 2021-31st Dec 2022.

Consent

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

References:

1. Sial SA, Detho AB, Memon FR, Parveen K, Memon S, Bukhari S, Ahmer A. Risk Factors during Pregnancy for Asymptomatic Bacteriuria. *Journal of Pharmaceutical Research International*. 2021 May 8;33(28B):21-6.
2. Sheppard M, Ibiebele I, Nippita T, Morris J. Asymptomatic bacteriuria in pregnancy. *Australian and New Zealand Journal of Obstetrics and Gynaecology*. 2023 May 8.
3. Nteziyaremye J, Iramiot SJ, Nekaka R, Musaba MW, Wandabwa J, Kisegerwa E, Kiondo P. Asymptomatic bacteriuria among pregnant women attending antenatal care at Mbale Hospital, Eastern Uganda. *PloS one*. 2020 Mar 19;15(3):e0230523.
4. Wingert A, Pillay J, Sebastianski M, Gates M, Featherstone R, Shave K, Vandermeer B, Hartling L. Asymptomatic bacteriuria in pregnancy: systematic reviews of screening and treatment effectiveness and patient preferences. *BMJ open*. 2019 Mar 1;9(3):e021347.
5. Werter DE, Kazemier BM, van Leeuwen E, de Rotte MC, Kuil SD, Pajkrt E, Schneeberger C. Diagnostic work-up of urinary tract infections in pregnancy: study protocol of a prospective cohort study. *BMJ open*. 2022 Sep 1;12(9):e063813.
6. Hafeez S, Hafeez Z, Hafeez S, Riaz S, Yousuf A. Frequency of Asymptomatic Bacteriuria and its Causative Organisms among Pregnant Women Visiting for Antenatal Checkup in Outpatient Department of Sims/Services Hospital. *Age (Years)*.;26:4-08.
7. Dube R, Al-Zuheiri ST, Syed M, Harilal L, Zuhaira DA, Kar SS. Prevalence, Clinico-Bacteriological Profile, and Antibiotic Resistance of Symptomatic Urinary Tract Infections in Pregnant Women. *Antibiotics*. 2022 Dec 25;12(1):33.
8. Taha LH, Almahmood AA, Farhan RK. Distribution of asymptomatic bacteriuria among women using Intrauterine Devices (IUDs) according sociodemographic characteristics factors in Takrit, Iraq. *Iraq Medical Journal*. 2021 Dec 26;5(4).
9. Edae, M., Teklemariam, Z., Weldegebreal, F. and Abate, D., 2020. Asymptomatic bacteriuria among pregnant women attending antenatal care at Hiwot Fana specialized university hospital, Harar, eastern Ethiopia: magnitude, associated factors, and antimicrobial susceptibility pattern. *International Journal of Microbiology*, 2020.

10. C. B. Ned, M. Petitti, T. G. DeWitt, and H. Russell, "Screening for asymptomatic bacteriuria in adults: U.S. Preventive services task force reaffirmation recommendation statement," *Annals of Internal Medicine*, vol. 149, no. 4, 2008
11. Mwei MK, Mchome B, John B, Maro E. Asymptomatic bacteriuria among pregnant women attending antenatal clinic at Kilimanjaro Christian Medical Centre in Northern Tanzania. *Clin Pract*. 2018;15(6):917-22.
12. Kafi AA, Khatun M, Shoshi BN. The Association of Asymptomatic Urinary Tract Infection (UTI) with Diabetes Mellitus (DM) in Pregnant Women.
13. Sujatha R, Nawani M. Prevalence of asymptomatic bacteriuria and its antibacterial susceptibility pattern among pregnant women attending the antenatal clinic at Kanpur, India. *Journal of clinical and diagnostic research: JCDR*. 2014 Apr;8(4):DC01.
14. Ali R, Afzal U, Kausar S. Asymptomatic bacteriuria among pregnant women. *Annals of Punjab Medical College (APMC)*. 2011 Dec 12;5(2):155-8.
15. Sonkar N, Banerjee M, Gupta S, Ahmad A. Asymptomatic bacteriuria among pregnant women attending tertiary care hospital in Lucknow, India. *Dubai Medical Journal*. 2021 Mar 11;4(1):18-25
16. Celen S, Oruç AS, Karayalçın R, Saygan S, Unlü S, Polat B, Danişman N. Asymptomatic bacteriuria and antibacterial susceptibility patterns in an obstetric population. *ISRN Obstet Gynecol*. 2011; 2011:721872. Doi: 10.5402/2011/721872. Epub 2011 Jan 24.
17. Akerele J, Abhulimen P, Okonofua F. Prevalence of asymptomatic bacteriuria among pregnant women in Benin City, Nigeria. *J Obstet Gynaecol*. 2001 Mar;21(2):141-4. doi: 10.1080/01443610020026038. PMID: 12521882.
18. Karikari, A.B., Saba, C.K.S. & Yamik, D.Y. Assessment of asymptomatic bacteriuria and sterile pyuria among antenatal attendants in hospitals in northern Ghana. *BMC Pregnancy Childbirth* **20**, 239 (2020). <https://doi.org/10.1186/s12884-020-02936-6>
19. Nicolle LE. Screening for asymptomatic bacteriuria in pregnancy. Ottawa Health, Canada: Canadian guide on preventive health care; 1994. pp. 100–106
20. Elzayat MA, Barnett-Vanes A, Dabour MF, Cheng F. Prevalence of undiagnosed asymptomatic bacteriuria and associated risk factors during pregnancy: a cross-sectional study at two tertiary centers in Cairo, Egypt. *BMJ open*. 2017 Mar 1;7(3):e013198.
21. Ullah MA, Barman A, Siddique MA, Haque AK. Prevalence of asymptomatic bacteriuria and its consequences in pregnancy in a rural community of Bangladesh. *Bangladesh Medical Research Council Bulletin*. 2007;33(2):60-4.
22. Ali, I.E., Gebrecherkos, T., Gizachew, M. and Menberu, M.A., 2018. Asymptomatic bacteriuria and antimicrobial susceptibility pattern of the isolates among pregnant women attending Dessie referral hospital, Northeast Ethiopia: A hospital-based cross-sectional study. *Turkish Journal of urology*, *44*(3), p.251.

23. Jayalakshmi J, Jayaram VS. Evaluation of various screening tests to detect asymptomatic bacteriuria in pregnant women. *Indian J Pathol Microbiol.* 2008 Jul-Sep;51(3):379-81. doi: 10.4103/0377-4929.42516. PMID: 18723963.
24. Jain V, Das V, Agarwal A, Pandey A. Asymptomatic bacteriuria & obstetric outcome following treatment in early versus late pregnancy in north Indian women. *Indian J Med Res.* 2013 Apr;137(4):753-8. PMID: 23703344; PMCID: PMC3724257.
25. Patnaik M, Panigrahi K, Das B, Pathi B, Poddar N, Lenka PR, Pattnaik D. Prevalence, risk factors and causative organisms of asymptomatic bacteriuria in pregnancy. *Int J Adv Med.* 2017 Sep;4(5):1348-54.
26. Vicar, E.K., Acquah, S.E., Wallana, W., Kuugbee, E.D., Osbutey, E.K., Aidoo, A., Acheampong, E. and Mensah, G.I., 2023. Urinary Tract Infection and Associated Factors among Pregnant Women Receiving Antenatal Care at a Primary Health Care Facility in the Northern Region of Ghana. *International Journal of Microbiology*, 2023.
27. Tadesse E, Teshome M, Merid Y, Kibret B, Shimelis T. Asymptomatic urinary tract infection among pregnant women attending the antenatal clinic of Hawassa Referral Hospital, Southern Ethiopia. *BMC Res Notes.* 2014 Mar 17; 7:155. Doi: 10.1186/1756-0500-7-155.