

Original Research Article

MODELLING THE LINEAR AND NONLINEAR EFFECTS OF MALARIA RISK FACTORS ON CHILD MORTALITY

Abstract

Generalized additive model was used to analyse data from Nigeria standard demographic and health survey (NDHS) 2018. The sample consists of 10609 children aged 6-59 months who were tested for malaria parasitemia through the rapid diagnostic test (RDT). Child mortality data was obtained by calculating the difference between the number of children ever born and the proportion of children alive during the survey. The analysis was carried out in R version 4.1.1 via mgcv package. The results obtained indicated linear and nonlinear effects of malaria risk factors on child mortality. The findings also revealed mosquito bed net usage, wealth index, maternal education, type of place of residence and malaria test outcome as significant predictors of child malaria mortality.

Key words: Malaria risk factors, Child mortality, Mortality rate, Nonlinear effects.

Introduction

Researchers made efforts to elucidate the link between malaria transmission and child mortality Snow and Marsh (1995), Trape and Rogier (1996), Snow *et al.*, (1999a), Snow *et al.*, (1999b), Smith *et al.*, (2001), Lindblade *et al.*, (2004), Rowe and Steketee (2007), Lim *et al.*, (2011). Ahmad *et al.*, (2000) observed that it is very difficult to determine whether the decline in the infant mortality rate is attributable to the success of a particular health intervention program, change in reproduction patterns, socio-economic development, or the combination of two or more of these factors. This is due to the scarcity of reliable data on the causes of infant death in developing countries. However, they opined that provided major health threats such as malaria and HIV/AIDS do not increase over the next few years in Africa particularly, child mortality might be reduced to half by 2030.

Regardless of age, everyone is liable to malaria infection. However, under five children and pregnant women account for highest burden of malaria infection. It is also observed that younger children under five years stand to be more vulnerable due to under developed immunity against mild forms of the disease Kazembe *et al.*, (2006), Kazembe *et al.*, (2008), Nigeria malaria facts sheet (2011), Herrera *et al.*, (2020). Child mortality in high malaria endemic regions is high Kalipeni (1993), Smith *et al.*, (2001)

It was suggested that 'all cause child mortality' (ACCM) can be proportionately reduced if malaria mortality decreases through scaling up of malaria intervention programs Lindblade *et al.*, (2004), Rowe and Steketee (2007). A strong positive association between infant mortality rate and entomological inoculation rate was revealed in Africa Smith *et al.*, (2001). On the other hand, Rowe *et al.*, (2006) cautioned the use of malaria burden research results to estimate number of deaths that would be prevented if malaria transmission were reduced or eradicated. However, they further explained that such estimate would lead to under estimation due to the fact that it would not include the large burden of indirect malaria mortality. They found a historical decline in all-cause infant mortality rates due to drastic reduction of malaria transmission.

Significant difference in child mortality rates between areas with insecticide treated nets interventions and the areas without interventions in Western Kenya was not observed Lindblade *et al.*, (2004). In a similar effort to find out clear and reliable association that may exist between malaria and child mortality rate, Gemperli (2003) linked Demographic and Health Surveys (DHS) and Mapping Malaria Risk in Africa (MARA) databases but no clear relationship was observed. Considering the research results and findings cited it is extremely difficult to conclude in generic terms the direct link between malaria transmission and child mortality. Also most of the researches were aimed at assessing and evaluation the effectiveness of the intervention programs rather than the relationship between malaria transmission and child mortality Rumisha *et al.*, (2014).

Several factors are attributable to child (under five) mortality in various countries. The factors include; sanitary conditions, access to health facilities, breast feeding, maternal, environmental, wealth index, age at first birth of the mother, place of residence, education, climate and so on. Saroj *et al.*, (2019) identified socio-economic, demographic, educational level, mother's age, size of the child, play a very important role in the under-five child mortality using Cox regression model. Poisson regression model was used to identify number of health facilities as the most significant variable in under-five mortality estimation Sudarno (2018). In another regression model, sanitation facilities were found to be a determinant factor in child mortality Cavalcanti *et al.*, (2019). Adetoro and Amoo (2014) identified maternal education as indicator for child mortality rate. Bourne (2012) observed log poverty and gross domestic product (GDP) per capita account for ninety percent (90%) of the explanation of changes in child mortality rate. Considering the foregoing, one can observe a direct interlink between malaria and child mortality risk factors.

Dansu and Asiribo (2007) attempted to investigate spatial association between malaria pandemic and mortality. Contrary to the title of their work, no data analysis or results were provided for valid association between malaria risk factors and mortality. Moreover, the data collected was historical, small and aggregated without a specified sample.

Snow *et al.*, (2004) made a comprehensive research on paediatric mortality in Africa with emphasis on malaria as mortality risk. The data and information were drawn from the burden of malaria in Africa (BOMA), unpublished ministry of health and conference materials and other archives in some African countries Snow *et al.*, (2004). Publications from the United Kingdom were used in addition to direct contact with the authors of published materials in order to clarify issues that were not clear in their original reports. Mortality reports were based on the definition of demographic surveillance system (DSS). The DSS data was used to provide estimate of annual mortality risk from birth to five years of life. The use of DSS data was borne out of the fact that the DSS restrict areas of definable malaria risk which allows the exploration of malaria transmission dynamics Snow *et al.*, (2004). Other technical and specialist definitions and medical parameters were extensively explained and the sources of information were provided.

The statistical methods used were descriptive using medians and interquartile ranges. Weighted least square regression was applied to model contiguous relationship between mortality and malaria prevalence. The model considered square of prevalence as a variable that allowed for the possibility of saturating or decreasing mortality risk at the highest prevalence level Snow *et al.*, (2004). Also, the stratified study site locations were considered as categorical variables. No regional influence was taken into consideration. They also revealed that the removal of infection risk might reduce the mortality by more than two folds.

In the second part of their work, causal relationship was considered. Because mortality rate might be due to interactions of malaria and other factors such as poverty that were capable of determining both malaria and mortality. To examine this effect, Snow *et al.*, (2004) attempted to investigate the malaria program effectiveness. In other words, how reduction in malaria risk might affect under-five mortality.

As in the case of logistic regression, ordinary regression model assumes linearity between response and explanatory variables. The model does not account for possible nonlinear relationship between dependent variable and independent variables.

Objective of the study

The following are the objectives of the study;

- i. To identify the malaria risk factors and related child mortality rate,
- ii. To assess linear and nonlinear effects of malaria risk factors on child mortality,
- iii. To assess the effects of categorical variables on malaria mortality risk,
- iv. To assess the effects of continuous covariates on malaria mortality risk.

Methodology

Data Source

The data used for the analysis in this study was obtained from the 2018 Nigeria standard demographic and health survey.

The Sample

A sample of 10609 children aged 6-59 months was collected, tested for malaria parasitemia and used for data analysis.

Variables of Malaria Risk Factors

Eleven (11) variables were selected for analysis. These are;

Age in Months, cluster altitude, Child weight, sleeping inside mosquito net by children, wealth index, mother's educational level, Mortality rate, place of residence, region, number of children and results of malaria test.

Child Mortality Data

The multiple indicator survey serves as reliable alternative for vital statistics data in Nigeria. The data was obtained from the information on the number of children ever born and the proportion of children alive during the survey. The difference gives the number of the dead children. That is, $\text{Number of dead children} = \text{Number of children ever born} - \text{Number of children alive during the survey}$. The difference is then aggregated into the children personal record data in the survey for the analysis in the model. The mortality rate was calculated using equation (9).

The Structured Additive Regression Model

The variable of interest (the dependent variable) y_i is the survival status of a child for malaria infection. Let y_i be the mortality status of a child i . Let the status "child is alive" (survived malaria) be 0 and 1 otherwise. That is;

$$y_i = \begin{cases} 0 & \text{if the child is alive (with probability } \pi) \\ 1 & \text{otherwise (with probability } 1-\pi) \end{cases} \quad (1)$$

y_i is exponential family distribution. That is, $y_i \in \text{EF}(\mu_i, \phi)$ with mean μ_i and scale parameter ϕ . That is, binary response data follows a Bernoulli distribution.

$$y_i \in \text{Bernoulli}(\pi_i) \quad (2)$$

Where π_i is the probability that the child is alive. And $\pi_i = \phi_i(\eta_i)$ where $\phi(\cdot)$ is the cumulative distribution function for the standard normal distribution and η_i is the predictor modelling the dependence of y on the covariates Fahrmeiret al., (2013). Now let

$$W_i' = (w_{i1}, w_{i2}, \dots, w_{ip})' \quad (3)$$

be a matrix of explanatory variables. Assuming linear predictor, we obtain

$$\eta_i = W_i' \beta \quad (4)$$

Since y_i has binary response, we use a logit regression model. Therefore the ordinary logistic regression with log link function is written as

$$A: \eta_i = \log\left(\frac{\pi_i}{1-\pi_i}\right) = \beta_0 + \beta_i' W_i' \quad (5)$$

where β_0 is the intercept, β_i is a vector of regression coefficients and W_i' is a categorical covariate vector. In generalized structured additive regression model, we replace the linear predictor by additive predictor Hastie and Tibshirani (1990), Fahrmeiret al., (2013). Therefore,

$$\log it(\pi) = \log\left(\frac{\pi}{1-\pi}\right) = f_0 + \sum_{i=1}^q f_i(x_{ik}) \quad (6)$$

The model can be extended to include parametric and non-parametric terms. Therefore we can write (5) and (6) as

$$B: \eta_i = \beta_0 + \beta_i' W_i' + \sum_{k=1}^q f_k(x_{ik}) \quad (7)$$

Where $\sum_{k=1}^q f_k(x_{ik})$ are nonlinear smooth effects of the continuous covariates. β_0 is the intercept, β_i is the parameter corresponding to the categorical fixed variables $W_i' = (w_{i1}, w_{i2}, \dots, w_{ip})'$.

The model can be further extended to include spatially unstructured random effect ϕ_i , spatially structured random effects u_i Okangoet al (2015) and mortality rate ψ_i . Therefore we obtain

$$C: \eta_i = \beta_0 + \beta_i W_i' + \sum_{k=1}^q f_k(x_{ik}) + \phi_i + \nu_i + \psi_i \quad (8)$$

Where ψ_i represents mortality rates due to malaria at time t . Mortality rate can be obtained through the Standard Mortality Rate (SMR) given by

$$\psi_i = SMR = \frac{\text{Observed child deaths due to malaria}}{\text{Expected number of child deaths}} \times 100\% \quad (9)$$

Variable Description

Dependent variable: The response variable for this study was the mortality status of a child aged 6-59 months. The response was dichotomous as dead (1) or alive (0).

Independent Variables

The explanatory variables used for this study were classified into two groups: Categorical variables such as; sleeping inside insecticide treated nets, wealth index, mother's educational level, type of place of residence, region and result of malaria test. Region was categorized into six geo-political zones as North East, North West, North Central, South South, South East and South West. Type of place of residence is categorical (Rural/Urban), Result of malaria test (positive or negative), sleeping inside bed net (all, none, some and no net), wealth index (poorest, poorer, middle, richer and richest), mother's educational level (no education, primary, secondary and higher). Continuous variables include; Cluster altitude (meters), number of children, age (months), child weight (kg) and mortality rate (%). (Ogunsakin and Chen (2020)).

Results

Structured additive regression analysis was carried out using R version 4.1.1 to determine the effects of each of the malaria risk factors on child mortality. Estimation of parameters was done through the restricted maximum likelihood (REML) in mgcv package.

Table 1 Output summary of the parametric component

| Predictor | Estimate | Std. Error | p-value |
|-------------------------------------|----------|------------|-----------------------|
| Intercept | -5.6414 | 0.4101 | 2.0×10^{-16} |
| Sleeping under bed net All children | -0.5256 | 0.1100 | 1.77×10^{-6} |

| | | | | |
|------------------------------------|----------------------|----------|--------|-------------------------|
| | Some Children | -0.6383 | 0.1527 | 2.92×10^{-5} |
| | No net | -0.5530 | 0.1194 | 3.62×10^{-6} |
| | None of the children | 1.00 | | |
| Wealth Index | Poorer | -0.3173 | 0.1023 | 0.0019 |
| | Middle | -0.4454 | 0.1168 | 0.0001 |
| | Richer | -0.8266 | 0.1538 | 7.60×10^{-8} |
| | Richest | -1.1335 | 0.2459 | 4.02×10^{-6} |
| | Poorest | 1.0 | | |
| Mother's level of Education | | | | |
| | Primary | 0.1218 | 0.1202 | 0.3109 |
| | Secondary | 0.1529 | 0.1225 | 0.2121 |
| | Higher | 0.5512 | 0.2636 | 0.0365 |
| | Don't know | 0.1585 | 0.1515 | 0.2955 |
| No Education | 1.0 | | | |
| Malaria test results | Positive | 4.9465 | 0.3443 | $< 2.0 \times 10^{-16}$ |
| Negative | 1.0 | | | |
| Region | North East | - 0.0361 | 0.1355 | 0.7899 |
| | North West | 0.1050 | 0.1321 | 0.4268 |
| | South East | - 0.2277 | 0.1781 | 0.2010 |
| | South South | - 0.2297 | 0.1930 | 0.2340 |
| | South West | 0.3607 | 0.1477 | 0.0146 |
| | North Central | 1.0 | | |
| Place of Residence | Rural | 0.3682 | 0.1125 | 0.0011 |
| | Urban | 1.0 | | |

Table 1 shows the results for the parametric component of the model. It gives the parametric coefficients as in the usual logistic regression model (linear model). The intercept gives the value of the child mortality when all numerical predictors equal to zero and the factor (categorical) variables were at their reference level. For instance, if the malaria test is negative, it means the average child mortality for malaria negative child is -5.64 . That is, the probability of child mortality for malaria negative child is 0.003 . The effects of sleeping under mosquito bed net, wealth index of the households, results of malaria rapid diagnostic test and place of residence were highly significant. Effect of mother's educational at higher level and the South West region were also significant.

Table 2 Output summary of the nonparametric component

| Predictor | edfRef.df | Chi. Sq. | p-value | |
|--------------------|-----------|----------|---------|--------|
| Cluster altitude | 4.2693 | 4.9896 | 17.653 | 0.0032 |
| Child weight | 1.0031 | 1.0060 | 7.422 | 0.0065 |
| Mortality rate | 1.0002 | 1.0003 | 4.868 | 0.0274 |
| Age | 1.8957 | 2.3328 | 2.708 | 0.2894 |
| Number of children | 1.0001 | 1.0002 | 1.563 | 0.2113 |

The results in table 2 shows that the cluster altitude has a statistically significant nonlinear effect (edf = 4.2693, $p = 0.0032$). the effects of child weight and mortality rate were linearly significant with edf 1.0 each, p-values 0.0065 and 0.0274 respectively. The effect of age was nonsignificant and nonlinear (edf =1.8957, $p = 0.2894$). Number of children has nonsignificant linear effect (edf = 1.0001, $p = 0.2113$).

Figure 1 shows the graph of nonlinear effect of the continuous covariates included in the modified model. The x-axis contains the values of cluster altitude (in meters), child weight (in kilogram), mortality rate (percent), age (in months) and number of children (in units) for respective graphs. The y-axis contains the partial effects of the covariates on the response (child mortality). The curvy shapes on the graphs for cluster altitude and age resulted from smoothing splines that models the nonlinearity in the data. The dotted lines around the main (bold) line signify the 95% confidence region.




Figure 1 smooth effect of the continuous covariates

Figure 1A indicates a nonlinear effect of cluster altitude on malaria related child mortality. The effect of malaria mortality was lowest at 500 meters and highest at 1000 meters above sea level. Figure 1B shows inverse linear relationship between child weight and its effects on malaria related mortality. The effect on child malaria mortality sharply decreases with increase in child weight. Figure 1C revealed a linear increase in the effect of mortality rate on child malaria mortality with increase in mortality rate with very wide confidence limits. Nonlinear partial effect of child's age on malaria mortality was observed in figure 1D. The highest effect of malaria related mortality was indicated in children of 35 months old and lowest in children aged less than 10 months. The effect of number of children was linear and decrease with increase in the number of children.

Discussion

The model was used to analyse data from the Nigeria national demographic and health survey (NDHS) 2018. The results of the analysis showed that the effects of sleeping under mosquito bed net, wealth index and results of malaria rapid diagnostic tests were highly significant while educational level of the mother was not. This contradicts the work of Ernst *et al.*, (2009), Seyum (2018) that reported significant effect of maternal level of education. A

significant association was observed between the mortality status of children and risk factors such as wealth index, sleeping inside mosquito bed net, regional location of children, place of residence and results of malaria test. The probability of malaria related mortality in children aged 6-59 months was highest in the South West (0.0175) and lowest in the South East (0.012). Use of mosquito bed net for sleeping significantly reduces the probability of malaria infection and consequently reduces the risk of malaria child mortality. Likelihood of child malaria mortality declined in the households where all children sleep inside mosquito bed net than in the households where none of the children sleep inside the bed net. This is in agreement with the work of Selemani *et al.*, (2016), Oguoma *et al.*, (2021) and Awosolu *et al.*, (2021). This however, contradicts the findings of Adigun *et al.*, (2015), Ugwu and Zewotir (2018) that the use of mosquito bed net has nonsignificant effect on malaria positivity in children.

Economic status of the household plays a key role in determining the probability of child malaria mortality. Direct measurement of household of household income seems to be extremely difficult due to a number of reasons. For instance, many respondents tend to conceal their income to interviewers. In addition, most of household income was persistently varied on daily, weekly or seasonal basis. Therefore, household wealth index serves as alternative variable that measure economic status of the households. Wealth index is a composite measure of a household cumulative living standard. This entails the ownership of some easy to measure household assets such as television sets, radio, mobile phone, car, bicycle, material used for building the house, sanitation facilities and so on. The indices were generated using principal component analysis.

Probability of malaria child mortality decreases significantly with increase in household wealth quantile. In other words, children from the poorest household were more likely to die of malaria infection than children from the richer households. This confirms the findings of Ahmed *et al.*, (2000), Bourne (2012) and Saroj *et al.*, (2019). Type of place of residence showed a significant effect on the probability of malaria risk of child mortality. The probability of malaria mortality risk was higher in the households living in the rural areas than those from the urban centres. This supports the research finding of Awosolu *et al.*, (2021).

Continuous covariates were modelled with smoothing splines to take care of possible nonlinear effects. The results showed that the cluster altitude and age of the subjects had nonlinear effect on the response variable. The effect of the cluster altitude was statistically significant while the effect of age was not. The risk of child malaria mortality was highest at 1000 meters above sea level and lowest at 500 above sea level. Similar finding was reported in Bødker *et al.*, (2003) and Siya *et al.*, (2020). However, this contradicts the findings of Chiromboet *et al.*, (2014) and Ugwu and Zewotir (2020). The nonlinear effect of age on child malaria mortality was parabolic indicating maximum value at 35 months. The risk of malaria child mortality was lower for younger children and stepwise increases with age. This supports the findings of Dondorpet *et al.*, (2008) that claimed significant association exists between child's age and mortality. This result was also supported by the research finding of Desai *et al.*, (2014) that reported decline in malaria related mortality in children of less than 36 months old and increase among children aged between three to five years.

The effects of child's weight, mortality rate and number of children in the household were linear. The likelihood of child malaria mortality decreases with increase in child weight. In other words, child weight had inverse negative effect on child mortality. The risk of malaria mortality declines with increase in child's weight. This confirms the findings of Alexandre *et al.*, (2015), Afoakwahet *et al.*, (2018) and Djimdeet *et al.*, (2019). The size of the households has

not shown nonlinear effect on the child malaria mortality. However Guerra *et al.*, (2018) reported significant association between household size and malaria risk. The partial effect of mortality rate on child malaria mortality was linear and directly proportional. This means as the mortality rate increases the child malaria mortality also increases. This finding was contrary to the findings of Spencer *et al.*, (1987) who reported lack of evidence to link mortality rates to malaria specific mortality. However recent researches confirm a strong link between malaria child mortality and mortality rate Streatfield *et al.*, (2014), Coetzer and Adeola (2020). The random effects of regional location and type of place of residence on child malaria mortality were statistically significant. This confirms the work of Chiromboet *et al.*, (2014).

Conclusion

The findings in this study showed that there was both linear and nonlinear relationship between the child malaria mortality and the predictors. Linear relations were indicated by categorical variables such as sleeping under mosquito bed net, wealth index, maternal education, results of malaria RDT test, regional location and place of residence. The continuous predictors that showed linear effects were; child weight, mortality rate and number of children in the household. The findings also revealed nonlinear relationship between child malaria mortality and smooth effects of cluster altitude and age. This study also revealed that the variables; sleeping under mosquito bed net, wealth index, mother's level of education, malaria test results, cluster altitude, child weight, mortality rate and type of place of residence were relevant predictors in modelling malaria risk factors of child mortality.

References

- Adetoro, G. W. and Amoo, E., O. (2014) A Statistical Analysis of Child Mortality: Evidence from Nigeria. *Journal of Demography and Social Statistics*1(1): 10 – 20
- Adigun, A. B., Gajere, E. N., Oresanya, O., Vounatsou, P. (2015) Malaria Risk in Nigeria: Bayesian Geostatistical Modelling of 2010 Malaria Indicator Survey Data. *Malaria Journal*14(156) DOI 10.1186/s12936-015-0683-6
- Afoakwah, C., Deng, X., Onur, I. (2018) Malaria Infection Among Children Under-five: The Use of Large-scale Interventions in Ghana. *BMC Public Health*18(536). <http://doi.org/10.1186/s12889-018-5428-3>
- Ahmad, O. B. Lopez, A. D. Inoue, M. (2000) The Decline in Child Mortality: A Reappraisal. *Bulletin of the World Health Organization* No. 78 1175-1191
- Ahmed, S., Reithinger, R., Kaptoge, S. K., Ngondi, J. M. (2020) Travel Is a Key Risk Factor for Malaria Transmission in Pre-Elimination Settings in Sub-Saharan Africa: A Review of the Literature and Meta-Analysis. *American Journal of Tropical Medicine and Hygiene*103(4) 1380–1387
- Alexandre, M. A. A., Benzecry, S. G., Siqueira, A. M., Vitor-Silva, S., Melo, G. C., Monteiro, W. M., Leite, H. P., Lacerda, M. V. G., Alecrim, M. G. C. (2015) The Association between Nutritional Status and Malaria in Children from a Rural Community in the Amazonian Region: A Longitudinal Study. *PLOS Neglected Tropical Diseases*, 9(4). doi:10.1371/journal.pntd.0003743

- Awosolu, O. B., Yahaya, Z. S., Haziqah, M. T. F., Oke, I. A. S., Fakunle, C. (2021) A cross-sectional study of the prevalence, density, and risk factors associated with malaria transmission in urban communities of Ibadan, South Western Nigeria. *Heliyon*7. <https://doi.org/10.1016/j.heliyon.2021.e05975>
- Bødker, R., Akida, J., Shayo, D., Kisinza, W., Msangeni, H. A., Pedersen, E. M., Lindsay, S. W. (2003) Relationship Between Altitude and Intensity of Malaria Transmission in the Usambara Mountains, Tanzania. *Journal of Medical Entomology* 40(5) 706-717. DOI: 10.1603/0022-2585-40.5.706
- Bourne, P. A. (2012) Under-Five Mortality, Health and Selected Macroeconomic Variables: The Children behind the Digits. *Epidemiology* 2(2) 2161-2165 <http://dx.doi.org/10.4172/2161-1165.1000115>
- Cavalcanti, A., Teixeira, A., Pontes, K. (2019) Regression Model to Evaluate the Impact of Basic Sanitation Services in Households and Schools on Child Mortality in the Municipalities of the State of Alagoas, Brazil. *Sustainability* 11(4150) doi:10.3390/su11154150
- Chirombo, J. Lowe, R. Kazembe, L. (2014) Using Structured Additive Regression Models to Estimate Risk Factors of Malaria: Analysis of 2010 Malawi Malaria Indicator Survey Data *PLoS ONE*9(7) www.plosone.org
- Coetzer, R. H. and Adeola, A. M. (2020) Assessing the Correlation between Malaria Case Mortality Rates and Access to Health Facilities in the Malaria Region of Vhembe District, South Africa. *Journal of Environmental and Public Health*. <https://doi.org/10.1155/2020/8973739>
- Dansu, B. M. and Asiribo, O. E. (2007) Spatial Association Between Malaria Pandemic and Mortality. *Data Science Journal*, 6(18) 145-153
- Desai, M., Buff, A. M., Khagayi, S., Byass, P., Amek, N., Eijk, A. V., Slutsker, L., Valule, J., Odhiambo, F. O., Howard, P. A. P., Lindblade, K. A., Laserson, K. F., Hamel, M. J. (2014) Age-specific Malaria Mortality Rates in the KEMRI/CDC Health and Demographic Surveillance System in Western Kenya 2003-2010. *PLOS ONE*9(9) www.plosone.org
- Djimde, M., Samouda, H., Jacobs, J., Niangaly, H., Tekete, M., Sombie, S. B., Mgina, E. J., Fofana, B., Sagara, I., Doumbo, O. K., Vaillant, M., Djimde, A. A. (2019) Relationship between Weight Status and Anti-malarial Drug Efficacy and Safety in Children in Mali. *Malaria Journal*18(40). <https://doi.org/10.1186/s12936-019-2673-6>
- Dondorp, A. M., Lee, S. J., Faiz, M. A., Mishra, S., Price, R., Tjitra, E., Than, M., Htut, Y., Mohanty, S., Yunus, E. B., Rahman, R., Nosten, F., Anstey, N. M., Day, N. P. J., White, N. J. (2008) Relationship Between Age and the Manifestations of and Mortality Associated with Severe Malaria. *Clinical Infectious Diseases* 47 151-157
- Ernst, K. C., Lindblade, K. A., Koech, D., Sumba, P. O., Kuwuor, D. O., John, C. C., Wilson, M. L. (2009) Environmental, Socio-demographic and Behavioural Determinants of Malaria Risk in the Western Kenyan Highlands: A Case-control Study. *Tropical Medicine and International Health* 14(10): 1258-1265

- Fahrmeir, L., Kneib, T., Lang, S., Marx, B. (2013) *Regression Model, Method and Applications*. Springer-Verlag New York 293-305
- Gemperli, A. (2003) *Development of spatial statistical methods for modelling point-referenced spatial data in malaria epidemiology*. Doctoral Thesis, University of Basel, Faculty of Science. http://edoc.unibas.ch/diss/DissB_6939 Accessed 12th April 2020 11:30 A.M.
- Guerra, M., Sousa, B., Mabale, N. N., Berzosa, P., Arez, A. P. (2018) Malaria Determining Risk Factors at the Household Level in two Rural Villages of Mainland Equatorial Guinea. *Malaria Journal* **17**(203). <https://doi.org/10.1186/s12936-018-2354-x>
- Hastie, T. and Tibshirani, R. (1990) *Generalized Additive Models*. Chapman & Hall/CRC: 86-173
- Herrera, S., Ivanovich, E., Ye, Y., Garley, A. (2020) *Surveillance, Monitoring and Evaluation of Malaria Programs: Online Course*. Chapel Hill, NC, USA: Measure Evaluation, University of North Carolina. 2-7
- Kalipeni E. (1993) Determinants of infant mortality in Malawi: a spatial perspective, *Social Science and Medicine* **37** 183–198
- Kazembe, L. N., Kleinschmidt, I. and Sharp, B. L. (2006) Patterns of Malaria-related Hospital Admissions and Mortality among Malawian Children: An Example of Spatial Modelling of Hospital Register Data. *Malaria Journal* **5**(93). <http://www.malariajournal.com/content/5/1/93>
- Kazembe, L. N., Chirwa, T. F., Simbeye, J. S. and Namangale, J. J. (2008) Applications of Bayesian Approach in Modelling Risk of Malaria-related Hospital Mortality. *BMC Medical Research Methodology*, **8**(6). <http://www.biomedcentral.com/1471-2288/8/6>.
- Lim, S. S., Fullman, N., Stokes, A., Ravishankar, N., Masiye, F., Murray, C. J. L., Gakidou, E. (2011) Net Benefits: A Multicountry Analysis of Observational Data Examining Associations between Insecticide-Treated Mosquito Nets and Health Outcomes. *PLoS Med* **8**(9): e1001091. doi:10.1371/journal.pmed.1001091.
- Lindblade, K. A., Eisele, T. P., Gimnig, J. E., Alaii, J. A., Odhiambo, F., Kuile, F. O., Hawley, W. A., Wannemuehler, K. A., Howard, P. A. P., Rosen, D. H., Nahlen, B. L., Terlouw, D. J., Adazu, K., Vulule, J. M. and Slutsker, L. (2004) Sustainability of reductions in malaria transmission and infant mortality in western Kenya with use of insecticide-treated bednets: 4 to 6 years of follow-up. *Journal of the American Medical Association (JAMA)* **291**: 2571–2580.
- NDHS (2018) National Population Commission (NPC) [Nigeria] and ICF. 2019. *Nigeria Demographic and Health Survey 2018 Key Indicators Report*. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF
- Nigeria Malaria Fact Sheet (2016) <http://photos.state.gov/libraries/nigeria/231771/Public/December-MalariaFactSheet2.pdf> Accessed 17-03-2020 9:30 AM
- Ogunsakin, R. E. and Chen, D. G. (2020) Bayesian Spatial-Temporal Disease Modelling with Application to Malaria. *Statistical Methods for Global Health and Epidemiology Principles, Methods and Applications*. Springer Nature Switzerland AG. 319-334 <https://doi.org/10.1007/978-3-030-35260-8>

- Oguoma, V.M.; Anyasodor, A.E.; Adeleye, A.O.; Eneanya, O.A.; Mbanefo, E.C. (2021) Multilevel Modelling of the Risk of Malaria among Children Aged under Five Years in Nigeria. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **115**, 482–494
- Okango, E., Mwambi, H., Ngesa, O., Achia T. (2015) Semi-Parametric Spatial Joint Modeling of HIV and HSV-2 among Women in Kenya. *PLoS ONE* **10**(8). doi:10.1371/journal.pone.0135212
- Rowe, A. K., Rowe, S. Y., Snow, R. W., Korenromp, E. L., Joanna, R. M., Schellenberg, A., Stein, C., Nahlen, B. L. and Bryce, J. (2006) The Burden of Malaria Mortality Among African Children in the Year 2000. *International Journal of Epidemiology*, **35**(3): 691–704. doi:10.1093/ije/dyl027.
- Rowe, A. K. and Steketee, R. W. (2007) Predictions of the Impact of Malaria Control Efforts on All Cause Child Mortality in Sub-Saharan Africa. *American Journal of Tropical Medicine and Hygiene* **77**(Suppl. 6): 48-55
- Rumisha, S. F., Smith, T. A., Masanja, H., Abdulla, S., Vounatsou, P. (2014) Relationship between child survival and malaria transmission: an analysis of the malaria transmission intensity and mortality burden across Africa (MTIMBA) project data in Rufiji demographic surveillance system, Tanzania. *Malaria Journal*, **13**(124). <http://www.malariajournal.com/content/13/1/124>
- Salemani, M., Msengwa, A. S., Mrema, S., Shamte, A., Mahande, M. J., Yeates, K., Mbago, M. C. Y., Lutanbi, A. M. (2016) Assessing the Effects of Mosquito Nets on Malaria Mortality Using a Space Time Model: A Case Study of Rufiji an Ifakara Health and Demographic Surveillance System Sites in Rural Tanzania. *Malaria Journal* **15**(257). <http://doi.org/10.1186/s12936-016-1311-9>
- Saroj, R. K., Murthy, K. H., Kumar, M., Singh, R., Kumar, A. (2019) Survival Parametric Models to Estimate the Factors of Under-Five Child Mortality. *Journal of Health Research and Reviews*. **6**:82-8
- Seyoum, S. (2018) Analysis of Prevalence of Malaria and Anemia Using Bivariate Probit Model. *Annals of Data Science* **5**(2):301–312
- Siya, A., Kalule, B. J., Ssentongo, B. Lukwa, A. T., Egeru, A. (2020) Malaria Patterns Across Altitudinal Zones of Mount Elgon Following Intensified Control and Prevention Programs in Uganda. *BMC Infectious Diseases* **20**(425). <https://doi.org/10.1186/s12879-020-05158-5>
- Smith T.A., Leuenberger R., Lengeler C. (2001) Child mortality and malaria transmission intensity in Africa, *Trends in Parasitology* **17** 145–149.
- Snow, R. W. and Marsh, K. (1995) Will Reducing *P. falciparum* Transmission Alter Malaria Mortality Among African Children? *Parasitology Today* **11**: 188-190.
- Snow, R.W., Craig, M. H., Deichmann, U. and Marsh, K. (1999a) Estimating Mortality, Morbidity and Disability due to Malaria Among Africa's Non-pregnant Population. *Bulletin of the World Health Organization*, **77**(8).
- Snow, R.W., Craig, M. H., Deichmann, U. and le Sueur, D. (1999b) A Preliminary Continental Risk Map for Malaria Mortality among African Children. *Parasitology Today*, **5**(3): 100-104.

- Snow, R. W., Korenromp, E. L., Gouws, E. (2004) Paediatric Mortality in Africa: *Plasmodium Falciparum* Malaria as a Cause or Risk? *American Journal of Tropical Medicine and Hygiene*, **71**(Suppl 2), 16–24
- Spencer, H.C., Kaseje, D. C. O., Mosley, W. H., Sempebwa, E. K. N., Huong, A. Y., Roberts, J. M. (1987) Impact on Mortality and Fertility of a Community-based Malaria Control Programme in Saradidi, Kenya, *Annals of Tropical Medicine and Parasitology*, **81**(1), 36-45, <http://dx.doi.org/10.1080/00034983.1987.11812187>
- Streatfield, P. K., Khan, W. A., Bhuiya, A., Hanifi, S. M. A., Alam, N., Diboulo, E., Sié, A., Yé, M., Compaoré, Y., Soura, A. B., Bonfoh, B., Jaeger, F., Ngoran, E. K., Utzinger, J., Melaku, Y. A., Mulugeta, A., Weldearegawi, B., Gomez, P., Jasseh, M., ... Byass, P. (2014) Malaria Mortality in Africa and Asia: Evidence from INDEPTH Health and Demographic Surveillance System Sites, *Global Health Action*, **7**(1), 25369, DOI: [10.3402/gha.v7.25369](https://doi.org/10.3402/gha.v7.25369)
- Sudarno, A. P. (2018) Modelling Infant Mortality Rate in Central Java, Indonesia use Generalized Poisson Regression Method. *Journal of Physics*, **1025**. doi :[10.1088/1742-6596/1025/1/012106](https://doi.org/10.1088/1742-6596/1025/1/012106)
- Trape, J. R. and Rogier, C. (1996) Combating Malaria Morbidity and Mortality by Reducing Transmission. *Parasitology Today*, **12**:236-240.
- Ugwu, C. L. J. and Zewotir, T. T. (2018) Using Mixed Effects Logistic Regression Models for Complex Survey Data on Malaria Rapid Diagnostic Test Results. *Malaria Journal* **17**(453). <https://doi.org/10.1186/s12936-018-2604-y>