

Seed quality behaviour variation of moth bean during storage in different packaging

ABSTRACT: An experiment was carried out in order to assess the behavior of the moth bean seed quality in 4 types of packaging including cloth, gunny, HDPE, and vacuum-packed bags for 18 months. After two months of storage, the moisture content of the seeds fluctuated greatly in cloth, gunny, and HDPE bags (conventional packaging) because these materials are pervious, but the moisture content of the seeds did not vary in vacuum packed bags as the polythene bag used for the vacuum package was thicker, had a lower water vapor and oxygen transmission rate. Further, seeds in conventional packaging's developed bruchid infestations after 4 months of storage but no infestation was found in vacuum-packed bags until 18 months. The bruchid infestation caused the germination rate of the seeds to drop to less than 20%. In comparison to the initial state, even after 18 months of storage, there was little change in the parameters used for measuring seed quality (germination, total seedling length, seedling dry weight, moisture content, and protein content). Therefore, without the use of chemicals, moth bean seeds can be safely stored using vacuum packaging technology.

Keywords: Bruchids, Conventional Packaging's, Moth bean seeds and Vacuum packaging

1. INTRODUCTION

Variations in temperature, rainfall patterns and the frequency and severity of climatic events such as floods and droughts may have a significant impact on agricultural production [1]. Furthermore, a shortage of suitable processing and storage facilities is causing significant damage to food grains following harvest. The FAO calculated that 10% of grains are lost annually worldwide during storage [2]. Multiple strategies are needed to ensure global food security, such as reducing post-harvest loss and improving and diversifying staple food crops. In India, the risk of damage from post-harvest storage to the tune of 25–30% is a serious problem [3] for pulses because of inadequate storage facilities, as well as a lack of expertise in post-harvest pulse management and storage.

Short-duration pulses, like moth bean, need to be preserved for a long time in order to be available year-round as a grain or as seeds. Therefore, in order to meet demand, these seeds need to be properly preserved. As pulses contain significant amounts of protein and are more susceptible to insect infestation, pulses are harder to store than cereals [4–5].

Callosobruchus spp., (Bruchids) are a major pest of grain legumes that can cause significant economic losses in a variety of pulses, including chickpea, mung, peas, cowpea, lentil, and arhar, both in the field and in storage [6, 7]. Because of the bruchids' higher rate of multiplication, which can result in a 30-fold increase in population with each generation that lasts for roughly four weeks [8]. The seeds can be treated with different hazardous chemicals such as methyl bromide, carbon disulfide, or phosphine, or numerous insecticides can be dust-applied to suppress these bruchids. Though these substances are effective, but their use in storage is discouraged since it threatens food safety and the environment [9].

Some plant-derived extracts, such as banana plant juice, soybean, maize and neem oil, and hot pepper powder, have shown benefits in addition to chemical control [10, 11]. Despite their many benefits, these plant-derived extracts have certain drawbacks, such as their slow action and easy degradation. Due to the drawbacks of using artificial and plant-derived chemicals to manage bruchids during storage, studies on chemical-free storage have been carried out to assess the storability of moth bean seeds using various packaging materials.

2. MATERIALS AND METHOD:

2.1) A storage experiment was conducted under ambient conditions at the University of Agricultural

Sciences, Dharwad, Karnataka, India for 18 months (15 November 2019 to 15 May 2021). Average data related to temperature ($^{\circ}\text{C}$) and relative humidity (%) that prevailed in the storehouse of ambient condition were recorded with the help of Anemometer. Freshly harvested healthy seeds of moth bean were collected from farmers' fields and then sun-dried. After sun drying, seeds were packed in different packaging cloth viz., cloth bag, gunny bag, HDPE bag and also vacuum packed. The seeds (3 kgs) were (Fig. 1) packed in the cloth, gunny and high density polythene bag and replicated five times. However, in the case of vacuum packaging, 1 kg of seeds were packed into 9 bags and replicated five times. It is also important to note that in storage, bags were not stacked, ensuring that each bag was equally exposed to the environment. The storehouse's average temperature and relative humidity is presented in Fig. 2. Table 1 lists the features of the polythene bags used for vacuum packaging. An OLPACK 501/V vacuum packer, made by INTERPRISE-BRUSSELS S.A., BRUXTAINER DIVISION, Belgium, was utilized for vacuum packaging.

2.2) Observation Recorded: Prior to storage, various parameters related to seed quality were measured and recorded, including germination percentage, total length of seedlings (cm), dry weight (mg), moisture content (%), and protein content (%). Further, samples were taken from all treatments on the fifteenth of every other month to evaluate the quality of the seed up to eighteen months.

a) Germination (%) and Total seedling length (cm): Germination (%) was calculated as per the standards described in the International Seed Testing Association, 2013. In order to measure the total length of the seedlings, which is expressed in centimeters, 10 normal seedlings were randomly chosen from each treatment on the last day of the above germination test.

b) Seedling dry weight: Ten standard seedlings, which were utilized to measure seedling length, were wrapped in butter paper and dried for twenty-four hours at seventy degrees Celsius in a hot air oven. Following cooling, the seedlings were weighed, and the average weight was determined.

c) Moisture content (%): It was calculated on a dry weight basis as per the International Seed Testing Association, 2013.

d) Protein content: The protein content was obtained by multiplying per cent nitrogen content of seeds with a factor 6.25 [13].

2.3) Data Analysis: Completely Randomized Design (CRD) was used to test the significance of various variables and the results are presented in graphical format[27].

3. RESULTS AND DISCUSSION

Moth bean seeds were stored in cloth, gunny, HDPE and vacuum-packed bags with 95.5% germination, total seedling length (44.1 cm), seedling dry weight (281.1 mg), moisture content (7-80%), and protein content (23.48%). With the exception of moisture content, none of the characteristics examined changed significantly after two months (Fig. 3). The pervious nature of the packaging materials used to store seeds is the reason for the rise in moisture content of the seeds in cloth, gunny, and HDPE bags. Furthermore, because they are hygroscopic by nature, seeds absorb and release moisture in response to environmental factors. Shankar *et al.* [14] in Blackgram have observed similar results of an increase in moisture content in pervious packaging material.; Malimath *et al.* [15] in gardenpa; Khaldun *et al.* [16] in cucumber.

Since vacuum packed bags have a higher thickness and a much lower rate of oxygen and water vapour transmission (Table 1) than pervious packaging materials (cloth, gunny and HDPE bags) and due to these characters, the hygroscopic nature of seeds holds good only when seeds are stored pervious packaging materials. Despite changes in the surrounding environmental conditions, the moisture content of the seeds remained constant during the storage period because of the unique properties of the polythene bag used for vacuum packaging. Similar findings indicating that the moisture content of vacuum-packed seeds remains unchanged have been reported by Chetti *et al.* [17] in chilli; Khanna *et al.* [18] in chickpea; Meena *et al.* [19] in soybean.

With the exception of vacuum-packed bags, there was an infestation of seeds with bruchids to the extent of more than 80 %. The bruchid infestation of seeds in cloth, gunny, and HDPE bags is caused due to the packaging material's permeable nature, which allowed the bruchids to enter material and grow. Similar results of an infestation of seeds with insects in pervious packaging were reported by Patel *et al.* [19] and Khadtar *et al.* [20]. Due to internal feeding behaviour of bruchids, seeds were completely deteriorated and because of complete deterioration there was a reduction in germination and other seed quality parameter's, which caused heavy losses and there by seeds

become unsuitable any purpose. Similar results of seed deterioration due to bruchids infestation in pervious packaging materials have been reported by Charjan *et al.* [21] in arhar; Miah *et al.* [22] in greengram; Swamy *et al.* [23] in black gram, greengram and redgram.

Germination was reduced to less than 20 per cent due to bruchid infestation in conventional packaging materials such as cloth, gunny, and HDPE bags. This was significantly lower than the minimum seed germination of 75 per cent for moth beans, as stated by the Central Seed Certification Board, Department of Agriculture & Co-operation, Ministry of Agriculture, Government of India. Hence, further observations in these treatments have been stopped. Furthermore, because of the type of packaging, the bruchid infestation was not noticed in the vacuum-packed bag even after an 18-month period of storage (Fig 3). Thakur *et al.* [24] in Blackgram reported similar results of non-infestation of seeds with bruchids in vacuum packed bags.

Due to the aging factor of the seeds, there was a slight decline in the percentage of germination, total length of seedlings, seedling dry weight, and protein content of vacuum-packed seeds as the storage period progressed. Our findings are consistent with those of Tripathi *et al.* [25] for onions, Khanna *et al.* [18] for chickpeas, and Ashok *et al.* [26] for onions, who reported that seeds can be stored in vacuum-packed bags for a longer period of time without significant deterioration.

CONCLUSION: The current study's findings indicate that vacuum packaging technology can be used to safely store moth bean seeds for a long period of time without any requirement of chemicals.

ACKNOWLEDGMENT: The authors greatly acknowledge the support Interprise Brussels, Bruxtainer Division (Belgium), Indian Institute of Packaging, Hyderabad and University of Agricultural Sciences, Dharwad

REFERENCES:

1. Arun KC, Aggarwal PK, Joshi PK, Vyas S. Farmers' prioritization of climate-smart agriculture (CSA) technologies. *Agric. Sys.* 2017; 151:184-91.
2. Parfitt J, Barthel M, Macnaughton S. Food waste within food supply chains: quantification and potential for change to 2050. *Philosophical Transactions of The Royal Society B: Biological Sci.* 2010;365(1554):3065-81.

3. Anonymous. Pulses in India: Retrospect and Prospects, 2017b, Government of India, Ministry of Agriculture & Farmers Welfare.
4. Deshpande SD, Singh G. Long term storage structures in pulses. In National Symposium on Pulses for Sustainable Agriculture and Nutritional Security, Indian Institute of Pulses Research, New Delhi 2001;17;17-19.
5. Lal, R. R. and Verma, P. Technical Bulletin – Post Harvest Management of Pulses. 2007; IIPR, Kanpur.
6. Aslam M, Khan KA, Bajwa MZ. Potency of some spices against *Callosobruchus chinensis* Linnaeus. J Bio Sci. 2002;2(7):449-52.
7. Srinivasan T, Durairaj C, Kumar BV. Damage potential of bruchids in different edible legumes and interspecific competition between two species of *Callosobruchus* spp.(Bruchidae: Coleoptera). Madras Agric. J. 2008; 95:1.
8. Sinha RN, Watters FL. Insect pests of flour mills, grain elevators, and feed mills and their control. Ottawa, ON: Canadian Government Publishing Centre, 1985;p. 290.
9. Mahmood I, Imadi SR, Shazadi K, Gul A, Hakeem KR. Effects of pesticides on environment. Plant Soil Microbes. 2016;1: 253-69.
10. Swella GB, Mushobozy DM. Evaluation of the efficacy of protectants against cowpea bruchids (*Callosobruchus maculatus* (F.)) on cowpea seeds (*Vigna unguiculata* (L.) Walp.). Plant Protection Sci. 2007;43(2):68.
11. Rozman V. Control of stored products pests by natural products.2015;111:295-9.
12. ISTA. International Rules of Seed Testing. Seed Sci Technol. 2013; 27:25-30.
13. Sadasivam S, Manickam A. Biochemical methods new age international (P) Ltd. Publishers, New Delhi. 1996.
14. Shankar SV, Pandiarajanb T, Ganapathy S. Biochemical Changes of Black Gram Grains during Hermetic Storage. Res. J. Chemical Environ Sci. 2018;6(6):15-21.
15. Malimath SD, Merwade MN. Effect of storage containers on seed storability of garden pea (*Pisum sativum* L.). Karnataka J Agric Sci. 2007;20(2):384-385.
16. Khaldun ABM. Haque, ME. Seed quality deterioration due to temporal variation of biotic and abiotic factors in cucumber. Bangladesh J Agri Res. 2009; 34(3):457-463.

17. Chetti MB, Deepa GT, Antony RT, Khetagoudar MC, Uppar DS, Navalgatti CM. Influence of vacuum packaging and long term storage on quality of whole chilli (*Capsicum annum L.*). J. Food Sci Technol. 2014;51(10):2827-32.
18. Khanna N, Jain P, Teckchandani CK. Comparative Study of Quality and Nutritive Parameters of Insect Infested Bengal Gram under Vacuum and Modified Atmosphere Storage in Laminated LDPE Bags. Int J Cur Microbiol App Sci. 2017;6(12):4303-4308.
19. Patel JV, Antala DK, Dalsaniya AN. Influence of different packaging materials on the seed quality Parameters of chickpea. Int J Current Microbiology Applied Sci. 2018;7(12):2458-67.
20. Khadtar UM, Shinde AK, Patil PP, Bhave SG. Physiological Studies on Effect of Different Packaging Material for Storage of Cowpea Seed. Adv Agric Res Techno J. 2018;2(2):202-5.
21. Charjan SKU, Gadewar RD, Ciiarde, PN, Lambat AP, Lambat PA. Effect of bruchids infestation on seed quality of arhar During storage. J Phytological Res. 2010;23(2): 383-384.
22. Miah MA, Ali MR, Husna A. Mollah MM. Efficacy of some botanicals against pulse beetle, *Callosobruchus maculatus* (Fab.) on stored greengram, *Vigna radiata*. Bangladesh J. Entomol. 2013;23: 11-20.
23. Swamy SG, Wesley BJ, Raja, DS. Vishnuvardhan, S. Feasibility of sand layer technique for small scale storage of pulses seed. J. Entomol Zoology Studies. 2018; 6(6):428-431.
24. Thakur V, Nawalagatti CM, Chetti MB, Hilli JS, Patil RV. Changes in Blackgram Seed Quality during Storage as Influenced by Various Packaging Materials and Storage Conditions. Indian J Ecology. 2022;49(6):2154-9.
25. Tripathi PC, Lawande KE. Effect of seed moisture and packing material on viability and vigour of onion seed. J Eng Computers Appl Sci. 2014;3(7):1-5.
26. Ashok BG, Doddagoudar SR, Vasudevan SN, Patil MG, Hosamani A. Evaluation of the Best Storage Methods for Maintaining Seed Quality of Onion. Int J Cur Microbiol Appl Sci. 2019;8(4):325-336.
27. Thakur V, Nawalagatti CM. Seed Health of Chickpea as Influenced by Different Packaging during Storage. International Journal of Plant & Soil Science. 2021 Sep 11;33(19):63-70.

S. No.	Characters	Unit	Results
1	Thickness (Microns)	Microns	149.40
2	Water vapor transmission rate	$\text{g/m}^2/24 \text{ hrs at } 38^\circ\text{C and } 90.0 \% \text{ Relative humidity}$	0.95
3	Oxygen transmission rate	$\text{cc}/(\text{m}^2 \times \text{day} \times \text{atm})$	0.91

Table 1. Characters of polythene bag used for vacuum packaging



Fig. 1. Different packaging used for moth bean storage




Fig. 2. Average temperature and relative humidity of store house



Fig.3. Germination (%), Total seedling length (cm), Seedling dry weight (mg), Moisture content (%) and Protein content (%) of moth bean seeds as influenced by different packaging at different time intervals of storage

Note: Observations in cloth, gunny, and HDPE bag treatments have been discontinued as a result of bruchid infestation after 4 months of storage