

Effect of organic manures and natural farming on quality and economics of carrot

Abstract-

The objective of the experiment was to examine the effects of organic manures and natural farming on the quality and economics of carrots. It was conducted at the research farm of the Deen Dayal Upadhyaya Centre of Excellence for Organic Farming of CCS Haryana Agricultural University, Hisar, during the *rabi* season 2022–2023. Three replications and eight treatments were included in the RBD design of the experiment. Three different organic manures were applied: FYM, vermicompost, poultry manure and in combination with biofertilizers (*Azotobacter* and PSB). Treatment of cow based bio-formulations (Ghanajeevamrit and Jeevamrit) and control was also included as treatment where no biofertilizers and manure were used. The study revealed significant improvements in almost all quality attributes due to the combined application of organic manures along with biofertilizers. Among different treatments, vermicompost + biofertilizer surpassed all other treatments by giving maximum root yield (283.4 qha⁻¹), maximum content of the total sugar (8.79%), reducing sugar (4.09%), non-reducing sugar (4.64%), TSS (11.5°Brix) and Chlorophyll content in leaves (4.37 nmol cm⁻²). Also, the treatment vermicompost + biofertilizers gave maximum gross return (Rs. 425100 ha⁻¹), net return (Rs. 232873 ha⁻¹), and BC ratio (2.21) during the experiment. It was also observed during the study that control treatments showed lowest findings among all the treatments.

Keywords- Biofertilizers, Organic manures, Quality, Sugar content.

Introduction

Vegetables are thought to be an essential dietary supplement that can help to maintain overall health and protect against some illnesses that are degenerative. Carrot (*Daucus carota* L.) is a winter season crop and member of the Apiaceae family and is utilised in human diets for a variety of purposes. It contains high amount of nutrients. India's main vegetable crop is carrot. India's major states for the production of carrots include Uttar Pradesh, Haryana, Andhra Pradesh, Karnataka, and Punjab.

Carrot roots are consumed as a raw vegetable in salad form or in boiled form, also soups and pickle is prepared from the carrot (Anjum and Amjad, 2002). Many refreshing drinks which is healthy and tasty is produced from carrot with combination of other vegetables (in combination with other vegetables). Carrot is the source of beta-carotene (precursor of vit- A). Carrot helps in maintaining nitrogen balance and increase the quantity of urine (Anjum and Amjad, 2002).

Organic manures are prepared from the product of plant or animal after decomposing. But raw materials are not collected from unsafe sources (such as industrial waste, toxic waste, hospital waste etc.). Using technologies like the frequent application of synthetic chemicals like pesticides and fertilisers, the adoption of high-yielding varieties, increased use of irrigation potentials, etc., can lead to a "green revolution" and increase production output in most situations. However, the persistent and careless utilisation of these substantial resources is currently causing a reduction in the yield and

efficiency of many crops, in addition to worsening the conditions of the soil and surrounding environments.

The organic manure could increase the fertility and productivity of the land and improve physical and chemical properties of soil as well as produce nutritive and chemical residue free vegetables (Ramesh *et al.*, 2005). The demand for organic produce is increasing with increasing awareness (Bhatta *et al.*, 2009). Both domestically and internationally, there is a growing demand for organic veggies. Consumers expect organic food to have a higher nutritional value, while being free of chemical residues (Ditlevsen *et al.* 2019). Food/vegetables that are produce under the organic system of cultivation contain a higher amount of secondary metabolites, vitamins and various mineral nutrients (Seufert and Ramankutty 2017). Organic manures are the alternate of the chemical fertilizers. Manures supply the required nutrients for the long time and in adequate amount, improve soil structure, increase microbial population and at the same time maintain the quality of crop produce (Wong *et al.*, 1999, Suresh *et al.*, 2004 and Dauda *et al.*, 2008). Farm yard manure is a natural source of available nitrogen. It adds humus and slow releasing nutrients to the soil and restores soil fertility by improving a wide range of natural properties of soil. Vermicompost is a valuable organic fertilizer. Earthworm promotes rapid decomposition with increased rate of mineralization, humification of organic matter and increased microbial biomass that improves the quality of the final product (Atiyeh *et al.*, 2002). Poultry manure is a more concentrated source of crop nutrients, especially nitrogen, phosphorus, potassium and calcium.

Material methods

The experiment was carried out at Research farm of Deen Dayal Upadhyaya Centre of Excellence for Organic Farming of CCS Haryana Agricultural University, Hisar during the *rabi* season 2022-23 under organic system of cultivation without using any chemicals is located at 29°8'20" N latitude and 75°42'04" E longitude at an elevation of 213 metre above mean sea level (MSL) in the subtropics of the country and characterized by semi-arid climatic zone. The soil of experimental field was sandy loam in texture containing 157, 15.70 and 364.00 kg ha⁻¹ available nitrogen, phosphorus and potassium, respectively in 0-15 cm soil depth with EC 0.78 dS m⁻¹, pH 7.80 and organic carbon content 0.67 per cent. The treatment combinations were setup as per recommended dose of Nitrogen for carrot crop in that is 60 kg ha⁻¹. The experiment was laid out in a randomized block design (RBD) with three replications and 8 treatment combinations. Seed of carrot cultivar Punjab carrot 161 was sown in net plot of 4.5m × 4.0m on 16 October, 2023. The treatments were

T₁= RDN through FYM

T₂= RDN through vermicompost

T₃= RDN through poultry manur

T₄= RDN through FYM + biofertilizers*

T₅= RDN through vermicompost + biofertilizers*

T₆= RDN through poultry manure + biofertilizers*

T₇= cow based bio-formulations (Ghanajeevamrit and Jeevamrit)

T₈= control.

(**Azotobacter* + PSB)

Whereas, organic manures i.e., FYM, vermicompost, poultry manures and biofertilizers applied @ 9210, 3157, 2143 kg ha⁻¹ and 50 ml each of *Azotobacter* and PSB for seed treatment. Cow based bio-formulations (Ghanajeevamrit and Jeevamrit) was applied @ 250 kg Ghanajeevamrit + 250 FYM kg/ha and 125 litre Jeevamrit in three split doses at 20, 40 and 60 DAS @ 5 %, 10% and 10%, respectively. Sugar % was analysed according to Lane and Eyon method. For the measurement of TSS Refractometer was used.

Data collected

Data of quality parameters viz., total sugar %, reducing sugar and non-reducing sugar % and TSS °Brix, respectively were taken from the plant's samples after harvest. Lane and Eynon method as given by Ranganna 1994 was used for estimation of sugar and expressed in percentage. TSS was calculated with the use of refractometer. Economics were taken from each plot and converted into per hectare. The economics were calculated on the basis of DDUCE-OF current price of input and output and stated in the form of net return.

Table 1- Analysis of nutrient composition of organic manures used during experiment

| Sr. no. | Organic Manure | Nitrogen (N) (%) | Phosphorous (P) (%) | Potassium (K) (%) |
|---------|----------------|------------------|---------------------|-------------------|
| 1. | FYM | 0.65 | 0.42 | 0.84 |
| 2. | Poultry manure | 2.80 | 1.59 | 1.72 |
| 3. | Vermicompost | 1.89 | 0.85 | 0.96 |

Result and discussion

Application of organic manures alone and with biofertilizers significantly influence the quality attributes of the carrot viz., total sugar, reducing sugar, non-reducing sugar and TSS nutrient content. The maximum content (8.79%) of the total sugar, reducing sugar (4.09%), non-reducing sugar (4.64%) and TSS (11.5°Brix) was recorded in T₅ with the application of RDN through vermicompost and seed was treated with the biofertilizers, which was at par with the treatment of T₂ and T₄ (table 2). However, in total sugar and non-reducing sugar treatment T₅ was also at par with the treatment T₁. The lowest or minimum content of total sugar, reducing sugar, non-reducing sugar and

TSS were recorded with the control treatment. The greater availability and absorption of nutrients in combination with manures contributed to the balanced C/N ratio and increased activity of plant metabolism, which may improve nutrition quality features in carrot as suggested by Emura and Hosoya (1979). Manures that include higher nitrogen levels appear to have encouraged photosynthetic activity and rapid vegetative growth, which in turn led to a bigger accumulation of food material i.e., carbohydrates and an increase in the TSS content of plants. The results were in close conformity with Wafaa (2013) and Mahala *et al.* (2018).

The data depicted in table 2 showed that there was no significant difference observed in chlorophyll content at 30 DAS, whereas at 60 and 90 DAS, application of organic manures with or without biofertilizers and cow based formulation significantly affected the chlorophyll content in leaves.

At 60 DAS, chlorophyll content was recorded significantly higher (3.23 nmol cm⁻²) in T₅ which was at par with T₁, T₂, T₄ and T₆ (3.11, 3.06, 3.15 and 2.92nmol cm⁻², respectively) and higher than all other treatments. At 90 DAS, the same trend was observed and found that T₅ that received RDN through vermicompost + biofertilizers have significantly maximum chlorophyll content (4.37 nmol cm⁻²), and was significantly higher than all other treatments except T₂ and T₄ (3.93 and 4.15 nmol cm⁻², respectively). Moreover, treatment cow based formulations showed no significant differences in chlorophyll content (3.01 nmol cm⁻²) as compared to the control (2.95 nmol cm⁻²). Total chlorophyll content is directly proportional to the availability of the nitrogen that plays an important role in the process of photosynthesis, also the increasing of nitrogen content and leaf area with organic fertilizers directly correlated with an increase in total chlorophyll content (Larimi *et al.*, 2014). The similar results were also found by Zeid *et al.* (2015) and Schmidt *et al.* (2018).

Economics

It was resulted that the application of recommended dose of nitrogen through vermicompost and seed treated with biofertilizer recorded the highest cost of cultivation (Rs. 192227), gross return (Rs. 425100), net return (Rs. 232873) and best benefit cost ratio (2.21) (table 3), which was highest because of higher root yield per hectare with application of RDN through vermicompost along with biofertilizers as compared to other treatments.

Table 2: Effect of organic manures and natural farming on total sugar, reducing sugar and non-reducing sugar and TSS

| Treatments | Total sugar (%) | Reducing sugar (%) | Non-reducing sugar (%) | TSS (°Brix) | Chlorophyll content in leaves | | |
|----------------|-----------------|--------------------|------------------------|-------------|-------------------------------|--------|--------|
| | | | | | 30 DAS | 60 DAS | 90 DAS |
| T ₁ | 8.47 | 3.94 | 4.55 | 10.2 | 1.93 | 3.06 | 3.65 |
| T ₂ | 8.64 | 3.98 | 4.5 5 | 10.9 | 2.00 | 3.11 | 3.93 |

| | | | | | | | |
|----------------|------|------|------|------|------|------|------|
| T ₃ | 8.14 | 3.89 | 4.25 | 10.0 | 1.83 | 2.83 | 3.31 |
| T ₄ | 8.53 | 4.04 | 4.57 | 10.6 | 2.10 | 3.15 | 4.15 |
| T ₅ | 8.79 | 4.09 | 4.64 | 11.5 | 2.13 | 3.23 | 4.37 |
| T ₆ | 8.22 | 3.89 | 4.33 | 10.1 | 1.83 | 2.92 | 3.43 |
| T ₇ | 8.01 | 3.83 | 4.18 | 9.8 | 1.83 | 2.74 | 3.01 |
| T ₈ | 7.93 | 3.80 | 4.13 | 9.2 | 1.80 | 2.64 | 2.95 |
| SEm (±) | 0.16 | 0.04 | 0.06 | 0.4 | 0.18 | 0.13 | 0.18 |
| CD at 5 % | 0.48 | 0.11 | 0.18 | 1.1 | N.S | 0.38 | 0.56 |

Table 3: Effect of organic manures and natural farming on yield, cost of cultivation, gross return, net return and BC ratio

| Treatments | Root yield (q ha ⁻¹) | Cost of cultivation (Rs/ha) | Gross return (Rs/ha) | Net return (Rs/ha) | BC Ratio |
|----------------|----------------------------------|-----------------------------|----------------------|--------------------|----------|
| T ₁ | 265.9 | 189754 | 398850 | 209096 | 2.10 |
| T ₂ | 275.1 | 192103 | 412650 | 220547 | 2.15 |
| T ₃ | 248.1 | 187697 | 372150 | 184453 | 1.98 |
| T ₄ | 271.2 | 189878 | 406800 | 216922 | 2.14 |
| T ₅ | 283.4 | 192227 | 425100 | 232873 | 2.21 |
| T ₆ | 254.7 | 187845 | 382050 | 194205 | 2.03 |
| T ₇ | 241.7 | 181838 | 362550 | 180712 | 1.99 |
| T ₈ | 220.8 | 178425 | 331200 | 152775 | 1.86 |
| SEm (±) | 7.5 | - | - | - | - |
| CD at 5 % | 22.9 | - | - | - | - |

Conclusion

Based on the experimental results it can be showed that the application of 100% RDN through vermicompost along with biofertilizers (*Azotobacter* and PSB) showed superior performance over other treatments which was at par with the treatments in which recommended dose of nitrogen was provided through vermicompost and RDN through FYM along with biofertilizers. Significantly maximum sugar content, net return (Rs 232879 ha⁻¹), BC ratio (2.21) closely followed by RDN through vermicompost (RS. 220547 and BC ratio 2.15), RDN through farm yard manure + biofertilizers (Rs. 216922 and BC ratio 2.14) and RDN through farmyard manure (Rs. 209096 and BC ratio 2.10) in organic cultivation system of carrot. In addition to enhancing soil health the application of organic based fertilizers can substitute the chemical fertilizers. So, the use of organic manures along with biofertilizers directly benefit to the farmers by reducing the cost of cultivation.




Figure 1: Effect of organic manures and natural farming on chlorophyll content in carrot leaves




Figure 2: Effect of organic manures and natural farming on TSS of carrot




Figure 3: Effect of organic manures and natural farming on cost of cultivation, gross return and net return

UNDER PEER REVIEW

References-

- Atiyeh, R. M., Lee, S., Edwards, C. A., Arancon, N. Q., and Metzger, J. D. (2002). The influence of humic acids derived from earthworm-processed organic wastes on plant growth. *Bioresource technology*, **84**(1), 7-14.
- Anjum MA, Amjad M (2002). Influence of mother root size and plant spacing on carrot seed production. *Journal of Resource Science*, **13**(2): 105-112.
- Bhatta, G. D., Doppler, W., and K., C., K. B. (2009). Potentials of organic agriculture in Nepal. *The Journal of Agriculture and Environment*, **10**, 1-11.
- Dauda SN, Ajayi FA, Ndor E. (2008). Growth and yield of water melon (*Citrullus lonatus*) as affected by poultry manure application. *Journal of agriculture and social sciences.*; 4: 121-124.
- Ditlevsen, K.; Sandøe, P. 2019. Lassen, J. Healthy food is nutritious, but organic food is healthy because it is pure: The negotiation of healthy food choices by Danish consumers of organic food. *Food Quality and Preference.*, **71**, 46–53.
- Emura, K. and Hosoya, T. (1979). Effect of fertilization on the quality and yield of spring sown carrots. *Bull. Saitawa Hort. Expt. Sta.*; pp. 13-23.
- Larimi SB, Shakiba M, Mohammadinasab AD, Vahed MM. 2014. Changes in nitrogen and chlorophyll density and leaf area of sweet basil (*Ocimum basilicum* L.) affected by biofertilizer and nitrogen application. *International Journal of Biosciences*.5(9):256-265.
- Mahala P, Chaudhary M., R, Garhwal O., P. 2018. Yield and quality of rabi onion (*Allium cepa* L.) influenced by integrated nutrient management. *International journal of current microbiology and applied sciences.*;7(5):3313-3321.
- Ramesh, P., Singh, M., and Rao, S.A. (2005). "Organic Farming: Its Relevance to the Indian Context. *Current Sciences*. 88 (4): 561-568.
- Schmidt L, Sorg S, Tittmann S, Max J, Zinkernagel J. 2018. Do extended cultivation periods and reduced nitrogen supply increase root yield and anthocyanin content of purple carrots. *Horticulturae*. 4(7):1-13.
- Seufert, V.; Ramankutty, N. 2017. Many shades of gray—The context-dependent performance of organic agriculture. *Science Advances*;3, e1602638.
- Suresh K., D., Sneh G., Krishna, K., K. and Mad, C., M. (2004). Microbial biomass carbon and Microbial activities of soils receiving chemical fertilizers and organic amendments. *Arich. Agron. Soil Sci.*, 50: 7-641.
- Wafaa, H.M. (2013). Yield, quality and micronutrients uptake of carrot (*Daucus carota* L.) and some soil properties as affected by organic fertilizers and elemental sulphur application. *Egyptian Journal of Soil Science*, **53**(4): 537-54.
- Wong J.W.C, Ma, K.K., Fang, K., M. and Cheung, C. (1999). Utilization of manure compost for Organic farming in Hong. *Bio-resource Technol.*, 67: 6-43.
- Zeid HA, Wafaa HM, Seoud AE, Alhadad WAA. 2015. Effect of organic materials and inorganic fertilizers on the growth, mineral composition and soil fertility of radish plants (*Raphane's sativus*) grown in sandy soil. *Middle East Journal of Agriculture Research.*;4(1):77-87