

PHYTOCHEMICAL CHARACTERIZATION, ANTIOXIDANT AND ANTIMICROBIAL ACTIVITIES OF THE LEAVES OF *COMBRETUM BAUCHIENSE* HUTCH. & DALZIEL (COMBRETACEAE)

ABSTRACT

Aim: The phytochemicals, antioxidant and antimicrobial investigations of *Combretum bauchiense* leaves used in folkloric medicine to treat majorly infectious diseases were explored in the present study.

Study design: Evaluation of antioxidant, antimicrobial and HPLC-DAD profiling of the extract and fractions of the leaves of *Combretum bauchiense*.

Place and Duration of study: Department of Pharmaceutical Chemistry, Enugu State University of Science and Technology, Enugu, Nigeria between January 2021 and May 2023.

Methodology: The methanol extract (MECB), n-hexane fraction (HFCB), ethyl acetate fraction (EFCB) and n-butanol fraction of *C. bauchiense* (BFCB) were screened for phytoconstituents using HPLC-DAD. The antioxidant activity of the extract and fractions were assayed by DPPH free radical scavenging model while the antimicrobial activity of extract and fractions were carried out by agar well diffusion method.

Results: The following phytoconstituents: dukurolide, shanzinmethyl ester, piperchabamide, isovitexin, straitisporolide, kaempferol, apigenin, protocatechuic acid, aureonitol, luteolin, rutin and quercetin were identified from the extracts and fraction by dereplication based on their retention time and wavelength. The result of antioxidant assay indicated that MECB showed strong free radical scavenging activity with IC_{50} value of 3.22 $\mu\text{g/mL}$. Among the fractions, EFCB and BFCB have moderate free radical scavenging activity (IC_{50} value of 5.69 and 6.35 $\mu\text{g/mL}$ respectively), compared to that of ascorbic acid (IC_{50} value of 0.21 $\mu\text{g/mL}$). HFCB showed the least free radical scavenging activity (IC_{50} value of 10.78 $\mu\text{g/mL}$).

Conclusion: The extract and the fractions at the concentration of 6.25-100 mg/mL showed inhibition zone diameters, which ranged from 10-27 mm against all tested bacteria. The antioxidant and antimicrobial properties, exhibited by the extracts and fractions may be as a result of the presence of high concentrations of the phenolic compounds identified by HPLC-DAD analysis. The results further substantiated the folkloric uses of the plant in treating infections and some disease conditions associated with oxidative stress.

Keywords: *Combretum bauchiense*; HPLC-DAD; Phytoconstituents; Antioxidant; Antimicrobial

INTRODUCTION

The genus *Combretum* are well known in African traditional medicine, and all parts of the *Combretum* species are used as medicine in treating stomachache, wound healing, edema, skin diseases, diarrhea, muscle pain, and malaria [1-4]. The major compounds derived from the *Combretum* genus include but not limited to the flavonoids, terpenoids and stilbenoids as previously reported [5-8]. Several scientific work have been reported on the potentials of combretum species as anti-inflammatory, antimicrobial, antimalarial, hepatoprotective, antioxidants, anthelmintics and cytotoxicity [9-15].

Combretum bauchiense is one of the poorly investigated species of combretum. It is a suffrutex with short, erect, usually herbaceous stems arising from a woody root stock. It appears in savanna soon after fire and lowers within a few weeks. The leaves are simple, opposite, whorled, sub opposite or alternate, almost always entire [16]. The plant is traditionally used in treating feverish condition, infectious diseases and as tonic. The biological activities of *C. bauchiense* reported so far are antidiarrheal, analgesic and antipyretic and antibacterial [17-19]. The preliminary phytochemical investigation revealed the presence of flavonoids, tannins, cardiac glycosides, terpenoids, and saponins. From the previous studies on the other species of combretum, they demonstrated a significant biological activities due to presence of bioactive compounds. This study was undertaken because of paucity of information on the phytochemicals and pharmacological properties of *Combretum bauchiense*. Therefore, we investigated for the first time the antimicrobial and antioxidant activities of extract and fractions of *Combretum bauchiense* leaves and employed HPLC-DAD analysis for the online detection and identification of the bioactive constituents.

MATERIALS AND METHODS

COLLECTION AND PREPARATION OF PLANT MATERIAL

The leaves of *Combretum bauchiense* were collected in Nsukka, Enugu State, Nigeria on the 23rd of January 2021. The plant was authenticated by Mr Felix Nwafor of Department of Pharmacognosy and Environmental Medicine University of Nigeria, Nsukka, Enugu State, Nigeria. A voucher specimen has been deposited at the herbarium (herbarium number 019/02). The leaves were washed on tap water, air dried for 14 days. The air dried leaves were pulverized using a laboratory scale slow speed electric blender to obtain a homogenous sample which was used for the extraction

CHEMICALS AND EQUIPMENT

Analytical grades of methanol, n-hexane, ethyl acetate and n-butanol (Honeywell, Germany), ascorbic acid (Qualikems), DPPH (Sigma Aldrich Germany) Levofloxacin (Kinvox), ketoconazole capsules (Greenlife), Nutrient agar (by TM Media United Kingdom), Sabouraud Dextrose Aga (by TM Media United Kingdom), Isopropyl alcohol (Moko Methylated spirit), distilled Water were used. HPLCMethanol (LiChroSolv HPLC; Merck) and nano-pure water (distilled and heavymetals free water obtained by passing distilled water through nano- andion-exchange filter cells; Barnstead, France). All other chemicals used were of AnalaR grade. Concentration of extract and fractions were done in rotary evaporator (Digital) TT-52 (by Techmel and Techmel, USA), Pressure cooker autoclave (by Bondtech Corporation, USA), incubator (by Genlab), and UV-visible spectrophotometer (6505, Jenway, England).

MICROORGANISMS

The microorganisms used in this research work are pathogenic in nature and were obtained from few different patients that visited Adonai Biomedical and Microbiological Laboratory and Research Centre, Nsukka. The test organisms were six in numbers comprising of two (2) gram positive bacteria, three (3) gram negative bacteria and one fungus. These organisms includes: *Staphylococcus aureus* and *Streptococcus pneumoniae*; *Salmonella typhi*, *Escherichia coli* and *Klebsiella pneumoniae*; and *Candida albican*. These microbial specimens were cultured and their pure isolates were obtained.

EXTRACTION AND FRACTIONATION

The pulverized plant sample weighing 1kg was macerated in methanol (7 L) for 72 hours with intermittent agitation. Thereafter the mixture was filtered, concentrated using rotary evaporator under reduced pressure, to obtain the methanol extract of *Combretum bauchiense* (MECB). The MECB weighing 100g was dispersed in 200 mL of water containing 20% methanol. The reconstituted solution was subjected to liquid-liquid extraction for fractionation based on solvent polarity using (n-hexane; 500 mL x 3), (ethyl acetate; 500 mL x 4) and (n-butanol; 300 mL x 2). The fractions were concentrated using rotary evaporator under reduced pressure to afford hexane fraction of *Combretum bauchiense* (HFCB), ethyl acetate fraction of *Combretum bauchiense* (EFCB) and buthanol fraction of *Combretum bauchiense* (BFCB) respectively.

HPLC-DAD PROFILING OF EXTRACT AND FRACTIONS

Each of the dried crude extract and fractions (2 mg) was dissolved 2 mLof HPLC grade methanol and the mixture was centrifuged at 3000 rpmfor 5 min. Then, 100 µL of the dissolved samples was transferred intoHPLC vials containing 500 µL of HPLC grade methanol.HPLC analysis was carried out on the samples with a Dionex P580 HPLCsystem coupled to a photodiode array detector. Detection was at 235, 254,280, and 340 nm. The separation column (125 mm x 4 mm; length x internaldiameter) was prefilled with Eurosphere C- 18 (Knauer, Germany), and alinear gradient of nanopure water, adjusted to pH 2 by addition of formicacid and methanol was used as the eluent. Compounds were detected usingdiode array and identified based on similarity with data in the inbuilt library.

DPPH FREE RADICAL SCAVENGING ACTIVITY OF EXTRACT AND FRACTIONS

The antioxidant activity of crude extract and fraction of *C. bauchiense* were determined using the 1, 1-diphenyl- 2-picrylhydrazyl (DPPH) free radical scavenging assay by previously described method [20]. A

stock solution 300 µg/mL of extract and fractions were prepared in methanol and test samples were prepared from stock solution by dilution with methanol to obtain different concentrations ranging from 18.75 µg/mL to 150 µg/mL. DPPH solution was prepared in methanol. A 1 mL of DPPH solution (0.6 mM) was added to 1 mL of different concentrations of the extract and fractions. The volume of the solution was adjusted with methanol to a final volume of 5 mL. The mixture was incubated in the dark for 30 min at room temperature and absorbance of the mixtures was obtained at 517 nm using VIS spectrophotometer (6505, Jenway, England). Ascorbic acid was used as standard. The absorbance of the control containing 1 mL of DPPH solution and 4 mL of methanol was used to calculate the free radical scavenging activities. The percentage free radical scavenging potentials of the extracts, fractions and standard (ascorbic acid) were calculated using the equation below.

Agar diffusion method

The agar well diffusion method previously reported was used [21]. The extract and fractions were dissolved in dimethyl sulfoxide (DMSO) to obtain a final concentration of 100 mg/mL for each of the test agents. From the 100 mg/mL stock solutions of each of the test agents, two (2) fold serial dilutions were done to obtain concentrations of 50 mg/mL, 25 mg/mL, 12.5 mg/mL and 6.25 mg/mL. 0.1mL of 0.5McFarland standard of the test microbial cells suspensions was placed in the petri-dish and 20 mL of the moderately warmed sterilized nutrient agar was added and the mixture was mixed properly by rotating the plate in a circular manner on the working bench for some time to ensure appropriate admixture. This was then allowed to seed at room temperature. Wells of about 5 mm in diameter were made in the seeded agar plate using a 5 mm sterile cork-borer. These wells were filled with 50 µL of the different dilutions of the test solutions using a sterilized microliter pipette. These procedures were followed using 10µg/mL of Levofloxacin as the positive control agent against the test organisms. A triplicate of each of these was obtained. These microbial cultures were then incubated for 24 h at 37°C for bacteria and 24°C for fungi. After the incubation period, the respective inhibition zone diameters around each well were measured in millimeters.

RESULTS

The HPLC-DAD analysis of extract and fractions indicated the presence of bioactive compounds with known pharmacological properties. Most of the bioactive compounds identified in the *C. bauchiense* leaves are moderately polar compounds: flavonoids, terpenoids, fatty acid and their derivatives as shown in Table 1-4 with their respective retention time, molecular weight, molecular formula and reported biological activity.

Table 1: Phytocompounds identified in the MECB

s/n	RT min	Name of compound	Mol. weight	Molecular formula
1	4.23	Protocatechuic acid	154.12	C ₇ H ₆ O ₄
2	6.62	Piperchabamide	411.23	C ₂₆ H ₃₇ N O ₃
3	14.03	Pestalotioprolide	266.33	C ₁₅ H ₂₂ O ₄
4	15.14	4-O-demethylisobutyrolactone	382.5	C ₂₂ H ₃₈ O ₅
5	16.98	Neobavaisoflavone	322.4	C ₂₀ H ₁₈ O ₄
6	18.30	Kaemferol-3-o-gall	594.5	C ₂₇ H ₃₀ O ₁₅
7	14.25	Rutin	610.5	C ₂₇ H ₃₀ O ₁₆




Figure 1: Chemical structures of phytochemicals identified in MECB

Table 2 Phytochemicals identified in HFCB

s/n	RT min	Name of compound	Molecular weight	Molecular formula
1	34.90	Striatisorolide	212.25	C ₁₁ H ₁₆ O ₄
2	35.57	Dukunolide F	484.5	C ₂₆ H ₂₈ O ₉
3	35.96	Piperchabamide H	411.23	C ₂₆ H ₃₇ N O ₃
4	36.39	Genistein -6-c-glucoside	432.4	C ₂₁ H ₂₀ O ₁₀
5	37.05	Shanzinmethyl ester	406.15	C ₁₇ H ₁₆ O ₁₁
6	37.54	Pestalotioprolide H	266.33	C ₁₅ H ₂₂ O ₄
7	39.58	Nidulalin B	302.28	C ₁₆ H ₁₄ O ₆





Figure 2: Chemical structures of phytochemicals identified in HFCB

Table 3: Phytochemicals identified in EFCB

s/n	RT min	Name of compound	Molecular weight	Molecular formula
1	4.85	Protocatechuic acid	154.12	C ₇ H ₆ O ₄
2	16.80	Aplysinamisn-2	650.2	C ₂₃ H ₂₈ Br ₃ N ₃ O ₄
3	22.11	Eriodictiol	288.25	C ₁₅ H ₁₂ O ₆
4	24.20	Luteolin	286.24	C ₁₅ H ₁₀ O ₆
5	25.34	Kaemferol	286.23	C ₁₅ H ₁₀ O ₆

6	26.03	Isovitexin	432.38	C ₂₁ H ₂₀ O ₁₀
7	30.15	Apigenin	270.05	C ₁₅ H ₁₀ O ₅
8	31.32	Aloeresin A	540.6	C ₂₈ H ₂₈ O ₁₁
9	32.14	Aureonitol	206.28	C ₁₃ H ₁₈ O ₂
10	35.90	Benzylpyridine	169.22	C ₁₂ H ₁₁ N
11	37.04	Shanzinmrthyl ester	406.15	C ₁₇ H ₁₆ O ₁₁
12	37.57	Striatisorolide A	212.25	C ₁₁ H ₁₆ O ₄
13	40.64	Nidulalin B	302.28	C ₁₆ H ₁₄ O ₆





Figure 3: Chemical structures of phytochemicals identified in EFCB

Table 4: phytochemicals identified in BFCB

s/n	RT min	Name of compound	Molecular weight	Molecular formula
1	14.77	Cyclo(tyrosylprolyl)	260.28	C ₁₄ H ₁₆ N ₂ O ₃
2	19.27	Rutin	610.5	C ₂₇ H ₃₀ O ₁₆
3	20.93	Apigenin	270.05	C ₁₅ H ₁₀ O ₅

4	37.37	Hydroxyanthranilic acid	153.14	C ₇ H ₇ N O ₃
5	38.47	Oleic acid	282.5	C ₁₈ H ₃₄ O ₂
6	47.18	Cyclophenol	310.3	C ₁₇ H ₁₄ N ₂ O ₄





Figure 4: Chemical structures of phytochemicals identified in BFCB

DPPH FREE RADICAL SCAVENGING CAPACITY DETERMINATION

The antioxidant potential of extract and fractions were determined by DPPH free radical scavenging activity. The extract demonstrated good free radical scavenging activity more than any of the fractions and to an extent comparable to the ascorbic acid (standard). The extract had, inhibition of 76.9% at 300 $\mu\text{g/mL}$ while different solvent fractions showed percentage inhibitions of 31.8% at 300 $\mu\text{g/mL}$ for HFCB, 46.3% at 300 $\mu\text{g/mL}$ for EFCB and 47.41% at 300 $\mu\text{g/mL}$ for BFCB respectively. The ascorbic acid indicated percentage inhibition of 91.8% at 300 $\mu\text{g/mL}$. The antioxidant capacity of the extract and fractions also expressed as IC₅₀, which is the concentration that showed 50% inhibition of the free radical (Table 5). The result indicated that MECB had the lowest IC₅₀ value of 3.22 $\mu\text{g/mL}$ showing good free radical scavenging activity. However, among the fraction EFCB and BFCB have significant free radical scavenging activity (IC₅₀ value of 5.69 and 6.35 $\mu\text{g/mL}$ respectively). HFCB showed the least free radical scavenging activity (IC₅₀ value of 10.78 $\mu\text{g/mL}$).

Table 5: IC₅₀ of the extracts and fractions

Samples	IC ₅₀ Values ($\mu\text{g/mL}$)
MECB	3.22
HFCB	10.78
EFCB	5.69
BFCB	6.35
AA	0.21

KEY: methanol extract of *C. bauchiense* (MECB), n-hexane fraction of *C. bauchiense* (HFCB), ethyl acetate fraction of *C. bauchiense* (EFCB), n-butanol fraction of *C. bauchiense* (BFCB) and ascorbic acid (AA)

ANTIMICROBIAL EVALUATION

The results of the antimicrobial sensitivity test of MECB, HFCB, EFAO and BFCB are shown in Table 6. The extract and the fractions at the concentration of 6.25-100 mg/mL showed inhibition zone diameters, which ranged 10-27 mm against all tested bacteria. The HFCB and EFAO had inhibition zone diameter ranged 10-25 mm against all tested bacteria, at concentration range of 6.25-100 mg/mL following 2-fold serial dilution. The BFCB exhibited inhibition zone diameter of 13-19 mm against only *S. aureus* at 6.25-25 mg/mL whereas 50 -100 mg/mL showed inhibition zone of 10-13 mm against *S. typhi*, *K pneumonia*, and *E coli*. The standard drugs had inhibition zone diameter, which ranged from 27-45 mm at concentration of 0.01 µg/mL.

Table 6: Inhibition zone diameter (mm) of MECB, HFCB, EFCB and BFCB against test organisms

Samples	Conc. mg/ML	IZD (mm)					
		<i>Staphylococcus aureus</i>	<i>Salmonella typhi</i>	<i>Klebsiella pneumonia</i>	<i>Escherichia coli</i>	<i>Streptococcus pneumonia</i>	<i>Candida albican</i>
MECB	100	27 ± 0.82	25 ± 1.41	25 ± 0.74	24 ± 0.82	26 ± 0.47	0
	50	24 ± 0.82	23 ± 0.82	21 ± 0.82	24 ± 0.82	23 ± 0.47	0
	25	20 ± 0	20 ± 0	21 ± 0.82	19 ± 1.63	19 ± 0	0
	12.5	20 ± 0.82	17 ± 0.82	18 ± 0	17 ± 0.82	16 ± 0.82	0
	6.25	17 ± 0.	15 ± 0.82	15 ± 0	14 ± 0	12 ± 0	0
HFCB	100	21 ± 0	19 ± 0.47	20 ± 0	20 ± 0	22 ± 0.82	0
	50	18 ± 0	17 ± 0	16 ± 0.82	19 ± 0.82	19 ± 0	0
	25	16 ± 0.82	15 ± 0	15 ± 0	19 ± 0.82	17 ± 0	0
	12.5	15 ± 0.82	14 ± 0	13 ± 0	17 ± 0.82	14 ± 0	0
	6.25	13 ± 0	12 ± 0.82	12 ± 0	13 ± 0.82	14 ± 0	0
EFCB	100	19 ± 0.82	18 ± 0.82	17 ± 0	20 ± 0	23 ± 0.82	0
	50	18 ± 0	16 ± 0	15 ± 0	18 ± 0.82	21 ± 0	0
	25	15 ± 0	14 ± 0.82	13 ± 0	16 ± 0	19 ± 0	0
	12.5	14 ± 0.82	14 ± 0	12 ± 0	14 ± 0	17 ± 0	0
	6.25	12 ± 0	12 ± 0	11 ± 0	10 ± 0	14 ± 0	0
BFCB	100	13 ± 0	16 ± 0	14 ± 0	13 ± 0	0	0
	50	10 ± 0	13 ± 0	10 ± 0	10 ± 0	0	0
	25	19 ± 0	0	0	0	0	0
	12.5	15 ± 0	0	0	0	0	0
	6.25	13 ± 0	0	0	0	0	0
LEVO	0.01	32 ± 0.05	32 ± 0.05	32 ± 0.05	32 ± 0.05	32 ± 0.5	-
KETO	0.01	-	-	-	-	-	27 ± 034

Key: methanol extract of *C. bauchiense* (MECB), n-hexane fraction of *C. bauchiense* (HFCB), ethyl acetate fraction of *C. bauchiense* (EFCB), n-butanol fraction of *C. bauchiense* (BFCB). LEVO (levofloxacin) and KETO (ketoconazole).

DISCUSSION

The HPLC-DAD analysis of the extract (MECB) and the fractions (HFCB, EFCB & BFCB) of *C. bauchiense* leaves detected a whole range of bioactive compounds, which are majorly of the flavonoids (Tables 1-4; Figures 1-4). The major components identified by dereplication include luteolin, quercetin, kaempferol, apigenin, isovitexin and rutin, which have been shown to demonstrated significant pharmacological activities in previous studies [22-23]. These included but not limited to antimicrobial,

antioxidant, anticancer, anti-inflammatory and antidiabetic activities [24-35]. The extracts and fractions demonstrated a dose dependent inhibition of the DPPH free radicals *in vitro* (Table 5). Our results indicated that EFCB, which showed the best antioxidant activity when compared with the other fractions had the highest concentration of phenolic compounds, mainly flavonoids (Table 3, Figure 3). It has been reported severally that flavonoids class of phytoconstituent possesses very good antioxidant properties [36-37]. Its antioxidant activity is due to presence of many phenolic hydroxyl groups in the molecule. These phenolic hydroxyl groups, particularly when they are in ortho-position have been shown to have immense ability to scavenge free radicals, chelate metal ions and inhibit lipid peroxidation [38]. The strong antioxidant activity of this plant may be associated with the high concentration of these phenolic compounds and this further justifies the use of the plant in the management of diseases associated with oxidative stress.

The preponderance of multi-drug resistant pathogens poses a serious threat to human and animal health. These microbial pathogens continue to develop intrinsic resistance to the available antibiotics [39]. The strategies could be by drug-degrading and -modifying enzymes, reduced membrane permeability for drugs, expression of drug efflux pumps, overproduction and alteration of drug targets, and bypassing the inhibitory pathway [39]. There is thus an urgent need to discover new antimicrobial agents with novel mechanism of action, to combat the multidrug resistance organisms. Our current study showed that the extract and fractions from *Combretum bauchiense* leaves possess broad spectrum antibacterial action but no antifungal property (Table 6). This is in agreement with previously reported antimicrobial activity of the other plants of *Combretum species* [40-41]. The broad spectrum antibacterial activity can also be attributed to the presence of the phenolic compounds in the plant. Flavonoids and other phenolic compounds have been severally reported to exhibit strong antimicrobial activity against resistance pathogens [42-43].

CONCLUSION

The result obtained demonstrated that leaves of *C. bauchiense* possesses antioxidant and antimicrobial properties, which may be as a result of the presence of copious concentrations of the phytoconstituents identified by HPLC-DAD analysis. Our findings also further validated the ethnomedicinal use of the leaves of *C. bauchiense* in the management of infections and various disease affecting man.

REFERENCE

1. Neuwinger HD. African Traditional Medicine. A Dictionary of Plant Use and Applications. Medpharm Scientific Publishers Stuttgart, Germany; 2000 p. 589
2. Fyhrquist P, Mwasumbi L, Hæggström CA, Vuorela H, Hiltunen R, Vuorela P. Ethnobotanical and antimicrobial investigations on some species of Terminalia and Combretum (Combretaceae) growing in Tanzania. J Ethnopharmacol. 2002; 79: 169–177
3. Gathirwa JW, Rukunga GM, Mwitari PG, Mwikwabe NM, Kimani CW, Muthaura CN, Kiboi DM, Nyangacha RM and Omar SA. Traditional herbal antimalarial therapy in Kilifi district, Kenya. Journal of Ethnopharmacology. 2011; 134, 434–442.
4. Nguta JM, Mbarian JM, Gakuya DW, Gathumbi PK and Kiama SG. Antimalarial herbal remedies of Msambweni, Kenya. Journal of Ethnopharmacology 2010;128, 424–432.
5. Welch CR. Chemistry and pharmacology of kinkéliba (*Combretum micranthum*), a West African medicinal plant; PhD Thesis, The State University of New Jersey, 2010; pp. 10-62
6. Kemvoufo PB, Barboni L, Teponno RB, Mbiantcha M, Nguielefack TB, Hee-Juhn P, Kyung-Tae L and Tapondjou LA. Polyhydroxyoleanane-type triterpenoids from *Combretum molle* and their anti-inflammatory activity. Phytochem. Lett., 2008;1(4),183- 187.
7. Martini ND, Katerere, DRP, Eloff, JN. Biological activity of five antibacterial flavonoids from *Combretum erythrophyllum* (Combretaceae). J. Ethnopharmacol. 2004; 93(2-3):207-212

8. Eloff JN, Famakin JO and Katerere DRP. Isolation of antibacterial stilbene from *Combretum Woodii* (Combretaceae) leaves. *Afr.J. Biotechnol.* 2005;4(10), 1167-1171.
9. Ogbu CP, Okagu, IU and Nwodo, OFC. (2019). Anti-inflammatory activities of crude ethanol extract of *Combretum zenkeri* Engl. & Diels leaves. *Comparative Clinical Pathology* 2019;29(2):397-409
10. Ntashanka MN, Ejidike IP, Mthunzi FM, Moloto MJ, Mubiayi KP (2020) Investigation into the phytochemical profile, antioxidant and antibacterial potentials of *Combretum molle* and *Acacia meurnsii* leaf parts. *Biomed pharmacol J* 2020; 13(4)
11. Sanon S, Gansane A, Ouattara LP, Traore A, Ouedraogo IN, Tiono A, et al. In vitro antiplasmodial and cytotoxic properties of some medicinal plants from western Burkina Faso. *Afr J Lab Med* 2013; 2:81.
12. Iyabo A, and Chinenye U. In-vivo hepatoprotective effects of *Combretum micranthum* leaves extract. *The FASEB journal* 35(S1) special issue: experimental biology meeting abstract. 2021
13. Rajalingam D, Varadharajan R, Palani S. Evaluation of Hepatoprotective and Antioxidant Effect of *Combretum albidum* G. Don against CCl₄ Induced Hepatotoxicity In Rats' *International Journal of Pharmacy and Pharmaceutical Sciences* 2016;8(9):218- 223
14. Simon MK, Ajanusi OJ, Abubakar MS, Idris, AL and Suleiman MM. The anthelmintic effect of aqueous methanol extract of *Combretum molle* (R. Br. x. G. Don) (Combretaceae) in lambs experimentally infected with *Haemonchus contortus*. *Veterinary Parasitol*, 2012;187(1-2):280-284
15. Wu XP, Han CR, Chen GY, Yuan Y and Xie JY. Cytotoxic pentacyclic triterpenoids from *Combretum oliviforme*. *Nat. Prod. Commun.*, 2010;5(7):1027-1030
16. Onyekwelu SSC. Germination, Seedling, Morphology and Establishment of *Combretum bauchiense* Hutch and Dalz. (combretaceae). *Botanical journal of the linnean society*, 1990;103 (2) 133-138
17. Odoh U.E, Agubata C.W, Obi P, Chikeoku I, Mba T, Onugwu O & Uzor G. Evaluation of the analgesic and antipyretic activity of methanol extract of *Combretum bauchiense* Hutch & Dalziel (Combretaceae) leaves. *Phytomedicine plus* 2022; 2, 100166.
18. Odoh U.E, Ugwu TN, Obi P, Chikeokwu I, Mba T, Onugwu O, Uzor G and Onyegbulam CM. Quantitative phytochemical analysis and anti-diarrhoeal activity of methanol leaf extract of *Combretum bauchiense* Hutch & Dalziel (combretaceae). *World journal of pharmaceutical research*, 2021;11 (1) 1607
19. Ali IJ, Okorie NH, Ugodi GW, Ujam TN, Ugwu CO, Okwori JM and Okoye FB. (2022) Phytochemical screening and antibacterial evaluation of extract and fractions of *Combretum bauchiense* leaves. *J.Pharmacognphytochem.* 11(2): 285-288
20. Osadebe PO, Okoye FBC, Uzor PF, Nnamani NR, Adiele IE and Obiano NC. Phytochemical analysis, hepatoprotective and antioxidant activity of *Alchornea cordifolia* methanol leaf extract on carbon tetrachloride-induced hepatic damage in rats. *Asian pacific journal of tropical medicine.* 2012;5(4)289-293.
21. Eloff JN. Avoiding pitfalls in determining antimicrobial activity of plant extracts and publishing the results. *BMC complement Altern Med.* 2019; 19, 106

22. Sanchez C. Reactive oxygen species and antioxidant properties from Mushrooms. *Synt and Sys Biotechnol.* 2017; 2:13-22.
23. Shashank K and Abhay KP. (2013) Chemistry and biological activities of flavonoids: An overview. *The scientific World Journal*; 2013:16 pages
24. Joung DK, Kang OH, Seo Y, Zhou T, Lee YS, Han SH, Mun SH, Kong R, Song HJ, Shin DW, Shin DW, et al: Luteolin potentiates the effects of aminoglycoside and β -lactam antibiotics against methicillin-resistant *Staphylococcus aureus* in vitro. *Exp Ther Med.* 2016; 11(6) : 2597-2601
25. Sheng JW, Liu DM, Jing L, Xia GX, Zhang WF, Jiang JR, Tang JB. Striatisorolide A, a butenolide metabolite from *Athyrium multidentatum* (Doll.) Ching, as a potential antibacterial agent. *Mol Med Rep.* 2019; 20(1):198-204
26. Ndjateu FST, Tsafack RBN, Nganon BK, Awouafack MD, Wabo HK, Tene M, Tane P and Eloff JN. Antimicrobial and antioxidant activities of extracts and ten compounds from three Cameroonian medicinal plants: *Dissotis perkinsiae* (Melastomaceae), *Adenocarpus mannii* (Fabaceae) and *Barteria fistulosa* (Passifloraceae). *South African Journal of Botany* 2014; 91, 37-42
27. Izuchi GW, Majekodunmi OF, Micheal D, Yousif MA and John H. Musanol: A new aureonitol-related metabolite from a chaetomium sp. *Chemoinform* 2007; 63(34):8174-8180
28. Liping Y, Wenlei Z, Shaoyang Z, Mingyu L, Mengjuan Z, Chaobin Q, Xiao Y, Junchang F, Guoxing N. Effects of genistein on glucose and lipid metabolism of common carp (*Cyprinus carpio*. L) in vivo and in vitro, *Aquaculture Reports*, 2022; 22, 100930,
29. Wei CC, Yen PL, Chang ST, Cheng PL, Lo YC, and Liao VHC. Antioxidative activity of both oleic acid and camellia tenuifolia seed oil are regulated by the transcription factor DAF-16/FOXO in *Caenorhabditis elegans*. *Plos One* 2016;11(6) e0157195
30. Owumi SE, Bello SA, Idowu TB, Arunsi UO, Oyelere AK. Protocatechuic acid protects against hepatorenal toxicities in rats exposed to Furan. *Drug Chem Toxicol.* 2022 ;45(4):1840-1850.
31. Liu S, Dai H, Makhoulfi G, Heering C, Janiak C, Hartmann R, Mándi A, Kurtán T, Müller WE, Kassack MU, Lin W, Liu Z, Proksch P. Cytotoxic 14-Membered Macrolides from a Mangrove-Derived Endophytic Fungus, *Pestalotiopsis microspora*. *J Nat Prod.* 2016;79(9):2332-40.
32. Zhang Y, Zhang R, Ni H. Eriodictyol exerts potent anticancer activity against A549 human lung cancer cell line by inducing mitochondrial-mediated apoptosis, G2/M cell cycle arrest and inhibition of m-TOR/PI3K/Akt signalling pathway. *Arch Med Sci* 2020;16:446–452
33. Szliszka E, Skaba D, Czuba ZP, Krol W. Inhibition of inflammatory mediators by neobavaisoflavone in activated RAW264.7 macrophages. *Molecules.* 2011;16(5):3701-12
34. Yeon MJ, Lee MH, Kim DH, Yang JY, Woo HJ, Kwon HJ, Kim JB. Anti-inflammatory effects of Kaempferol on *Helicobacter pylori*-induced inflammation. *Bioscience, Biotechnology, and Biochemistry* 2019; 83(1): 166-173
35. Haddad P and Eid H. The antidabetic potential of quercetin: underlying mechanism. *Current medicinal chemistry.* 2017;24 (2) : 355-364
36. Mbikay M, Sirois F, Simoes S, Mayne J and Chretien M. Quercetin-3-glucoside increases low-density lipoprotein receptor (LDLR) expression, attenuates proprotein convertase subtilisin/kexin

9 (PCSK9) secretion, and stimulates LDL uptake by Huh7 human hepatocytes in culture. *FEBS Open Bio*. 2014;4(1):755–762.

37. Okoye FBC, Osadebe PO. A new anti-inflammatory flavonol glycoside from *Alchorneafloribunda* leaves. *Natural Product Research* 2010; 24 (3): 266-273.
38. Wahyuni MH, Malaka NA, Yanti R, Hartati S and Sahidin I. Radical scavenging and antibacterial activity of phenolic compounds from *anacardiumoccidentale* stem barks from south-east Sulawesi- Indonesia. *Indian J Pharm Sci*, 2018;80(1):143-149
39. Ogundipe FO, Bamidele, FA, Adebayo-Oyetoro AO, Ogundipe OO and Tajudeen OK. Incidence of Bacteria with Potential Public Health Implications in Raw *Lycopersicon esculentum* (Tomato) Sold in Lagos State, Nigeria, *Nigerian Food Journal*. 2012; 30(2): 106-113
40. Nounagnon SM, N'tcha C, Sina H, Noumavo AP, DurandDah-Nouvlessounon, Assogba MF, Gbénou DJ and Baba-Moussa L. Antimicrobial activities of *Combretum micranthum* extracts on *Staphylococcus aureus* strains isolated from skin infections and some reference strains. *Journal of Plant Science and Research*. 2016; 6(4):40-47
41. Eloff JN, Famakin JO and Katerere DRP. Isolation of an antibacterial stilbene from *Combretum woodii* (Combretaceae) leaves. *African Journal of Virology Research*. 2020; 14 (2): 001-005
42. Okeke VO, Okoye NN, Ngwoke KG, Okoye FBC. Antimicrobial Screening and HPLC/DAD-MS Characterization of the Flavonoid-Rich Fractions of the Methanol Leaf-Extract of *Lawsoniainermis*Linn. *Trop J Nat Prod Res*. 2021; 5(8):1500-1505
43. Okoye, N. N and Okoye, C. O. B. Antimicrobial and antioxidant flavonoid glycosides from the leaves of *Alstoniaboone*De Wild. *Journal of Pharmaceutical Research International*(formerly British Journal of Pharmaceutical Research). 2016; 10(6):1-9