

Original Research Article

Seasonal occurrence of melon fruit flies (*Zeugodacus cucurbitae* Coq.) and its natural enemies on Bitter gourd

ABSTRACT

Aim: To study the seasonal occurrence of cucurbit melon fly and correlation study with weather parameters to know the impact of abiotic environmental factors on the activity of melon fruit fly concerning infesting the fruits of bitter gourd in crop ecosystem.

Place and duration of study: College of Agriculture, VNMKV Parbhani, Maharashtra. Studies during two seasons Rainy and Summer 2021 and 2022 respectively.

Methodology: The experimental plot was kept unsprayed throughout the crop season. The observation was recorded as soon as the incidence was noticed from five randomly selected plants at weekly intervals. The observations of fruit flies were recorded from the day of fruit formation to the last picking of the fruit. The damaged and healthy fruits were recorded at each picking to knowing the percentage fruit infestation by fruit flies.

Results: Fruit fly was the major pest of bitter gourd in fruit damage. The fruit damage record range was 15.65 to 59.33 per cent in the rainy season whereas, 28.99 to 61.14 per cent in the summer season. The per cent fruit infestation shows a significant and positive correlation with no. of maggots per fruit ($r = 0.857$ and 0.905) during both seasons s respectively. Weather parameters like morning relative humidity (MRH) and rainfall the during rainy season show significant and positive correlation ($r = 0.87$ and $r = 0.71$ respectively) whereas, during the summer season significant and positive correlation ($r = 0.77$) with MRH and significant and negatively correlated with ($r = -0.66$) minimum temperature. The record of larval pupal parasitoid *Psytalia fletcheri* (Silvestri) and it was discovered that 16% of the population was parasitized during the experimental period.

Conclusion: The infestation may vary from season to season, region to region and concerning variety also. Here the infestation range was higher during the summer season, which may be because of the coincidence with the fruit season of mango in that area.

Keywords: Melon fruit fly, Bitter gourd, Weather parameters, Rainy, Summer, correlation

1. Introduction

Melon fly is a serious pest of vegetable crops, especially cucurbits such as melon, pumpkin, squash, zucchini and cucumber. It damages over 81 plant species, but plants belonging to the family Cucurbitaceae are preferred most (Allwood *et al.* 1999). As one of the most serious, even though enough

research has been done so far in India, there is still very little information accessible for melon fly. Moreover, due to regional variations in meteorological circumstances, the seasonal incidence of any pest may vary from place to place.

Fruit flies use a wide spectrum of hosts, from severe monophagy to extreme polyphagy. Some tephritid species, particularly those of the subfamilies Dacinae and Trypetinae, have frugivorous larvae that feed on the fruit pulp of both cultivated and wild plants, giving rise to their common name of "fruit flies." Larvae of [the](#) remaining species feed on stems, shoots and flowers (Christenson and Foote, 1960). It infests more than 125 species of plants, including cucurbits, tomatoes, and many other [vegetables](#), [which](#) have been recorded as hosts of the melon fly. Preferred hosts include: cantaloupe, cowpea, cucumber, gourd, pumpkin, squash, string bean, tomato and watermelon. Occasional hosts include: eggplant, fig, mango, orange, papaya and peach. Wild hosts include: balsam apple; Chinese cucumber, *Momordica spp.*; colocynth; two genera of cucurbits- *Sicyos sp.*; *Cucumistrigonus*; *Diplocyclospalmatus*; and passion-flower, *Passiflora spp.* However, White and Elson-Harris (1994) claim that a lot of these reports might have been based on haphazard observations of adults trapped in traps placed on non-host trees or resting on plants.

As this pest is polyphagous it may cause significant economic losses, which can range from 30% to 100% depending on the crop and season (Dhillon *et al.* 2005). Nowadays *Z. cucurbitae* has [become a](#) threat to intensive agriculture. Therefore, the purpose of the current proposal was to consider the seasonal occurrence of *Z. cucurbitae* and to investigate the impact of meteorological parameters on the population dynamics of the bitter melon crop pests to forecast the most effective management strategies [concerning](#) both space and time for this pest.

2. Materials and Methods:

The experiment was conducted during [the](#) Rainy 2020-21 and Summer 2021-22 [at](#) [the](#) Experimental Farm of [the](#) Department of Agricultural Entomology, Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani Maharashtra. To study the population dynamics of fruit flies [the](#) experimental plot was kept unsprayed throughout the crop season. The observation [was](#) recorded as soon as the incidence was noticed from five randomly selected plants at weekly intervals.

The observations of fruit flies were recorded from [the](#) day of fruit formation to [the](#) last picking of the fruit. The damaged and healthy fruits were recorded at each picking for record fruit infestation by fruit flies by following [the](#) formula. The mean percent fruit infestation was taken and [the](#) standard deviation has been worked out.

$$\text{Percent Fruit infestation} = \frac{\text{Number of infested fruits}}{\text{Total number of Fruits}} \times 100$$

The fruit infestation was subjected to correlation and multiple regression with weather parameters like maximum temperature, minimum temperature, morning relative humidity, evening relative humidity, rainfall, wind velocity, number of rainy days, and sunshine hours. The weather data was collected from the observatory of the Department of Agricultural Meteorology, Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani.

3. Results and Discussion:

3.1 Fruit fly infestation during *Rainy* 2021

The per cent fruit infestations during *Rainy* shows that (Table.1) the per cent fruit infestation varies from 15.65 to 59.33 per cent with a mean of 41.23 per cent of fruit infestation a on number basis whereas, on a weight basis the per cent fruit infestation during this season vary 12.41 to 65.00 per cent with mean 12.41 to 65.00 per cent fruit infestation. The number of maggots per fruit varied from 4.49 to 21.00 maggots per fruit with a mean 11.96 number of maggots per fruit

3.1.1 The correlation study between fruit infestation and maggot count with weather parameters

The correlation studies with weather parameters (Table 2) and Fig.1. shows that a significant and positive correlation existed between per cent fruit infestation (on a number basis) and morning relative humidity. While, non-significant and positive correlation with maximum temperature, minimum temperature, evening relative humidity, rainfall and Rainy days $r = 0.21$, $r = 0.61$, $r = 0.51$, $r = 0.62$ and $r = 0.59$ respectively, non-significant and negative correlated with evaporation, bright sunshine hours and wind speed $r = -0.49$, $r = -0.09$ and $r = -0.59$, respectively.

The correlation between weather parameters and per cent fruit infestation on a weight basis shows (Fig.2) a significant and positive correlation with morning relative humidity and rainfall $r = 0.84$ and $r = 0.71$ respectively. The non-significant and positive correlation has existed with maximum temperature ($r = 0.21$), minimum temperature ($r = 0.55$), evening relative humidity ($r = 0.46$) and Rainy days ($r = 0.63$), non-significant negative correlation with evaporation ($r = -0.48$), bright sunshine hours ($r = -0.02$) and wind speed ($r = -0.644$), respectively.

The correlation with number of maggots per fruit with weather parameter shows (Fig.3) that significant and positive correlation with minimum temperature ($r = 0.73$), morning relative humidity ($r = 0.92$), evening relative humidity ($r = 0.76$), rainfall ($r = 0.89$) and Rainy days ($r = 0.89$) whereas, significant and negative correlation with evaporation ($r = -0.78$). The maggot population was non-significant and negatively correlated with maximum temperature ($r = -0.086$) bright sunshine hours ($r = -0.44$) and wind speed ($r = -0.302$).

3.1.2 The association between per cent fruit infestation and maggots count per fruit during season *Rainy* 2021

The association between [several](#) maggots per fruit and per cent fruit infestation on bitter gourd during *Rainy* season 2021 in (Table 3). It shows that percent fruit infestation both on [a](#) number basis and weight basis showed [a](#) significant and positive correlation with [many](#) maggots per fruit $r = 0.838$ and $r = 0.857$ respectively.

3.2 Fruit fly infestation during *Summer* 2022

The per cent fruit infestation of bitter gourd during [the](#) *Summer* season recorded data presented in (Table 4) it shows that the per cent fruit infestation varies from 28.99 to 47.57 per cent with mean per cent fruit infestation 47.57 per cent on number basis and weight basis varied from 61.93 to 25.84 per cent with mean per cent fruit infestation of 48.66 per cent. While number of maggots per fruit varies from 27.40 to 6.53 maggots per fruit with the mean number of maggots [being](#) 18.59 per fruit during [the](#) *Summer* season.

3.2.1 Correlation study with weather parameters

The correlation association between fruit infestation and their maggots with weather parameters during *Summer* season represented in Table 3 and Fig.4 it shows that percent fruit infestation on number basis was [non-significant](#) and positively correlated with morning relative humidity ($r = 0.52$), evening relative humidity ($r = 0.25$), rainfall ($r = 0.12$), *Rainy* days ($r = 0.13$) bright sunshine hours ($r = 0.17$), wind speed ($r = 0.09$) whereas, [non-significant](#) and negatively correlated with maximum temperature ($r = -0.44$), minimum temperature ($r = -0.37$) and with evaporation ($r = -0.45$).

The per cent fruit infestation recorded on [a](#) weight basis showed [a](#) significant and positive correlation with morning relative humidity ($r = 0.77$) whereas, significant and negative correlated with minimum temperature ($r = -0.66$). [Non-significant](#) and positively correlated with evening relative humidity ($r = 0.58$), rainfall ($r = 0.34$), *Rainy* days ($r = 0.34$) evaporation ($r = 0.62$), wind speed ($r = 0.11$) while [non-significant](#) and negatively correlated with maximum temperature ($r = -0.65$) shown in Fig.5

The correlation study (Fig.6) of [several](#) maggots per fruit during [the](#) *Summer* season with weather parameters shows that significant and positive correlation with the morning relative humidity ($r = 0.88$) and evening relative humidity ($r = 0.83$) whereas, significant but negatively correlated with maximum temperature and minimum temperature $r = -0.73$ and $r = -0.77$ respectively. [Non-significant](#) and positively correlated with rainfall ($r = 0.34$), *Rainy* days ($r = 0.34$), evaporation ($r = 0.62$) and wind velocity ($r = 0.19$) whereas, [non-significant](#) and negative correlation with bright sunshine hours ($r = -0.14$).

3.2.2 The association between per cent fruit infestation and maggots count per fruit

The correlation association between [several](#) maggots per fruit and per cent fruit infestation during [the](#) *Summer* season in (Table 4) shows that the percent fruit infestation both on [a](#) number basis and weight basis shows [a](#) significant and positive correlation with [several](#) maggots per fruit $r = 0.682$ and $r = 0.905$ respectively.

More or less similar results were recorded by [the](#) following authors at different topographic regions. Laskar *et al.* (2013) noticed during warm *Rainy* months flies were more active as compared to dry and winter months in the foothills of [the](#) Himalayas. Similarly, Nair *et al.* (2020) concluded in their work


[that](#) maximum temperature and minimum temperature have [a](#) significant influence [on the Z.tau](#) population of cucurbits [in](#) Tripura ecosystem. Beer *et al.* (2021) concluded that the trapped *B.cucurbita* had [a](#) positive correlation with minimum temperature, maximum temperature and relative humidity during [the Kharif](#) season. Similarly, Bhowmik *et al.* (2014) recorded [the](#) highest fruit infestation i.e. 51.66 and 58.88 per cent on pointed gourd while on bitter gourd the highest infestation [was](#) 40.14 and 54.71 per cent during [the](#) 2012 and 2013 respective years.

3.2.3 Record of natural enemies:

A total of 50 pupae were maintained under observation, and their [color](#) gradually changed from light cream to dark brownish. Eight of the pupae turned out [to be](#) parasitoid. In the last week of October, it was discovered that 16% of the population was parasitized. This process was conducted twice to ensure the associated parasitoid. The emerged parasitoids were confirmed as *Psytalia fletcheri* (Silvestri) Fig.1 b It was a larval pupal parasitoid, it lay egg on the larvae of melon fly and emerged as [an](#) adult from pupae.



Fig.1 a) Fruit fly (Male and Female)



b) Parasitoid

Table 1: Percent fruit infestation of bitter gourd during both seasons Rainy 2021 and Summer 2022

Rainy Season 2021				Summer Season 2022			
SMW	Per cent fruit infestation		No. of Maggots/fruit	SMW	Per cent fruit infestation		No. of Maggots/Fruit
	Number basis	Weight basis			Number basis	Weight basis	
38	48.61*	46.15	16.67	21	28.99	25.84	6.53
39	54.41	54.66	21.00	22	36.81	26.45	8.33
40	56.12	65.00	21.74	23	61.14	53.68	14.73
41	59.33	52.16	13.04	24	38.66	40.77	17.67
42	42.53	45.99	10.26	25	55.12	55.93	20.47
43	37.67	41.24	8.12	26	49.00	54.68	19.40
44	33.16	34.33	5.30	27	51.04	61.86	25.67
45	23.55	22.34	7.00	28	53.53	56.76	27.40
46	15.65	12.41	4.49	29	53.86	61.93	27.07
Max.	59.33	65.00	21.74	Max.	61.14	61.93	27.40
Min.	15.65	12.41	4.49	Min.	28.99	25.84	6.53
Mean	41.23	41.59	11.96	Mean	47.57	48.66	18.59
STDV	14.24	15.44	6.17	STDV	9.84	13.36	7.23

Table 2: Correlation between percent fruit infestation by fruit flies and their maggots/fruit with weather parameters during the Rainyseason 2021.

Percent fruit infestation	Temperature(*C)		Relative Humidity (%)		R.F (mm)	Rainy days	Evaporation (%)	BSS (Hours)	WS (Km/hr)
	Max.	Min.	MRH	ERH					
Number Basis	0.21	0.61	0.87**	0.51	0.62	0.59	-0.49	-0.09	-0.59
Weight basis	0.21	0.55	0.84**	0.46	0.71*	0.63	-0.48	-0.02	-0.644
No. of maggots/fruit	-0.086	0.73*	0.92**	0.76*	0.89**	0.89**	-0.78*	-0.44	-0.302

* Significance at 5 %, ** Significance at 1 %

Table 3: Correlation between percent fruit infestation by fruit flies and their maggots/fruit with weather parameters during the Summerseason 2022.

Percent fruit infestation	Temperature(*C)		Relative Humidity (%)		Rain fall (mm)	Rainy days	Evaporation (%)	BSS (hrs/day)	Wind Speed (Km/hr)
	Max.	Min.	Morning	Evening					
No. Basis	-0.44	-0.37	0.52	0.25	0.12	0.13	-0.45	0.17	0.09
Weight basis	-0.65	-0.66*	0.77*	0.58	0.34	0.34	0.62	-0.05	0.11
No. of maggots/fruit	-0.73*	-0.77*	0.88**	0.83**	0.34	0.34	0.62	-0.14	0.19

* Significance at 5 %, ** Significance at 1 %

Table 4: Association between per cent fruit infestation and maggots of melon fruit fly in bitter gourd crop during both *Rainy* 2021 and *Summer* season 2022

<i>Rainy Season 2021</i>				<i>Summer Season 2022</i>			
SMW	% Fruit Infestation		Maggots/ Fruits	SMW	% Fruit Infestation		Maggots/ Fruits
	No.basis	Wt.basis			No.basis	Wt.basis	
38	48.61	46.15	16.67	21	28.99	25.84	6.53
39	54.41	54.66	21.00	22	36.81	26.45	8.33
40	56.12	65.00	21.74	23	61.14	53.68	14.73
41	59.33	52.16	13.04	24	38.66	40.77	17.67
42	42.53	45.99	10.26	25	55.12	55.93	20.47
43	37.67	41.24	8.12	26	49.00	54.68	19.40
44	33.16	34.33	5.30	27	51.04	61.86	25.67
45	23.55	22.34	7.00	28	53.53	56.76	27.40
46	15.65	12.41	4.49	29	53.86	61.93	27.07
Maggots	0.838*	0.857*		Maggots	0.682*	0.905**	

Summary and Conclusion:

Fruit fly was the major pest of bitter gourd in fruit damage. The fruit damage record range was 15.65 to 59.33 per cent in the rainy season whereas, 28.99 to 61.14 per cent in *summer* season. Per cent fruit infestation was positively correlated with relative humidity and rainfall of weather factors.

References:

- Allwood, A.J., Chinajariyawong, A., Kritsaneepaiboon, S., Drew, R.A.I., Hamacek, E.L., Hancock, D.L., Hengsawad, C., Jipanin, J.C., Jirasurat, M., Krong, C.K. and Leong, C.T.S., Host plant records for fruit flies (Diptera: Tephritidae) in Southeast Asia. *Raffles Bulletin of Zoology*, 47 (Supplement 7), 199; .1-92.
- Beer, D., Chandra, U., Yadav, J. K., & Kumar, S. Population dynamics of fruit flies in bitter gourd (*Momordica charantia* L.) and its relationship with weather factors. *The Pharma Innovation Journal* 2021; 10(10): 975-978
- Bhowmik, P., Devi, L. L., Chatterjee, M., & Mandal, D Seasonal Bionomics Of Melon Fruit Fly, *Bactrocera Cucurbitae* Coquillett On Bottle Gourd In Laboratory Condition. *The Eco Scan* 2014; 8(1&2): 157-162,
- Christenson, L.D. and Foote, R.H. Biology of fruit flies. *Ann. Rev. Entomol*; 1960; 5:171-192.
- Dhillon MK, Singh R, Naresh, J.S. and Sharma HC. The melon fruit fly, *Bactrocera cucurbitae*: A review of its biology and management. 16pp. 2005; *Journal of Insect Science* 5:40,

Laskar, N., & Chatterjee, H. Fruit Infestation and Larval Density of Melon Fly, *Bactrocera cucurbitae* (Coq.) as Influenced by Morphological Traits of Bitter Gourd (*Momordica charantia* L.). *International journal of Bio-resource and Stress Management*, 2013; 4(1), 54-57.

Nair, N., Pal, P., & Nath, D. Seasonal Incidence of Fruit Fly (*Zeugodacus tau*) in Cucurbit Ecosystem in Tripura. *Int J Curr Microbiol App Sci*, 2020; 9(11), 971-977.

White, I. M., and Elson-Harris, M.M. Fruit Flies of Economic Significance: Their Identification and Bionomics. Commonwealth Agriculture Bureau International, Oxon, UK, 1994; 595-774

UNDER PEER REVIEW